walking, moderate, and vigorous intensity, according to SLEDAI-2K and SDI were not found. In addition to high PCS of SF-36 ($p=0.006$) and SLEDAI-2K ($p=0.038$), less vigorous physical activity were related with lupus nephritis ($p=0.033$). However, the risk of CVD was not associated with physical activity of SLE.

Conclusions: This study showed that patients with lupus nephritis had less vigorous physical activity. It implicates that SLE-related organ damage might be associated with levels of physical activity.

REFERENCES:

- [1] Costenbader KH, Wright E, Liang MH, Karlson EW. Cardiac risk factor awareness and management in patients with systemic lupus erythematosus. *Arthritis Rheum* 2004;51:983–8.
- [2] Katz P, Julian L, Tonner MC, Yazdany J, Trupin L, Yelin E, Criswell LA. Physical activity, obesity, and cognitive impairment among women with systemic lupus erythematosus. *Arthritis Care Res (Hoboken)* 2012;64:502–10.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3585

AB0628 INFECTIONS IN HOSPITALISED PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS- A PROSPECTIVE OBSERVATIONAL STUDY FROM A TERTIARY CARE CENTRE IN SOUTHERN INDIA

S. Alur¹, M.M. Thabab¹, V.S. Negi², S. Sistla³. ¹Medicine; ²Clinical Immunology; ³Microbiology, Jawaharlal Institute of Post Graduate Medical Education and Research, Puducherry, India

Background: Infections are one of the main causes of morbidity and mortality in SLE.¹ The high prevalence of infection is attributed to use of immunosuppressive agents and disease activity. There is paucity of literature from India on infections, types of infections, predictors and outcome.

Objectives: To determine the frequency, types and determinants of major infections in hospitalised patients with SLE.

Methods: Patients with SLE (2012 SLICC-ACR criteria) aged ≥ 13 years, admitted to Clinical Immunology and Medicine wards were included. Clinical evaluation for major organ involvement of SLE, disease activity was assessed by SLEDAI-2K. Whenever there was a suspicion of infection appropriate work up to find the etiology of infections was done. Major infection was defined as presence of either one of the following a) sepsis b) pneumonia, or pyelonephritis, or endocarditis, or meningitis. c) skin or soft tissue infection requiring hospital admission for treatment d) Infection requiring IV antibiotic therapy. The diagnosis of infection was made by compatible clinical, microbiological and imaging criteria.

Results: From December 2016 till January 2018, 63 (55 women and 8 men) were included; Mean age was 28.8 ± 9.8 years; mean SLEDAI-2K was 14.6 ± 7.03 . Median duration of hospital stay was 14 days; range 2–64 days. Disease manifestation (including present and ever present) were arthralgia/arthritis in 51 (81%), oral ulcers 56 (88.9%), myositis 10 (15.9%), lupus nephritis 35 (55.6%), CNS lupus 14 (22.2%), serositis 14 (22.2%), cardiac involvement 13 (20.6%), APS 3 (4.8%). At the time of enrollment thrombocytopenia was present in 13 (20.6%), and leukopenia 11 (17.5%).

Twenty-three of 63 (36.5%) had infections (table 1), 14/23 (60.8) were microbiologically proven, remaining (39.2%) were confirmed by compatible clinical features and imaging.

Seven of 63 (11.1%) died; 5 (21.7%) in the infection group versus 2 (5%) in no infection group, $p=0.08$ OR 5.58 [CI 0.98–31.7]. Thirty-four of 63 (54%) had fever on admission; 17/23 (74%) patients with infection had fever compared to 17/40 (42.5%) in no infection group, $p=0.02$ [OR=3.8, CI 1.2–11.7]. There was no association of lupus nephritis, CNS lupus, leukopenia, receipt of cyclophosphamide, rituximab, methylprednisolone pulse therapy with occurrence of infection.

Abstract AB0628 – Table 1. Infections in SLE patients

Type of Infections	n=23 (%)
Bacterial Pneumonia	3 (13.7)
Fungal Pneumonia	2 (8.7)
(Invasive Aspergillosis)	
Pulmonary tuberculosis	2 (8.7)
Disseminated tuberculosis	2 (8.7)
Urinary tract infection	4 (17.4)
Skin and soft tissue infection	4 (17.4)
Blood stream infections	3 (13)
Suspected brain abscess	1 (4.3)
Pyogenic meningitis	1 (4.3)
Disseminated Varicella	1 (4.3)
Viral bronchiolitis	1 (4.3)

Conclusions: Almost 36% of SLE patients in hospital have infections. There appeared to be an increase in number of deaths among patients with infections. There was no association of infections with the dose of prednisolone or previous immunosuppression with cyclophosphamide.

REFERENCE:

- [1] Shyam C, Malaviya AN. Infection-related morbidity in systemic lupus erythematosus: a clinico-epidemiological study from northern India. *Rheumatol Int.* 1996; 16: 1–3.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1383

AB0629 PREVALENCE OF ASYMPTOMATIC VERTEBRAL FRACTURES IN POSTMENOPAUSAL WOMEN WITH SYSTEMIC LUPUS ERYTHEMATOSUS

S. Shkireeva^{1,2}, O.M. Lesnyak¹, E.G. Zotkin². ¹North-Western State Medical University named after I.I. Mechnikov, Saint-Petersburg; ²Research Institute of Rheumatology named after V.A. Nasonova, Moscow, Russian Federation

Objectives: To investigate the prevalence of vertebral fractures and to identify risk factors associated with vertebral fractures in postmenopausal women with systemic lupus erythematosus (SLE).

Methods: 86 consecutive postmenopausal women with SLE were included in this study. Bone mineral density (BMD) measurements of the hip and spine were performed using the dual energy X-ray absorptiometry (DEXA). Vertebral fracture assessment (VFA) was done for detection vertebral fractures using a method described by Genant. Accumulated damage was scored using the SLICC/ACR damage index (SDI).

Results: Vertebral fractures were defined in 42 (48.8%) postmenopausal women with SLE. Half of all patients with SLE ($n=22$, 25.6%) had asymptomatic vertebral fractures which were diagnosed only in this study. Every patient had 1.7 vertebral fractures on average (from 0 to 4). Distribution of vertebral fractures along the spine showed one peak at Th7. Univariate analyses of variables associated with fractures were longer duration of postmenopausal status in women with SLE, longer glucocorticoids intake, higher cumulative dose of glucocorticoids, higher SDI, lower BMD of the spine. Multivariate analysis showed longer duration of postmenopausal status in women with SLE (odds ratio 4.05, $p=0.007$), cumulative dose of glucocorticoids ≥ 60000 mg of prednisone (odds ratio 3.68, $p=0.009$), SDI ≥ 7 (odds ratio 2.94, $p=0.028$) and low BMD ($T \leq -2.0$) of the spine (odds ratio 2.48, $p=0.015$) were significantly associated with vertebral fractures in the thoracic and/or lumbar spine.

Conclusions: Vertebral fractures were occurred in 48.8% of postmenopausal women with SLE and half of them were asymptomatic. The current method using DEXA to predict the presence of vertebral fracture has limited value and there is a need for assessment of bone quality. Vertebral morphometry in patients with SLE is recommended and early therapeutic intervention is necessary to prevent vertebral fractures in postmenopausal women with SLE.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5498

AB0630 EXPRESSION OF TNF-A AND IL-6 IN SYSTEMIC LUPUS ERYTHEMATOSUS: RELATIONSHIP WITH DISEASE ACTIVITY

T.A. Panafidina, T.V. Popkova, A.A. Novikov, E.L. Nasonov. V.A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: Cytokines play an important role in the pathogenesis of SLE. TNF- α , IL-6 are the cytokines with suggested proinflammatory and immunoregulatory actions in the pathogenesis of SLE with differential effects on B or T cells, as well as on programmed cell death. Fas/APO-1 and Fas ligand (FasL) are involved in the apoptosis. FasL is a marker of apoptosis, it induces cell apoptosis when binds to Fas/APO-1. Both are expressed in membrane-associated and soluble forms (sFas/APO-1 and sFasL). sFas/APO-1 is known to inhibit the apoptosis via blocking the binding of Fas/APO-1 to FasL/sFasL.

Objectives: We evaluated the levels of IL-6 and TNF- α and their possible association with disease activity, apoptosis markers.

Methods: The study included 52pts (89% female, age 30.0 [26.5–44.5 years] (median [interquartile range 25% >75%])) with SLE ACR, 1997 and 20 controls (100% female without any rheumatic and infectious diseases, age 30.0 [25.0–39.5 years]). SLE-related factors, including disease duration, clinical features, SLE Disease Activity Index (SLEDAI 2K) and Systemic Lupus International Collaborating Clinics (SLICC) damage index were evaluated in parallel with relevant laboratory findings, autoantibodies. Serum levels of IL-6 and TNF- α (pg/ml), sFas/

APO-1 (pg/ml) and sFasL (ng/ml) were measured by ELISA (Bender MedSystem GmbH, Austria).

Results: Median SLE duration was 5 [1–11] years, SLEDAI 2K–9.5 [4.0–16.0], SLICC-0 [0–1], current prednisone dose–5.0 [0–11.9] mg/day, 44% of pts received hydroxychloroquine, 17% >cyclophosphamide, 10% >azathioprine, 4% >mycophenolate mofetil, 8% >biologics. SLEpts had higher level of IL-6 vs control (2.9 [1.5–9.7] vs 1.4 [0.7–2.1] pg/ml, $p<0.01$) and lower level of TNF- α (1.9 [0.5–3.2] vs 4.5 [3.5–6.4] pg/ml, $p<0.0001$), lower levels of sFas/APO-1 vs control (657.5 [483.2–791.4] vs 1456.9 [426.5–2060.8] pg/ml, $p>0.05$) and lower levels of sFasL vs control (0.01 [0.01–0.03] vs 0.07 [0.05–0.10] ng/ml, $p<0.0001$). In SLE pts serum level IL-6 correlated with IgM ($r=0.69$, $p<0.01$), K⁺ ($r=0.43$, $p<0.01$), TNF- α ($r=0.50$, $p<0.001$), APRIL ($r=0.39$, $p<0.01$) and sFas/APO-1 levels ($r=0.29$, $p<0.05$); TNF- α level correlated with SLICC ($r=0.32$, $p<0.05$), APRIL ($r=0.42$, $p<0.01$), IL-6 ($r=0.50$, $p<0.001$), haemoglobin concentration ($r=0.39$, $p<0.01$), sFasL ($r=0.44$, $p<0.01$) and total cyclophosphamide dose ($r=0.28$, $p<0.05$). We divided SLEpts on two groups: the 1st-pts ($n=31$) with high activity (SLEDAI 2K ≥ 8), the 2nd-pts ($n=21$) with low (SLEDAI 2K < 8). The pts in the 1st group had lower level of TNF- α compared to 2nd (1.6 [0.3–2.2] vs 2.8 [1.3–4.5] pg/ml, $p<0.05$), with no difference in IL-6 level.

Conclusions: SLEpts demonstrated higher IL-6 and lower TNF- α levels as compared to healthy controls. Pts with high activity of SLE had decreased level of TNF- α . There is no correlation between disease activity and IL-6 level, as well as between elevated IL-6 and acute phase indicators. We suggest an immunoregulatory not proinflammatory role of IL-6 in SLE and probable protective role of TNF- α in SLE pathogenesis (it's deficiency can lead to the development and high activity of SLE). There was no correlation between IL-6, TNF- α levels and any SLE features. The correlation of IL-6 with sFas/APO-1 level (apoptosis inhibitor that blocks binding of sFas/APO-1 to sFasL) and TNF- α with sFasL concentration (a marker of apoptosis) suggests their different roles in the mechanisms of apoptosis.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3220

AB0631 IS HYPERHOMOCYSTEINEMIA (HHC) ADDITIONAL RISK FACTORS OF THROMBOSES IN PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS AND ANTIPHOSPHOLIPID SYNDROME

T.M. Reshetnyak^{1,2}, N.V. Seredavkina¹, E.L. Nasonov¹, L.I. Patrushev³. ¹Vascular Rheumatology, V.A. Nasonova Research Institute of Rheumatology; ²Russian Medical Academy of Postgraduate Education; ³Shemyakin-Ovchinnikov Institute of Biorganic Chemistry of RAS, Moscow, Russian Federation

Background: HHC may be additional factor of thrombosis in SLE patients

Objectives: To assess the role of HHC as additional risk factors in development of vascular complications in SLE and antiphospholipid syndrome (APS).

Methods: A total of 125 patients (24 M and 101 F, mean age 38 $\pm$ 13 years and the disease duration 14 $\pm$ 11 years) were divided into three groups: group 1 – SLE patients ($n=51$); group 2 – SLE +APS patients ($n=49$); group 3 – primary APS patients ($n=25$). Homocystein (HC) was assayed by high performance liquid chromatography. DNA diagnostics by the method of polymerase chain reaction was used to determine gene mutation C677T methylenetetrahydrofolate (MTHFR) reductase in 93 out of 125 patients.

Results: HHC (HC >15 mcg/L) was diagnosed in 82 of 125 (66%) patients: in 59% patients of group 1, 67% – of group 2 and 76% – of group 3. There was a relationship between HHC and digital necrosis (DN): 80% of patients with DN had HHC while HHC was in 57% patients without DN ($\chi^2=4.76$, $p=0.03$). Elevated level of HC was registered in 43 of 55 (78%) APS patients with thromboses vs. 9 of 19 (47%) aPL-positive SLE patients without thromboses ($p=0.03$). HHC occurred significantly more frequently in patients with arterial thromboses (in all 14 patients) than in patients with venous thromboses – in 16 of 23 (69.9%) pts ($p=0.03$) and in the absence of thromboses ($p=0.04$). HHC was associated with thromboses of cerebral (in 90%), peripheral arteries (in 84%) and myocardial infarction (in 79%) vs. 47% of patients without thromboses ($p=0.005$; $p=0.04$, $p=0.04$ respectively), (79 vs. 47%). Mutation of C677T MTHFR was revealed in 33.3% of SLE patients, in 57.7% – of SLE +APS and in 63.2% – PAPS. Mutation C677T in the gene MTHFR was detected in 57% of examined patients, 20% of them had homozygous variant and 80% are heterozygous. Patients with a homozygous C677T MTHFR gene was observed the tendency to develop arterial thrombosis

Conclusions: More than 50% APS pts had elevated level of HC as additional thromboses factors. Correlation between HHC and thromboses, primarily arterial, in APS patients gives grounds for the role of HHC in development of vascular complications in SLE and APS pts.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2484

AB0632 ESSDAI DOMAIN EVALUATION OF PRIMARY SJÖGREN'S SYNDROME (PSS) PATIENTS ENROLLED IN TWO INDEPENDENT POC STUDIES INDICATES DIFFERENTIAL UTILITY OF DOMAINS FOR TRIAL INCLUSION AND COMPOSITE ENDPOINTS IN PSS TRIALS

T. Dörner¹, B.A. Fisher², X. Ren³, P. Faerber³, P. Gergely³, L. Mooney³, Y. Li³, S. J. Oliver³, W. Hueber⁴, A.M. Wright³. ¹Charité Hospital, Berlin, Germany; ²Rheumatology Research Group, University of Birmingham, Birmingham, UK; ³Novartis Institutes for BioMedical Research; ⁴Novartis Pharma, Basle, Switzerland

Background: According to international consensus, disease activity in primary Sjögren's syndrome (pSS) patients shall be scored across 12 different domains according to the European Sjögren Syndrome Disease Activity Index (ESSDAI¹). Two separate phase 2 proof-of-concept (PoC) studies using either BAFF receptor inhibitor (VAY736; iganalumab) or CD40 inhibitor (CFZ533) have recently been completed in comparable pSS cohorts with ESSDAI as the primary endpoint.

Objectives: To evaluate contribution of individual domains to composite ESSDAI scores at baseline and after interventional treatment with either VAY736 or CFZ533, or placebo. Aggregate efficacy and safety results were presented at previous meetings.^{2,3}

Methods: Key inclusion criteria for both studies were fulfilling revised European US consensus criteria for pSS,⁴ autoantibody positivity, exclusion of secondary SS and ESSDAI ≥ 6 . Permitted background medications in both trials included stable doses of hydroxychloroquine, methotrexate and prednisone ≤ 10 mg/d and also azathioprine in the CFZ533 trial. The primary endpoint was change in ESSDAI at week 12. Descriptive and principal component analysis were done with the goal to identify distinct subgroups of patients based on ESSDAI involvement at baseline (BL), and the relative importance of single domain contribution to overall ESSDAI responses were explored.

Results: 27 patients received single i.v. dose VAY736 10 mg/kg ($n=12$) or 3 mg/kg ($n=6$) or placebo, and 44 patients received multiple doses of CFZ533 10 mg/kg i.v. ($n=21$) or 3 mg/kg s.c. ($n=8$) or placebo. ESSDAI breakdown at BL showed a predominance of the articular, glandular, biological, constitutional and lymphadenopathy domains in both trials. Activity in more than 3 domains was recorded for 12/27 (44%) and 12/44 (27%) of patients in the VAY736 and CFZ533 studies, respectively. Principal component analysis identified the articular domain as the key component describing the difference between patients in their ESSDAI domains at BL. Two other domains that explained the variability between patients were the biological and glandular domains. Treatment effects in domains with low BL scores were more difficult to assess using ESSDAI. The majority of ESSDAI domains were not amenable to quantitative analysis, due to absence or low incidence at baseline.

Conclusions: The most frequently observed ESSDAI domain was articular involvement. Our results provide insights into ESSDAI domain frequency and distribution in the randomised controlled trial setting that may have implication for future trial design in pSS.

REFERENCE:

- [1] Seror R, et al. (2016) Ann Rheum Dis 75:382–389.
- [2] Dörner Th, et al. (2016) Arthritis Rheumatol 68 Suppl 10;
- [3] Fisher B, et al. (2017) Arthritis Rheumatol 69; Suppl 10;
- [4] Vitali C, et al. (2002) Ann Rheum Dis 61:554–558.

Disclosure of Interest: T. Dörner Grant/research support from: Novartis, Charité CRO, Consultant for: Novartis, B. Fisher Consultant for: Novartis, Roche, BMS and AstraZeneca/MedImmune., X. Ren Employee of: Novartis Pharmaceuticals, P. Faerber Employee of: Novartis Pharmaceuticals, P. Gergely Shareholder of: Novartis Pharmaceuticals, Employee of: Novartis Pharmaceuticals, L. Mooney Employee of: Novartis Pharmaceuticals, Y. Li Shareholder of: Novartis Pharmaceuticals, Employee of: Novartis Pharmaceuticals, S. J. Oliver Shareholder of: Novartis Pharmaceuticals, Employee of: Novartis Pharmaceuticals, W. Hueber Shareholder of: Novartis Pharmaceuticals, Employee of: Novartis Pharmaceuticals, A. M. Wright Shareholder of: Novartis Pharmaceuticals, Employee of: Novartis Pharmaceuticals

DOI: 10.1136/annrheumdis-2018-eular.4441

AB0633

MEASURING WHAT MATTERS TO LUPUS PATIENTS: TRANSLATING PATIENT VIEWS INTO NOVEL PATIENT-REPORTED OUTCOMES

S. Cleanthous¹, A. Regnault², M. Schneider³, C. Gordon⁴, S. Bartlett⁵, S. Cano¹, T. Morel⁶. ¹Modus Outcomes, Cambridge, UK; ²Modus Outcomes, Lyon, France; ³Policlinic and Hiller Research Unit of Rheumatology, UKD, Heinrich-Heine-University, Düsseldorf, Germany; ⁴University of Birmingham, Birmingham, UK; ⁵McGill University, Montreal, Canada; ⁶UCB Pharma, Brussels, Belgium

Background: The 2010 U.S. FDA guidance¹ on systemic lupus erythematosus (SLE) acknowledged that 'improvements in clinical outcome measures in patients with SLE may not always translate to improvements in how patients feel or function' and encourages the use of patient-reported outcome instruments (PROs) to assess changes in outcomes that matter most to patients. Fatigue, identified by patients as their chief complaint during the Lupus Patient-Focused Drug Development (PFDD) meeting², is not optimally measured by existing PROs¹. There is a need for new patient-centred outcome measures as they are pivotal to PFDD. To address this, UCB has been conducting a multi-faceted, patient-centric, mixed methods research programme including evidence generated from >2100 SLE patients.

Objectives: There were three main objectives of the programme: 1) identify what concepts are most important to measure for SLE patients when evaluating treatment benefit; 2) critically evaluate the extent to which these concepts are captured by existing PROs; and 3) explore the opportunity to develop and evaluate new conceptually-based PROs in SLE.

Methods: There were three stages: 1) development of a preliminary treatment benefit conceptual model related to SLE including disease symptoms and impacts (generated from literature review, and patient and clinician input); 2) formal comparison of i) the conceptual model and the content validity (qualitative review) of existing SLE PROs and ii) psychometric properties (ie. SF-36, FACIT-F, and LupusQoL in the discontinued EMBODY Phase 3 studies for epratuzumab); 3) development of new conceptually-based PROs in SLE (figure 1).

Abstract AB0633 – Figure 1

Results: A preliminary treatment benefit conceptual model in SLE was developed. Among the many symptoms and disease impacts reported by SLE patients, physical fatigue, mental fatigue ('brain fog'), sudden fatigability, joint/muscle stiffness and pain, skin symptoms and mobility difficulties featured as most troublesome. The content validity of 10 widely used PROs was found to be limited when compared to this conceptual model. Psychometric analysis (based on Rasch Measurement Theory) confirmed the limitations of the SLE PROs used in the EMBODY studies and suggested post-hoc reconceptualisation would improve the ability of the scales to detect clinical change. This led to the development of five new conceptually-based SLE PROs (ie. fatigue, pain, mobility, symptom severity, and emotional state), which are currently being explored in >500 SLE patients in a Phase 2b dapirolizumab pegol study (NCT02804763) and two cross-sectional, non-interventional, observational studies. Data on the new PROs are currently being assessed at several sites in the USA, Latin America, UK and Germany and will be available in the coming months.

Conclusions: New, well-defined and reliable PROs to better capture the patient perceptions of the symptoms and impact of SLE are needed. UCB has developed

five new PROs which offer the promise of improved patient-centred outcome measurements in SLE and other autoimmune diseases.

REFERENCES:

- [1] Food and Drug Administration. Guidance for Industry. Systemic Lupus Erythematosus. June 2010.
- [2] Lupus Patient-Focused Drug Development meeting, 25 September 2017.

Disclosure of Interest: S. Cleanthous Employee of: Modus Outcomes Ltd, A. Regnault Employee of: Modus Outcomes Ltd, M. Schneider Consultant for: UCB Pharma, C. Gordon Grant/research support from: UCB Pharma, Consultant for: UCB, Merck Serono, GSK, S. Bartlett: None declared, S. Cano Shareholder of: Modus Outcomes Ltd, Employee of: Modus Outcomes Ltd, T. Morel Shareholder of: UCB Pharma, Employee of: UCB Pharma

DOI: 10.1136/annrheumdis-2018-eular.5435

AB0634

CHARACTERISATION OF SLE PATIENTS WHO UNDERWENT CARDIOVASCULAR SURGERY

T. Kawamoto¹, S. Matsushita², H. Amano¹, K. Minowa¹, Y. Asai¹, K. Yamaji¹, A. Amano², N. Tamura¹. ¹Internal medicine and rheumatology; ²Cardiovascular Surgery, Juntendo University School of Medicine, Tokyo, Japan

Background: SLE is characterised by alternate occurrences of flares and remissions throughout its course. During these flares, immune complexes are excessively produced and deposited in affected tissues. Previous reports suggest that immune complexes deposited in the blood vessels, atrioventricular valves, and other associated components of the heart cause disorders in these tissues. Although, the characteristics of patients in whom tissue injuries progress and ultimately require surgery have not been clarified.¹

Objectives: Heart diseases are categorised into valvular disease, ischaemic heart disease, and aortic aneurysm. This study was conducted to determine whether there was a correlation between the activity of SLE and the progress and aggravation of heart disease in each category.

Methods: Of the 2707 patients who were diagnosed with SLE at Juntendo University Hospital from 2012–2017, 35 underwent cardiac surgery. Six patients met the exclusion criteria, and thus, 29 patients were included. Patients with infectious endocarditis and those in whom SLE had not been diagnosed prior to surgery were excluded. Indices for SLE activity were anti-DNA Ab and CH50. Peak and bottom values of these indices before surgery were recorded, and the values obtained during the pre-surgery 180 day period were compared with those that had been determined prior to this period.

Results: The mean disease duration at the time of cardiac surgery was 24.2 years. The cardiac diseases were heart valve disease (V) in 10, ischaemic heart disease (I) in 7, and aortic aneurysm (A) in 12 patients. The integrated values obtained by the activity indices, anti-DNA Ab, and disease duration were 14 020 for V, 32 966 for I, and 29 444 for A. Thus, there was a tendency for values to be slightly lower in the heart valve disease category. The peak/bottom values for CH50 and anti-DNA Ab in the pre-surgery 180 day period were no different from those obtained prior to this period. Investigation of anti-phospholipid antibodies indicated that the rates of positive result for either CL β_2 GPI, anti-CL antibodies, or LA were 60% for V, 28% for I, and 33% for A, and thus, higher in the heart valve disease category. All patients were treated with steroids, and five patients (17.2%) of the 29 used immunosuppressants. This rate tended to be lower than the previous frequency of the combined use of immunosuppressive drugs at our hospital (onset: 18.4%, first flare 33.3%).²

Conclusions: There is a possibility that the long-term duration of SLE may lead to cardiac disease requiring surgery. The integrated value for DNA Ab was lower in the heart valve disease category; therefore, the presence of anti-phospholipid antibodies may be a risk factor for valvular disease. The low frequency of combined use of immunosuppressants in SLE patients with cardiovascular manifestations requiring surgery, suggest the possibility that treatment with a single steroid may require heart surgery.

REFERENCES:

- [1] Yasuhiro Sakaguti, Sachio Kawai, Ryoza Okada, et al. Immunohistochemical study of the endomyocardial biopsy of Systemic Lupus Erythematosus. Journal of Cardiology. 1995;25:181–188
- [2] Kentaro Minowa, Hirofumi Amano, Yoshinari Takasaki, et al. Disease flare patterns and predictors of systemic lupus erythematosus in a monocentric cohort of 423 Japanese patients during long-term follow-up: The JUDE study. Mod Rheumatol. 2017;27:72–76

Acknowledgements: none

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5041