musculoskeletal pain (n=76; RR, 1.37; IC, 0.45) and myasthenia gravis (n=66; RR, 1.42; IC, 0.50);² PEMBROLIZUMAB: arthralgia (n=151; RR, 1.43; IC, 0.52) and pain in an extremity (n=58; RR, 1.35; IC, 0.43);³ DURVALUMAB: polymyositis (n=2; RR, 4.41; IC, 2.15), rhabdomyolysis (n=4; RR, 2.68; IC, 1.42), and autoimmune arthritis (n=2; RR, 8.83; IC, 3.14);⁴ IPILIMUMAB: muscular weakness (n=157; RR, 1.70; IC, 0.76) and back pain (n=105; RR, 1.27; IC, 0.34). In general, rates of rheumatic and musculoskeletal adverse events were higher in men and in the elderly population (>65 years).

Conclusions: A wide spectrum of rheumatic and musculoskeletal toxicity signals were detected with ICI's. Clinicians need to be vigilant about these rare but debilitating complications. Future studies to explore mechanisms and optimal management strategies of these toxicities are warranted.

REFERENCE:

- [1] Sarangdhar M, Tabar S, Schmidt C, et al. Data mining differential clinical outcomes associated with drug regimens using adverse event reporting data. *Nat Biotechnol* 2016;34:697–700.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6113

THURSDAY, 14 JUNE 2018

Can we halt progression of structural damage in axial SpA?

OP0198 COMBINED EFFECTS OF TUMOUR NECROSIS FACTOR INHIBITORS AND NSAIDS ON RADIOGRAPHIC PROGRESSION IN ANKYLOSING SPONDYLITIS

L.S. Gensler¹, M. Gianfrancesco¹, M.H. Weisman², M.A. Brown³, M. Lee⁴, T. Leach⁵, M. Rahbar⁴, J.D. Reveille⁶, M.M. Ward⁷. ¹Medicine, University of California, San Francisco, San Francisco; ²Medicine, Cedars Sinai Medical Center, Los Angeles, USA; ³Medicine, Queensland University of Technology, Brisbane, Australia; ⁴Epidemiology/Biostatistics, The University of Texas Health Science Center at Houston, Houston; ⁵Radiology, Cedars Sinai Medical Center, Los Angeles; ⁶Medicine, The University of Texas Health Science Center at Houston, Houston; ⁷NIH/NIAMS, Bethesda, USA

Background: The potential of TNFi or NSAIDs to reduce radiographic progression in AS is uncertain and causal effects of both exposures on radiographic progression have not been convincingly demonstrated. In addition, no study has evaluated whether effects are comparable among different NSAIDs in this setting. **Objectives:** The objective of this study was to explore causal effects of NSAIDs and TNFi on radiographic progression in Ankylosing Spondylitis (AS) and to compare effects of celecoxib to other NSAIDs.

Methods: We included all patients meeting the modified New York criteria in a prospective cohort with at least 4 years of clinical and radiographic follow up. Clinical and medication data were collected every 6 months and radiographs were performed at baseline and every 2 years. We used longitudinal targeted maximum likelihood estimation to estimate the causal effect of TNFi and NSAIDs (using the NSAID index) on radiographic progression as measured by the modified Stoke Ankylosing Spondylitis Spine Score (mSASSS) at 2 and 4 years, accounting for time-varying covariates. We controlled for sex, race/ethnicity, education, symptom duration, enrollment year, number of years on TNFi, symptom duration at time of TNFi start, baseline mSASSS, ASDAS-CRP, current smoking, and missed visit status.

Abstract OP0198 – Table 1

	TNF use	No TNF use	Mean Difference	P-value
Comparing TNF use vs no TNF use, given no NSAID (=0) at time t:				
mSASSS @ 2 years	13.94	14.92	-0.98 (-2.77, 0.81)	0.28
mSASSS @ 4 years	16.12	15.62	0.50 (-0.63, 1.64)	0.38
Comparing TNF use vs no TNF use, given low NSAID (>0 and <50) at time t:				
mSASSS @ 2 years	15.43	15.49	-0.06 (-1.63, 1.51)	0.94
mSASSS @ 4 years	15.52	16.76	-1.24 (-1.80, -0.68)	<0.001
Comparing TNF use vs no TNF use, given high NSAID (>=50) at time t:				
mSASSS @ 2 years	14.79	15.13	-0.34 (-1.46, 0.78)	0.56
mSASSS @ 4 years	14.17	17.47	-3.31 (-4.02, -2.59)	<0.001
Comparing TNF use vs no TNF use, NSAID=celecoxib at time t:				
mSASSS @ 2 years	11.63	15.62	-3.98 (-4.51, -3.45)	<0.001
mSASSS @ 4 years	14.37	19.06	-4.69 (-5.08, -4.30)	<0.001

Results: Of the 519 patients, 75% were male with a baseline mean (SD) age and symptom duration of 41.4 (13.2) and 16.8 (12.5) years respectively. The baseline mean (SD) mSASSS was 14.2 (19.6). At baseline, NSAIDs were used in 66% of

patients, of which ½ used an index <50 and ½ an index ≥50). TNFi were used in 46% of patients at baseline. In the setting of TNFi use, the addition of NSAID therapy was associated with less radiographic progression in a dose-related manner at 4 years. When NSAID specific effects were examined, celecoxib in combination with TNFi use was associated with the greatest reduction in radiographic progression and this was significant at both 2 and 4 years (table 1).

Conclusions: Dose related use of NSAIDs together with TNFi in AS patients has a synergistic effect in slowing radiographic progression with the greatest effect in those using both high-dose NSAIDs and TNFi. Celecoxib appears to confer the greatest benefit in decreasing progression with effect at both 2 and 4 years.

Disclosure of Interest: L. Gensler Grant/research support from: Amgen, AbbVie, UCB, Consultant for: Janssen, Lilly, Novartis, M. Gianfrancesco: None declared, M. Weisman Consultant for: Celltrion, Bayly, Novartis, Lilly, GSK, M. Brown Grant/research support from: AbbVie, Janssen, UCB, Leo Pharma, Consultant for: AbbVie, Janssen, Pfizer, Speakers bureau: AbbVie, UCB, Pfizer, M. Lee: None declared, T. Leach: None declared, M. Rahbar: None declared, J. Reveille Grant/research support from: Lilly UCB, Consultant for: Novartis Janssen Lilly UCB, M. Ward: None declared

DOI: 10.1136/annrheumdis-2018-eular.4027

OP0199 SUSTAINED REMISSION OF INFLAMMATION IS ASSOCIATED WITH REDUCED STRUCTURAL DAMAGE ON SACROILIAC JOINT MAGNETIC RESONANCE IMAGING IN PATIENTS WITH EARLY AXIAL SPONDYLOARTHRITIS: EVIDENCE TO SUPPORT THE CONCEPT OF TREAT-TO-TARGET

W.P. Maksymowych¹, P. Claudepierre², M. de Hooze³, R.G. Lambert¹, R. Landewé⁴, A. Molto⁵, D. van der Heijde³, J.F. Bukowski⁶, H. Jones⁶, I. Logeart⁷, L. Marshall⁶, R. Pedersen⁶, A. Szumski⁸, B. Vlahos⁶, M. Dougados⁵. ¹University of Alberta, Edmonton, Canada; ²Université Paris Est Creteil, Paris, France; ³Leiden University Medical Center, Leiden; ⁴Amsterdam Rheumatology and Immunology Center, Amsterdam, Netherlands; ⁵Paris Descartes University, Hôpital Cochin, Paris, France; ⁶Pfizer, Collegeville, USA; ⁷Pfizer France, Paris, France; ⁸inVentiv Health, Princeton, USA

Background: Treat-to-target is acceptable in RA; however, it is unknown whether it will reduce/prevent disability, impairment of mobility, and structural damage in early axial spondyloarthritis (axSpA) without radiographic sacroiliitis.

Objectives: To evaluate the impact of sustained clinical remission on MRI structural parameters. We hypothesised that patients with sustained inactive disease according to the ankylosing spondylitis disease activity score (ASDAS <1.3) are more likely to achieve reduction in erosion (structural damage) and increase in backfill (a reparative process) on MRI of the SI joints (SIJ).

Methods: EMBARK (NCT01258738) and DESIR (NCT01648907) enrolled patients with early axSpA. EMBARK included 12 weeks of double-blind placebo-control, then open-label etanercept for 92 weeks. Patients in the DESIR observational cohort had no history of biologics and received no biologics for 2 years. T1 weighted MRI images of SIJ at baseline and 104 weeks were combined and anonymized; readers were unaware of film chronology and original patient cohort. Three experienced readers evaluated MRI images using the SpondyloArthritis Research Consortium of Canada SIJ Structural Score. Lesion change was considered present if ≥2 of 3 readers measured change in same direction. ASDAS endpoints were assessed sequentially: sustained (≥2 visits 6 months apart) inactive disease (ASDAS <1.3) or moderate disease activity (≥1.3 to <2.1); or no sustained response (≥2.1). Net proportions of patients with decrease in erosion and increase in backfill were determined, unadjusted and adjusted for covariates that may affect development of lesions on MRI.

Results: From EMBARK and DESIR, 161 and 76 patients, respectively, were included. For patients in EMBARK with sustained ASDAS <1.3, a greater percentage had decrease in erosion (34/104, 32.7%) than increase (5/104, 4.8%); p<0.0001; without sustained response, 5/24 (20.8%) had decrease in erosion and 1/24 (4.2%) had increase. This trend was also present in DESIR. Patients with sustained ASDAS <1.3 in EMBARK: 22.1% had increase in backfill, 0% had decrease; p<0.0001; in DESIR, 21.7% had increase, 0% had decrease; p<0.05. For those without sustained response, difference between increase and decrease was smaller. Net percent of patients (adjusted) with sustained ASDAS <1.3 and erosion decrease: 22.6% and 9.3% for EMBARK and DESIR, respectively; without sustained response: 13.3% and -10.1%. Net percent of patients with sustained ASDAS <1.3 and backfill increase: 19.6% and 25.7% for EMBARK and DESIR, respectively; without sustained response: 8.7% and 6.0%.

Conclusions: These data demonstrate a link between sustained ASDAS inactive disease and MRI structural endpoints. Clinical relevance of change in MRI SIJ erosion and backfill and their relationship to ankylosis development requires study.

Disclosure of Interest: W. Maksymowych Grant/research support from: AbbVie, Pfizer, Consultant for: AbbVie, Janssen, Lilly, Merck, Novartis, Pfizer, UCB, P. Claudepierre Consultant for: AbbVie, BMS, Celgene, Janssen, Novartis, Merck,

Pfizer, Roche, UCB, Lilly, M. de Hooge Employee of: Selfemployed without other personnel (registered company under the Belgium law) MdH Research. Additional Affiliation: Ghent University, Ghent, Belgium., R. Lambert Consultant for: AbbVie, Bioclinica, Janssen, Parexel, UCB, R. Landewé Grant/research support from: Abbott, Amgen, Centocor, Novartis, Pfizer, Roche, Schering-Plough, UCB, Consultant for: AbbVie, Ablynx, Amgen, Astra-Zeneca, Bristol Myers Squibb, Celgene, Janssen, Galapagos, GlaxoSmithKline, Novartis, Novo-Nordisk, Merck, Pfizer, Roche, Schering-Plough, TiGenix, UCB, Employee of: Director of Rheumatology Consultancy BV, which is a registered company under Dutch law, Speakers bureau: Abbott/AbbVie, Amgen, Bristol Myers Squibb, Janssen, Merck, Pfizer, Roche, Schering-Plough, UCB, A. Molto Grant/research support from: Pfizer, Consultant for: Merck/UCB, D. van der Heijde Consultant for: AbbVie, Amgen, Astellas, AstraZeneca, BMS, Boehringer Ingelheim, Celgene, Daiichi, Lilly, Galapagos, Gilead, GlaxoSmithKline, Janssen, Merck, Novartis, Pfizer, Regeneron, Roche, Sanofi, Takeda, UCB, Employee of: Director of Imaging Rheumatology bv., J. Bukowski Shareholder of: Pfizer, Employee of: Pfizer, H. Jones Shareholder of: Pfizer, Employee of: Pfizer, I. Logeart Shareholder of: Pfizer, Employee of: Pfizer, L. Marshall Shareholder of: Pfizer, Employee of: Pfizer, R. Pedersen Shareholder of: Pfizer, Employee of: Pfizer, A. Szumski Employee of: InVentiv Health, B. Vlahos Shareholder of: Pfizer, Employee of: Pfizer, M. Dougados Grant/research support from: Pfizer, AbbVie, UCB, Merck, Lilly, Consultant for: Pfizer, AbbVie, UCB, Merck, Lilly
DOI: 10.1136/annrheumdis-2018-eular.1114

THURSDAY, 14 JUNE 2018

Reproductive issues in rheumatology

OP0200

CHARACTERISTICS AND OUTCOMES OF PROSPECTIVELY REPORTED PREGNANCIES EXPOSED TO CERTOLIZUMAB PEGOL FROM A SAFETY DATABASE

M.E.B. Clowse¹, A.E. Scheuerle², C. Chambers³, A. Afzali⁴, A. Kimball⁵, J. J. Cush⁶, M. Cooney⁷, L. Shaughnessy⁷, M. Vanderkelen⁸, F. Förger⁹. ¹Duke University Medical Center, Durham, NC; ²UT Southwestern Medical Center, Dallas, TX; ³UC San Diego School of Medicine, La Jolla, CA; ⁴Ohio State University, Wexner Medical Center, Columbus, OH; ⁵Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA; ⁶Baylor Scott and White Research Institute, Dallas, TX; ⁷UCB Pharma, Raleigh, NC, USA; ⁸UCB Pharma, Braine-l'Alleud, Belgium; ⁹Inselspital, University of Bern, Bern, Switzerland

Background: Anti-tumour necrosis factor medications (anti-TNFs) are effective in controlling chronic inflammatory diseases, but information about their use and safety in pregnancy is limited. Consequently, anti-TNFs are often discontinued early in gestation. Certolizumab pegol (CZP), an Fc-free, PEGylated anti-TNF approved for treatment of rheumatic diseases and/or Crohn's disease, has no active placental transfer.

[a] Includes 10 sets of twins.

Abstract OP0200 – Figure 1 Overview of pregnancy reports

Objectives: This project provides information on pregnancy outcomes in women receiving CZP, especially those with early pregnancy exposure.

Methods: Prospective and retrospective data on maternal CZP exposure, including timing and duration, outcomes, comorbidities, and major malformations were extracted from the UCB Pharma safety database through 6 March 2017. This analysis was limited to prospective reports to avoid bias associated with retrospective submissions. Numbers of live births, miscarriages, elective abortions, stillbirths, and major congenital malformations were ascertained.

Results: From a total of 1541 maternal CZP-exposed pregnancies, 1137 were reported prospectively, of which 528 pregnancies (including 10 twin pregnancies) had 538 known outcomes: 459 live births (85%), 47 miscarriages (9%), 27 elective abortions (5%), and 5 stillbirths (1%) (figure 1). Of the 459 live births, 8 (2%) cases of major congenital malformations were reported (vesicoureteral reflux, club foot, congenital heart disease, cerebral ventricle dilatation, polydactyly, anal fistula, accessory auricle, and hydronephrosis). Of the 528 prospective pregnancies with known outcomes, 436 (83%) were exposed during the 1 st trimester, when most organogenesis occurs; 201 pregnancies were exposed during the entire pregnancy.

Conclusions: This analysis represents the largest published cohort of pregnant women exposed to an anti-TNF for management of chronic inflammatory diseases. Analysis of pregnancy outcomes does not indicate a malformative effect of CZP compared to the EU general population (2%–3%), nor an increased risk of foetal death. These data are reassuring for women of childbearing age considering treatment with CZP; however, the ongoing collection of post-marketing surveillance data, including the ongoing MotherToBaby study from the Organisation of Teratology Information Specialists, will provide further important information.

Acknowledgements: This study was funded by UCB Pharma. The authors thank the patients and their caregivers in addition to the investigators and their teams who contributed to this study. Editorial services were provided by Costello Medical. All costs associated with development of this abstract were funded by UCB Pharma.

Disclosure of Interest: M. Clowse Grant/research support from: Pfizer and Janssen, Consultant for: UCB Pharma, A. Scheuerle Grant/research support from: UCB Pharma, INC Research and Genentech, C. Chambers Grant/research support from: AbbVie, Amgen Inc., Bristol-Myers Squibb, Celgene, GlaxoSmithKline, Janssen, Pfizer, Hoffman-La Roche, Genentech, Genzyme Sanofi-Aventis, Seqirus, Takeda and UCB Pharma, A. Afzali Grant/research support from: UCB Pharma, Consultant for: UCB Pharma, AbbVie, Takeda and IBD Horizons, A. Kimball Consultant for: UCB Pharma, Dermira, Janssen and AbbVie, J. Cush Grant/research support from: Pfizer, Janssen, AbbVie, Celgene, Novartis, AstraZeneca and Genentech, Consultant for: Janssen, AbbVie, Novartis, Amgen, Genentech, Lilly and Horizon, M. Cooney Employee of: UCB Pharma, L. Shaughnessy Employee of: UCB Pharma, M. Vanderkelen Employee of: UCB Pharma, F. Förger Grant/research support from: UCB Pharma, Speakers bureau: Mepha, Roche and UCB Pharma

DOI: 10.1136/annrheumdis-2018-eular.2417

THURSDAY, 14 JUNE 2018

Do we still need biopsies to diagnose Sjögren's and autoimmune myositis?

OP0201

CLINICAL OUTCOMES AND RESPONSE TO ANTI-THROMBOTIC TREATMENT AMONG PATIENTS WITH CONCOMITANT LUPUS NEPHRITIS AND THROMBOTIC MICROANGIOPATHY: A MULTICENTER COHORT STUDY

S. Sciascia¹, J. Yazdany², M. Cuadrado³, M. Radin⁴, M. Dall'Era⁵, I. Aggarwal², R. Fenoglio¹, A. Barreca⁶, M. Papotti⁶, I. Cecchi¹, E. Rubini¹, K. Schreiber³, D. Roccatello¹. ¹Center of Immunopathology and Rare Diseases, University of Torino, Torino, Italy; ²UCSF, San Francisco, USA; ³Lupus Unit, GSTT, London, UK; ⁴University of Torino, Torino, Italy; ⁵Center of Immunopathology and Rare Diseases, UCSF, San Francisco, USA; ⁶Pathology, University of Torino, Torino, Italy

Background: In addition to glomerular lesions, renal vascular involvement is an important prognostic marker of lupus nephritis (LN). Among patients with various vascular changes, individuals with thrombotic microangiopathy (TMA) present with severe clinical manifestations and have a high mortality.

Objectives: We sought to assess renal outcomes and response to anti-thrombotic treatments in addition to conventional immunosuppression in patients with biopsy proven LN and TMA.

Methods: Clinical and renal histopathological data for 97 patients with biopsy-proven LN and TMA were retrospectively analysed. Antibody profiles, induction and maintenance therapies for LN, and anti-thrombotic treatments were collected. TMA lesions were classified into acute and chronic (table 1). A complete renal response (CR) was defined as proteinuria <0.5 g/24 hour and normal or near-