AB0734 SERUM IL-35 LEVELS IN SYSTEMIC SCLEROSIS AND RELATIONSHIP WITH CLINICAL FEATURES

M.E. Yayla¹, M. Torğutalp¹, I.E. Okatan¹, E. Uslu Yurteri¹, O. Küçükşahin², A. B. Keleşoğlu Dinçer¹, E.G. Aydemir Gülöksüz¹, S. Sezer¹, E. Us³, T.M. Turgay¹, G. Kınıklı¹, A. Ateş¹. ¹Department of Rheumatology, Ankara University Faculty of Medicine; ²Department of Rheumatology, Yıldırım Beyazıt University Faculty of Medicine; ³Department of Clinical Microbiology, Ankara University Faculty of Medicine, Ankara, Turkey

Background: Interleukin (IL)-35 is a member of the IL-12 family. Studies show that IL-35 is an important anti-inflammatory cytokine and suppresses effector T cell activity. Systemic sclerosis (SSc) is a chronic disease characterised by vascular damage, autoimmunity and fibrosis.

Objectives: In this study, we aimed to evaluate serum IL-35 level in SSc patients and its potential relation with disease findings.

Methods: Fifty-five SSc patients and 25 healthy controls were included in the study. All patients were diagnosed according to 2013 ACR/EULAR systemic sclerosis classification criteria. Serum IL-35 was measured using a commercial ELISA kit (Cloud-Clone Corp, Wuhan, China).

Results: SSc patients and healthy controls had similar ages and genders. The mean of serum IL-35 level was significantly higher in SSc patients (8.2±1.3 pg/ml) than in healthy controls (6.9±0.7 pg/ml) ($p < 0.001$). Serum IL-35 levels were similar in lcSSc (n=44) and dcSSc (n=11) groups (8.3±1.2 and 8±0.1, 1.6 pg/ml, respectively). There were no significant relationships between serum IL-35 level and digital ulcer, interstitial lung disease, pulmonary hypertension, tendon friction rub, gastrointestinal motility disorder, cardiac involvement and Anti-SCL70, anti-CENP-B positivities (table 1). There was not any correlation between IL-35 level and disease duration, modified Rodnan skin score, Valentini disease activity score, Medsger disease severity score, erythrocyte sedimentation rate and C-reactive protein (CRP) levels. Serum IL-35 levels between patients treated with or not vasodilating agents such as calcium channel blockers, endothelin receptor antagonists and prostaglandin analogues, and also immunosuppressive drugs did not differ significantly.

Abstract AB0734 – Table 1. Relationship between clinical and laboratory features and IL-35

		IL-35 (pg/ml)	P
		*	
Digital ulcer	- (n=32) + (n=23)	8.3±1.4 8.1±1	0.5
Gastrointestinal involvement	- (n=43) + (n=12)	8.3±1.2 8±1.5	0.512
Cardiac involvement	- (n=52) + (n=3)	8.1±1.3 9.5±0.8	0.073
Interstitial pulmonary disease	- (n=34) + (n=21)	8.3±1.3 8.1±1.3	0.593
Pulmonary arterial hypertension	- (n=51) + (n=4)	8.2±1.3 7.9±1.2	0.617
Arthritis	- (n=36) + (n=19)	8.4±1.3 7.9±1.2	0.160
Tendon friction rub	- (n=47) + (n=8)	8.2±1.3 8.3±1	0.798
Anti-Scl 70	- (n=34) + (n=20)	8.3±1.3 8.2±1.2	0.793
Anti-CENP B	- (n=33) + (n=21)	8.1±1.3 8.4±1.3	0.442

*mean ±std, + and – denote presence and absence, respectively.

Conclusions: It has been reported that serum IL-35 level is higher SSc patients than healthy volunteers and is associated with interstitial lung disease. In our study, the level of IL-35 was higher SSc patients than healthy controls. There is no relationship between clinical parameters and IL-35 levels, this may be due to small sample size. Our results suggest that IL-35 may have a role on the pathogenesis of SSc. Further investigations are needed in large sample size of SSc patients.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2605

AB0735 NAILFOLD CAPILLARY COUNTS CORRELATE MOST WITH COMPLICATIONS OF SYSTEMIC SCLERODEMA IN JAPANESE PATIENTS

A. Kondo¹, H. Dobashi¹, T. Kameda², M. Izumikawa², S. Nakashima², H. Shimada², R. Wakiya², M. Kato², N. Kadowaki². ¹Kagawa university, KIta, Japan; ²Department of Internal Medicine, Division of Hematology, Rheumatology and Respiratory Medicine, Kagawa university, KIta, Japan

Background: Connective tissue diseases (CTD) patients have various clinical manifestation including pulmonary hypertension (PH) and digital ulcer (DU). Especially, Raynaud's phenomenon (RP) is one of the common symptoms. On the other hand, "abnormal nailfold capillaries" are important as the findings that suggest RP associated with CTD. They are included in one of the items of classification criteria for Scleroderma (SSc) in 2013 Classification Criteria for Systemic Sclerosis (ACR/EULAR) ¹. Recently, some reports presented the usefulness of nailfold capillaroscopy (NFC) in SSc ². NFC findings showed enlarged/giant capillaries, fresh or old haemorrhages, avascular areas, ramified/bushy capillaries. These findings are known as "scleroderma pattern" features. Additionally, there are some reports that nailfold capillary density in SSc related with DU, PH and interstitial pneumonia (IP).

In this study, we clarify the association between nailfold capillary density and clinical manifestation by analysis of the case in our institution.

Objectives: To clarify the association between nailfold capillary density and clinical manifestation.

Methods: We enrolled Japanese SSc patients from May 2016 to May 2017 in our institution.

We measured total nailfold capillary count per 1 μm from second to fifth finger by NFC "OptiPIX Capillaroscopy Clinic 1.7.x". In addition, we investigated relationship with NFC findings and clinical manifestation such as current and previous DU, PH and IP.

Results: We enrolled 42 SSc patients. Total nailfold capillary counts significantly decreased, especially in SSc patients with, previous DU ($p = -0.4122$, $p = 0.0054$), and PH ($p = -0.3226$, $p = 0.0023$). In addition, there is significant difference between DU and existence of giant capillaries ($p = 0.3019$, $p = 0.0464$), existence of avascular areas ($p = 0.4057$, $p = 0.0063$). On the other hand, there is no significant difference between IP, PH and other "scleroderma pattern" features.

Conclusions: Our study revealed that total nailfold capillary counts were associated with PH and DU in SSc patients. We suggest that nailfold capillary density may most predict complications with SSc patients.

REFERENCES:

- [1] Markus Mueller, et al. Relation of Nailfold Capillaries and Autoantibodies to Mortality in Patients With Raynaud Phenomenon. *Circulation*. 133:509–517, 2016.
- [2] Maurizio Cutolo, et al. Nailfold Videocapillaroscopic Features and Other Clinical Risk Factors for Digital Ulcers in Systemic Sclerosis. *Arthritis & Rheumatology*. 68 (10): 2527–2539, 2016.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7510

AB0736 SYSTEMIC SCLEROSIS SINE SCLERODERMA: CHARACTERISTICS OF A SOUTH INDIAN COHORT FROM A TERTIARY CARE CENTRE

B. Chilukuri^{1,2}, S. Chinnadurai², B. Mahendran³, R. Ramamoorthy², B. Selvakumar², R. Sankaralingam⁴. ¹Dept of Rheumatology, Parvathy Hospital; ²Institute of Rheumatology, Madras medical college, Chennai; ³Dept of Rheumatology, KIMS Hospital, Trivandrum; ⁴Dept of Rheumatology, SRMC, Chennai, India

Background: Systemic Sclerosis is a complex disorder characterised by autoimmunity, vasculopathy and fibrosis. The hallmark of Systemic Sclerosis is skin thickening. Systemic Sclerosis sine scleroderma is a variant of Systemic sclerosis which shares visceral, serological and vascular manifestations but lacks skin thickening. Systemic Sclerosis sine scleroderma still remains a rarely reported subtype of Systemic Sclerosis unlike commonly reported limited cutaneous and diffuse cutaneous types.

Objectives: Aim of this study was to analyse the characteristics of patients with Systemic Sclerosis sine scleroderma from a cohort of Systemic Sclerosis patients.

Methods: This study was done at Institute of Rheumatology, Madras Medical College, Chennai, India. It was a retrospective observational study (January 2006 to November 2016). Patients satisfying ACR/EULAR 2013 Classification criteria for Systemic Sclerosis were included in this study. Patients with Overlap syndrome, MCTD, UCTD, children and pregnant women were excluded. Poormoghim criteria was used for diagnosis of Systemic Sclerosis sine scleroderma: Absent skin