Abstract AB0841 – Table 1. Comparison of the perceived social support, depression and loneliness scores with the Morisky Green Levine Scale subgroups.

Characteristics	Morisky Green Levine medication adherence groups				F	p
	Overall	High (n=43) ^a	Medium (n=66) ^b	Low (n=10) ^c		
Multidimensional Scale of Perceived Social Support total score ^{12–84}	63.52 ±17.13	65.09 ±16.85	64.65 ±16.59	49.30 ±17.01	3.958	0.022
Beck Depression Inventory (0–63) [*]	13.65 ±10.41	13.07 ±11.29	12.15 ±8.51	26.10 ±10.73	8.952	<0.001
UCLA Loneliness Scale ^{20–80}	37.34 ±11.22	35.19 ±11.26	37.56 ±10.90	45.20 ±10.56	3.388	0.037

F: Oneway ANOVA, Statistically significant between a and c, and b and c (p<0.05). * It shows the lowest-highest scores that can be taken from the scale and its sub-dimensions.

Conclusions: In this study, it was shown that as the average age and social support scores of patients decreased, and as the BASDAI, spinal pain, spinal pain at night, fatigue, patient global assessment, loneliness and depression scores increased, adherence to treatment were decreased. It is thought that patients should be handled holistically in terms of biopsychosocial aspect in order to improve adherence to medical treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5873

AB0842 DRUG SURVIVAL AND EFFECTIVENESS OF THE FIRST TNF INHIBITORS IN PATIENT WITH LATE ONSET SPONDYLARTHROSIS: TREASURE REAL-LIFE RESULTS

N. Alpay Kanitez¹, S. Kiraz², I. Ertenli², O. Küçükşahin², E. Dalkılıç³, C. Bes¹, T. Kaşifoğlu⁴, H. Emmungil⁵, S.S. Koca⁶, M. Çınar², S. Yılmaz², V. Yazısız⁷, B. Keleşoğlu Dinçer², S. Akar⁸, O. Gercik⁸, A. Ateş², S. Erten², S. Turan⁵, B. Yağız², B. Öz⁶, S. Yaşar Bilge⁴, L. Kılıç², G.K. Yardımcı², B.N. Coşkun³, U. Kalyoncu². ¹Treasure biological registry, İstanbul; ²Treasure biological registry, Ankara; ³Treasure biological registry, Bursa; ⁴Treasure biological registry, Eskişehir; ⁵Treasure biological registry, Edirne; ⁶Treasure biological registry, Elazığ; ⁷Treasure biological registry, Antalya; ⁸Treasure biological registry, İzmir, Turkey

Background: Spondylarthritis (SpA) are usually observed in young patients and a clinical onset after 45 years is rare. The average life expectancy is getting longer and the proportion of late onset of SpA in rheumatology practice may become more common. SpA patients with late onset may have a distinctive clinical pattern in terms of functional impairment, extra-articular disease, co-morbidities and treatment status.

Objectives: The aim of this study was to compare the drug survival and effectiveness of tumour necrosis factor (TNF) inhibitors in patients with late onset SpA (LoSpA) compared with early onset SpA (EoSpA).

Methods: TREASURE is a prospective, multicenter biological treatments registry from Turkey since 2016. It includes 15 different rheumatology centres. Patients with SpA fulfilling the ASAS criteria from TREASURE database were divided into two groups as LoSpA (symptom onset >45 years of age) and EoSpA (symptom onset ≤45 years of age). Drug retention rates of first TNF inhibitors were calculated using the time until drug discontinuation independent of the reason that drug interruption. Specific reasons for discontinuing drugs were also assessed.

Results: Of 1382 SpA patients treated with TNF inhibitors, 9.4% (n=130) were included in the LoSpA and 90.6% (n=1252) were included in the EoSpA group. LoSpA had more female, enthesitis and psoriasis. The median treatment duration was 53 months in LoSpA and 61 months in EoSpA. The baseline disease activity measures were similar except from ASDAS-ESh which is higher in LoSpA (table 1). The rate of major treatment response (BASDAI50) was lower in LoSpA than EoSpA at the last visit (26.1% vs 46.2%; p=0.009). Regarding the survival rates of TNF inhibitors, there was no significant difference between the patient groups (figure 1). The major cause of discontinuation in both of the groups was drug ineffectiveness (68.6% in LoSpA and 60% in EoSpA, p>0.05).

Abstract AB0842 – Table 1

	LoSpA n: 130	EoSpA n:1252	p values
Age [median (min-max)]	56.5 (45–75)	35 (0–76)	<0.001*
Female n (%)	90 (69)	494 (40)	<0.001*
Disease duration [median (min-max)]	61 (1–334)	108 (0–686)	<0.001*
BASDAI [median (min-max)]	6.1 (0–10)	6 (0–10)	0.748
BASFI [median (min-max)]	5.3 (0–10)	4.5 (0–10)	0.098
ASDAS-ESH [median (min-max)]	3.29 (1.23–4.92)	2.91 (0–5.7)	0.001*
ASDAS-CRP [median (min-max)]	3.55 (1.49–5.92)	3.35 (0.7–8)	0.251

Abstract AB0842 – Figure 1

Conclusions: LoSpA patients were almost 10% of all biological registry. LoSpA patients were predominantly female, and they had relatively higher baseline disease activity and lower biological treatment response. On the other hand the drug survival rate and discontinuation reasons of TNF inhibitors in the LoSpA group was comparable to that in the younger group.

REFERENCE:

- [1] - Chen HA, Chen CH, Liao HT, Lin YJ, Chen PC, Chen WS, et al. Clinical, functional, and radiographic differences among juvenile-onset, adult-onset, and late-onset ankylosing spondylitis. *J Rheumatol.* 2012; 39:1013–8

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4844

AB0843 EFFECT OF TNF INHIBITORS ON BONE MICROARCHITECTURE IN PATIENTS WITH ANKYLOSING SPONDYLITIS: A LONGITUDINAL STUDY BASED ON HIGH-RESOLUTION PERIPHERAL QUANTITATIVE BASED (HRPQCT)

N. Nigil Haroon¹, E. Szabo², A.M. CHEUNG³, R. Inman⁴. ¹Medicine, Northern Ontario School of Medicine, Sudbury; ²Osteoporosis program, UHN; ³Medicine, Univ of Toronto; ⁴Medicine, unit of Toronto, Toronto, Canada

Background: Ankylosing spondylitis (AS) is associated with high risk of fractures. BMD, bone microarchitecture and strength are negatively affected in AS. TNF inhibitors such as etanercept, adalimumab, golimumab and infliximab are the mainstay of treatment in AS. However no data is available on the effect of TNF inhibitors on bone microarchitecture and strength.

Objectives: This study aimed to assess the effect of TNF inhibitors on bone microarchitecture in patients with AS.

Methods: AS was defined by Modified New York criteria. Areal BMD was measured by DXA. Volumetric BMD (vBMD) and bone microarchitecture were measured using high-resolution peripheral quantitative CT (HRpQCT) at the radius and tibia at baseline and after one year of treatment with TNF inhibitors. Intake of calcium and vitamin D were optimised.

Results: There were 31 subjects (58% men). Mean (±SD) age and BASDAI were 40±14 years and 4.1±2.1 respectively. Median duration of disease was 14 (IQ: 6.5–25.5) years. Mean duration of follow-up was 15 months. Areal BMD (n=22) at lumbar spine (1.053±0.235 vs. 1.049±0.202, p=0.89), total hip (0.944±0.152 vs. 0.912±0.164, p=0.5), and femoral neck (0.955±0.151 vs. 0.954±0.191, p=0.2) did not change significantly. HRpQCT (n=31) on follow-up demonstrated that total, trabecular and cortical volumetric BMD were unchanged at both radius and tibia (table 1). Also, HRpQCT based trabecular parameters such as trabecular number, thickness and separation, BV/TV and cortical parameters such as cortical porosity and thickness remained stable (table 1). FEA estimates of bone stiffness and

stress tended to be lower at the radius on follow-up however these parameters were not significantly different at the tibia (table 1).

Abstract AB0843 – Table 1

	RADIUS			TIBIA		
	Baseline	Follow-up	p	Baseline	Follow-up	p
Total volumetric BMD (mg/cm ³)	318.93 ±311.80	317.37 ±378.61	0.49	301.89 ±48.99	302.06 ±50.71	0.70
Cortical volumetric BMD (mg/cm ³)	857.22 ±51.43	851.48 ±52.96	0.08	869.69 ±38.21	866.62 ±39.91	0.11
Trabecular volumetric BMD (mg/cm ³)	158.33 ±42.34	156.81 ±41.88	0.21	155.200 ±44.34	154.26 ±45.25	0.35
BV/TV (%)	0.132 ±0.035	0.131 ±0.035	0.19	0.129 ±0.037	0.129±0.04	0.37
Trabecular number (per mm ³)	1.985 ±0.302	2.017 ±0.319	0.32	1.793 ±0.36	1.809±0.37	0.56
Trabecular thickness (mm)	0.067 ±0.013	0.065 ±0.013	0.12	0.072 ±0.01	0.071±0.01	0.17
Trabecular separation (mm)	0.435 ±0.104	0.429 ±0.106	0.46	0.483 ±0.13	0.478±0.13	0.06
Cortical thickness (mm)	0.765 (0.72–0.85)	0.760 (0.71–0.82)	0.80	0.751 (0.58–0.84)	0.745 (0.55–0.83)	0.78
Cortical porosity (%)	0.019 (0.71–0.82)	0.017 (0.016–0.024)	0.81	0.049 (0.02–0.06)	0.048 (0.017–0.065)	0.80
Stiffness (N/mm)	1.37±0.42	1.33±0.34	0.03	1.57±0.31	1.58±0.31	0.72
Stress (MPa)	27.23 ±10.52	26.30±9.83	0.03	32.78±7.8	32.70±7.71	0.71

Conclusions: This is the first study to document the changes in bone strength in AS patients with the use of TNF inhibitors. Treatment with TNF inhibitors might maintain bone microarchitecture at cortical and trabecular sites in patients with AS.

Disclosure of Interest: N. Nigil Haroon Consultant for: AMGEN Canada, E. Szabo: None declared, A. CHEUNG Consultant for: AMGEN, ELI LILLY, R. Inman Grant/research support from: AMGEN, Consultant for: AMGE, ABBVIE, NOVARTIS, JANNSEN, MERCK

DOI: 10.1136/annrheumdis-2018-eular.1023

AB0844

EFFECTIVENESS AND RETENTION RATE OF CERTOLIZUMAB PEGOL IN SPONDYLOARTHRITIS. REAL LIFE DATA

R. Expósito-Molinero¹, R. Garcia-Portales², J.R. Lamua-Riazuelo³, A. Urruticoechea-Arana⁴, P. Navarro-Alonso⁵, J.S. Rey-Rey⁶, M. Fernandez-Prada⁷, C.M. Gonzalez Fernandez⁸, on behalf of RENACER Study Group.
¹Rheumatology, Hospital Comarcal de Laredo, Laredo; ²Rheumatology, Hospital Virgen de la Victoria, Malaga; ³Rheumatology, Hospital del Henares, Alcala de Henares; ⁴Rheumatology, Hospital Can Misses, Ibiza; ⁵Rheumatology, Hospital de Fuenlabrada, Fuenlabrada; ⁶Rheumatology, Hospital Virgen de la Salud, Toledo; ⁷Rheumatology, Hospital Universitario de Guadalajara, Guadalajara; ⁸Rheumatology, Hospital General Universitario Gregorio Marañón, Madrid, Spain

Background: Certolizumab pegol (CZP) is available for patients with axial spondyloarthritis (axSpA), including ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA). The efficacy and the safety of CZP are well established from clinical trials. However, evidence of its effectiveness in regular clinical practice is limited.

Objectives: To evaluate the effectiveness and safety of CZP in a real word setting in axSpA patients.

Methods: Multicentric cohort of SpA patients treated with CZP according to routine clinical practice. The study was approved by the local Ethics Committee. Maximum follow-up was 12 months. Clinical response was evaluated through BASDAI, ASDAS, BASFI and MASES scores. Safety variables: discontinuation rate.

Results: 336 patients with axSpA were included: 56.5% male, mean age 45.8 (±12.1) years, median disease time 4.3 (0, 49.5) years, 68.5% of patients were HLAB27 positive, and never smokers 64.7%. Prior bDMARD received (27.2% none; 37.9% 1, 35%≥2). At baseline 36.8% had concomitant DMARDs and 82.8% NSAIDs. 31.8% of patients had peripheral arthritis and 42.7% entesitis at baseline. CZP retention time 10.3 months. Statistically significant differences in BASDAI, BASFI, ASDAS y MASES were observed at the last visit comparing to baseline (table 1). In the last observation, 41.0% of the patients achieved BASDAI50, 34% were in ASDAS remission (ASDAS <1.3) and 49% presented a minimal clinical improvement (ASDAS at least 1.1). 46.3% of patients had resolution of the entesitis (MASES=0). According to Kaplan-Meier analysis, the drug survival of CZP was 83.3%, and no differences were observed in retention rates

when CZP were used as first option (83.3%) or after failure to other biologicals (79.1%) (figure 1). 56/336 (16.7%) withdrawn CZP treatment: 34/336 (10.1%) due to lack of effectiveness, 16/336 (4.8%) due to adverse event and 6/336 (1.8%) for other reason. Serious adverse event was found in 23/336 (6.8%) patients.

Abstract AB0844 – Table 1. Evolution of clinical variables of activity

	Baseline	Last visit
BASDAI evolution	6,2±1.6(n=263)	3,9±2.2(n=260)*
BASFI evolution	5,6±2.1(n=225)	3,7±2.3 (n=241)*
ASDAS evolution	3,7±2.7(n=322)	2,4±2.2 (n=321)*
MASES score	3,9±2.7(n=78)	1,2±1.8(n=51)*

*p<0,001, Wilcoxon's test

Abstract AB0844 – Figure 1. CZP Retention rate in axSpA patients without and with previous biological experience.

Conclusions: Real life experience from this nationwide rheumatology study, demonstrated the effectiveness and safety of CZP in patients with axSpA, with a significant reduction of BASDAI, BASFI, ASDAS and MASES scores. No differences were observed in the retention rate regardless previous biological treatment.

Disclosure of Interest: R. Expósito-Molinero: None declared, R. Garcia-Portales Consultant for: Celgene, Speakers bureau: UCB, Pfizer, Roche, J. R. Lamua-Riazuelo: None declared, A. Urruticoechea-Arana: None declared, P. Navarro-Alonso: None declared, J. S. Rey-Rey Speakers bureau: UCB, Abbvie, Pfizer, BMS, Roche, Celgene, M. Fernandez-Prada: None declared, C. Gonzalez Fernandez Consultant for: MSD, Janssen, Novartis, Celgene, Speakers bureau: Abbie, Janssen, MSD, Novartis, Roche, UCB, BMS

DOI: 10.1136/annrheumdis-2018-eular.5552

AB0845

THE EFFECT OF BIOLOGIC DISEASE-MODIFYING ANTIRHEUMATIC DRUGS PATIENT REPORTED OUTCOMES IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS; A SYSTEMATIC LITERATURE REVIEW AND A CALL FOR ACTION

S. Rodrigues-Manica^{1,2}, J. Leite Silva³, A.R. Machado⁴, C. Coelho⁵, J. Duarte⁶, E. Vieira-Sousa⁴, J. Tavares-Costa³, F.M. Pimentel-Santos^{1,2}. ¹CEDOC; ²Egas Moniz, Centro Hospitalar de Lisboa Ocidental, Lisbon; ³Unidade Local de Saúde do Alto Minho, Ponte de Lima; ⁴Centro Hospitalar de Lisboa Norte, EPE, Hospital de Santa Maria, Lisbon Academic Medical Centre; ⁵Genetics Laboratory, Institute of Environmental Health, Lisbon School of Medicine, University of Lisbon; ⁶Medical Department, Novartis Pharma, Pharmaceutical products, Lisbon, Portugal

Background: Patient reported outcomes (PROs) have gained relevance in the evaluation of several diseases such as axial spondyloarthritis (axSpA). They allow the clinician to have a quantitative measurement of several aspects of the disease, according to the patient perspective.

Objectives: In this review we intended to evaluate the efficacy of different biologic disease-modifying anti-rheumatic drugs (bDMARD) in achieving the minimum clinically important difference (MCID) in several PROs. Data from randomised controlled trials (RCT) conducted in radiographic axSpA (r-axSpA) and non-radiographic axSpA (nr-AxSpA) patients were included.

Methods: A systematic literature review (SLR) was performed using the MEDLINE database (August 17 2017) with the filters "published in the last 10 years" and "humans". Abstracts from the EULAR 2017 were also considered. The PICO (P,