AB0983

HYALURONAN DERIVATIVE HYMOVIS® INCREASES CARTILAGE VOLUME AND TYPE II COLLAGEN TURNOVER IN OSTEOARTHRITIC KNEE: DATA FROM MOKHA STUDY

Y. Henrotin¹, R. Bannuru², M. Malaise³, K. EA⁴, C. Confavreux⁵, J. Bentina⁶, D. Urban-Choffray⁷, T. Conrozier⁸, Brasseur⁹, T. Thomas¹⁰, A. Marrinello¹¹, N. Giordan¹², P. Richette¹³. ¹Bone and Cartilage Research Unit, University of Liege, Liège, Belgium; ²Center for Treatment Comparison and Integrative Analysis, Tufts Medical Center, Boston, USA; ³Service de Rhumatologie, CHU Sart-Tilman, Liège, Belgium; ⁴Service de Rhumatologie, Hôpital Lariboisière, Paris; ⁵Department of Rheumatology, Hospices Civils de Lyon, Lyon, France; ⁶Service de Rhumatologie, CHU Brugmann, Bruxelles; ⁷Service de Médecine Physique, CHR Citadelle, Liège, Belgium; ⁸Service de Rhumatologie, Hôpital Nord Franche-Comté, Trévenans, France; ⁹Service de Rhumatologie, CHU UCL Namur, Yvoir, Belgium; ¹⁰Service de Rhumatologie, CHR Metz-Thionville, Metz, France; ¹¹Fidia Farmaceutici, S.p.A.; ¹²Fidia Farmaceutici, S.p.A., Fidia, Abano, Italy; ¹³Centre Viggo Petersen, Hôpital Lariboisière, Paris, France

Background: Intra-articular injections of hyaluronan represent one of the well-accepted standard of care for treating symptomatic knee osteoarthritis (OA). Until now, not much is known about the structural-modifying effect of this treatment justifying this pilot study.

Objectives: This exploratory non-controlled study aims to study effects of HYMOVIS on imaging, biological and clinical variables.

Methods: Forty six patients with symptomatic knee OA (mean age 61.4 years [min.35-max.80; 67.4% female; Kellgren and Lawrence grade II and III (63% and 37%, respectively); mean BMI 30.6 kg/m²) were enrolled in this open-label, prospective, multicenter, pilot study. Patients received two treatment cycles of intra-articular injections (3 mL) of Hymovis (8 mg/mL of the 500–730 kDa hyaluronic acid hexadecylamide) at 6 months interval. Each treatment cycle corresponded to two intra-articular injections one week apart. All patients had MRI of the target knee at baseline and 1 year, and blood samples (D30, D90, D180, D210 and D360) to assess joint biomarkers. The primary outcome was the change in type II collagen-specific biomarkers (Coll2-1, Coll2-1NO2 and CTX-II) after Hymovis treatment versus baseline. Secondary endpoints included levels changes in AGG, COMP, PIINP, MMP-3, MPO and IL-6 serum biomarkers, the ratio Coll2-1/PIINP, CTX-II/PIINP, MRI cartilage volume and KOOS index.

Results: Coll2-1 serum levels significantly increased overtime (0.05<P<0.001) while Coll2-1NO2 levels was only increased at D360 (p<0.05). Serum PIINP levels also progressively and significantly enhanced with time (p<0.001). In contrast, other serum biomarker levels including CTX-II, AGG, COMP, MMP-3, MPO or IL-6 did not change significantly overtime. Interestingly, the ratios Coll2-1/PIINP and CTX-II/PIINP decreased (p<0.005), indicating a decrease of cartilage catabolism. Compared to baseline value, MRI cartilage volume increased in lateral femoral and lateral trochlea compartments (p<0.05) and not in medial compartment. Interestingly, WOMBS effusion score, an indicator of synovitis, significantly decreased (p<0.016). Finally, global KOOS score and subscales significantly increased overtime (p<0.001) while pain at rest, walking pain and patients or investigators global assessment of disease activity decreased (p<0.001). The safety profile was favourable with a low incidence of injection-site pain.

Conclusions: Hymovis, a well-tolerated intra-articular treatment, significantly enhanced type II collagen turnover as suggested by the increase in Coll2-1 and PIINP levels and cartilage volume observed by MRI in lateral knee compartment. Importantly, this study highlighted the potential symptomatic benefit of Hymovis on pain and function and provides critical information for the design of a larger phase III clinical trial.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4198

AB0984

CADMIUM TOXICITY AS A PROBABLE CAUSE OF SMOKING INDUCED BONE LOSS

A.A. Elbeily^{1,2}, H.A. Eldosouky¹. ¹Rheumatology, Al-Azhar University of Medicine, Cairo, Egypt; ²Rheumatology, St Thomas Hospital, London, UK

Background: Cigarette smoking supposed to be a risk factor for osteoporosis. There is an inverse relationship between smoking and both bone mass and fracture risk. Tobacco smoking is the most important single source of cadmium exposure in the general population. The absorption of cadmium from the lungs is much more effective than that from the gut.

Objectives: This study was designed to evaluate the effect of cigarette smoking on bone mineral density, due to cadmium toxicity.

Methods: this study was carried on 100 persons, selected from AL-Azhar university hospital and divided into three groups: group I: included 40 persons with active smokers; group II: included 40 persons with passive smokers and group III included 20 nonsmokers. All persons were subjected to full history taking,

thorough clinical examination, routine lab tests, serum and urinary cadmium and lead, and bone mineral density was measured by DEXA.

Results: Serum and urinary cadmium and lead were statistically significantly higher in group I in comparison to groups II or III and in group II in comparison to group III. Also, there was statistically significant decrease of BMD in group I in comparison to either group II or group III and in group II in comparison to group III. There was an inverse statistically significant correlation between serum and urinary cadmium and bone mineral density.

Abstract AB0984 – Table 1. Comparison between studied groups as regard serum and urinary cadmium levels

		Mean	S. D	Minimum	Maximum	F	P
Serum cadmium (µg/l)	Group I	0.449	0.287	0.18	1.80	27.92	<0.001(S)
	Group II	0.232*	0.040	0.16	0.32		
	Group III	0.094*	0.047	0.01	0.16		
Urinary cadmium (µg/l)	Group I	0.555	0.304	0.30	2.00	27.16	<0.001(S)
	Group II	0.335*	0.042	0.26	0.42		
	Group III	0.181*	0.052	0.11	0.26		

*=statistically significant decrease in comparison to group I or group II

#=statistically significant decrease in comparison to group I

Conclusions: Results of the present study revealed that: there are harmful effects of smoking on the bone mineral density and it may be occurred by direct (increased blood and urinary levels of both cadmium and lead) or indirect effects (effects of both renal and liver functions) of cadmium and lead.

REFERENCES:

- Aoshima K, Katoh T, Kasuya M. Renal effects of environmental exposure to cadmium in middle-aged population of Jinzu River basin in Toyama, Japan: 2000–2002 study. *Nippon Eiseigaku Zasshi* 2006;61:69–80.
- Bernard A. Cadmium & its adverse effects on human health. *Indian J Med Res* 2008; 128:557–64.
- Bhattacharyya MH. Cadmium Osteotoxicity in Experimental Animals: Mechanisms and Relationship to Human Exposures. *Toxicol Appl Pharmacol* 2009; 238(3): 258–265.
- Hapidin H, Othman F, Soelaiman IN, Shuid AN, Luke DA, Mohamed N: Negative effects of nicotine on bone-resorbing cytokines and bone histomorphometric parameters in male rats. *J Bone Miner Metab* 2007, 25 (2):93–8.
- Noël L, Huynh-Delerme C, Guérin T, Huet H, Frémy JM, Kolf-Clauw M. Cadmium accumulation and interactions with zinc, copper, and manganese, analysed by ICP-MS in a long-term Caco-2 TC7 cell model. *Biometals*. 2006; 19(5):473–81.
- Satarug S. Long-term exposure to cadmium in food and cigarette smoke, liver effects and hepatocellular carcinoma. *Curr Drug Metab*. 2012 Mar; 13 (3):257–71.
- Schutte R, Nawrot TS, Richart T, Thijs L, Vanderschueren D, Kuznetsova T, Van Hecke E, Roels HA, Staessen JA. Bone Resorption and Environmental Exposure to Cadmium in Women: A Population Study. *Environ Health Perspect* 2008; 116:777–783.
- Tamaki J, Iki M, Fujita Y, Kouda K, Yura A, Kadowaki E, Sato Y, Moon JS, Tomioka K, Okamoto N, Kurumatani N: Impact of smoking on bone mineral density and bone metabolism in elderly men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) study. *Osteoporos Int* 2010, 22 (1):133–141.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3369

AB0985

PERSISTENCE AND ADVERSE EVENTS IN PATIENTS TREATED WITH DENOSUMAB

A. Lopez, I. Janta, J.C. Nieto, J. Martinez Barrio, B. Serrano, F. Garcia, J. C. Sanchez, C. Garaballu, T. Del Rio, C. Gonzalez, J. Lopez Longo, I. Monteagudo. *Rheumatology, Hospital Gregorio Marañón, Madrid, Spain*

Background: Denosumab (DNS) is a human monoclonal antibody directed against RANKL, which blocks the maturation of the osteoclast, inhibiting bone resorption. The binding of DNS to RANKL suppresses bone resorption mediated by osteoclasts and decreases bone turnover.