Results: This analysis included 29 987 patients, representing 56 951 patient-years of exposure (table 2). The majority of adalimumab exposure was in RA studies. The most frequently reported SAE of interest was infection (highest incidences in CD, RA, UV, and UC). Overall and for most of the adalimumab populations (AS, PsA, Ps, UC, CD, and RA), the observed number of deaths was below what would be expected in an age- and sex-adjusted population (table 1). For HS, nr-axSpA, pSpA, and UV studies, the small size of these trials precluded accurate assessment of the standardised mortality ratio, and the 95% CIs all included 1.0.

Abstract OP0233 – Table 1 Standardised mortality ratios across indications

Indication	SMR	95% CI
RA (n=15,511)	0.74	0.63–0.87
AS (n=2026)	0.14	0.00–0.77
nr-axSpA (n=863)	1.22	0.14–4.40
pSpA (n=165)	1.84	0.21–6.65
PsA (n=837)	0.34	0.04–1.24
Ps (n=3732)	0.34	0.15–0.64
HS (n=733)	1.50	0.40–3.84
CD (n=3896)	0.44	0.14–1.02
UC (n=1739)	0.37	0.12–0.87
UV (n=464)	1.23	0.45–2.68
Total (n=29,986)	0.65	0.57–0.74

AS, ankylosing spondylitis; CD, Crohn's disease; HS, hidradenitis suppurativa; nr-axSpA, non-radiographic axial SpA; Ps, plaque psoriasis; PsA, psoriatic arthritis; pSpA, peripheral SpA; RA, rheumatoid arthritis; SMR, standardised mortality ratio; SpA, spondyloarthritis; UC, ulcerative colitis; UV, uveitis.

Abstract OP0233 – Table 2 Incidence rates of serious adverse events of interest*

Characteristic	RA	AS	nr-axSpA	pSpA	PsA	Ps	HS	CD	UC	UV	Total†
n	15,512	2026	863	165	837	3732	733	3896	1739	464	29,987
Exposure, PYs	24,922	2120	709	391	998	5479	1198	4359	3407	1151	56,951
Serious infection	4.6	1.8	2.5	1.0	2.8	1.8	2.8	6.9	3.5	4.1	3.7
Tuberculosis	0.3	0.1	0.1	0.3	0.2	0.2	0	0.2	<0.1	0.4	0.2
Active	0.3	0.1	0.1	0	0.2	0.2	0	0.1	<0.1	0.2	0.2
Latent	<0.1	0	0	0.3	0	0	0	<0.1	0	0.3	<0.1
Opportunistic infection‡	<0.1	0	0.1	0	0	0	0	<0.1	<0.1	0.4	<0.1
Demylinating disorder	<0.1	<0.1	0	0	0	0	0	0.1	<0.1	0.3	<0.1
Lupus-like syndrome	<0.1	<0.1	0.1	0	0	0	0	<0.1	<0.1	<0.1	<0.1
CHF	0.2	<0.1	0	0	0.1	0.2	0	<0.1	<0.1	0.2	0.2
Ps new onset/worsening	<0.1	<0.1	0	0	0.1	<0.1	<0.1	<0.1	<0.1	0	<0.1
Malignancy‡	0.7	0.2	0.1	0.3	0.2	0.5	0.5	0.4	0.6	0.7	0.6
Lymphoma	0.1	<0.1	0	0	0.2	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
NMSC	0.2	0.2	0	0	0.1	0.1	<0.1	<0.1	<0.1	0.2	0.1
Melanoma	<0.1	<0.1	0	0	0.2	0	0	<0.1	0	<0.1	<0.1
Sarcoidosis	<0.1	<0.1	0	0	0	0	0	0	0	<0.1	<0.1
Any AE leading to death	0.7	<0.1	0.3	1.0	0.3	0.2	0.5	0.1	0.1	0.6	0.5

AE, adverse event; AS, ankylosing spondylitis; CD, Crohn's disease; CHF, congestive heart failure; HS, hidradenitis suppurativa; NMSC, non-melanoma skin cancer; nr-axSpA, non-radiographic axial SpA; Ps, plaque psoriasis; PsA, psoriatic arthritis; pSpA, peripheral SpA; PY, patient-year; RA, rheumatoid arthritis; SpA, spondyloarthritis; UC, ulcerative colitis; UV, uveitis.

*Rates in events/100 PYs.

†Total includes the 10 populations shown plus 20 patients with Behcet's disease (35.5 PY).

‡Excludes oral candidiasis and tuberculosis.

§Excludes lymphoma, hepatosplenic T-cell lymphoma, leukemia, NMSC, and melanoma.

Conclusions: This analysis of data from clinical trials of adalimumab demonstrated an overall safety profile consistent with previous findings and with the TNF inhibitor class. No new safety signals or tolerability issues with adalimumab treatment were identified and, for most indications, the mortality rate was below what would be expected in an age- and sex-adjusted population. Efficacy and safety data continue to support the well-established benefits of adalimumab for the approved indications.

Acknowledgements: AbbVie funded the study, contributed to its design, and participated in data collection, analysis and interpretation of the data, and in writing, review, and approval of the publication. Medical writing (funded by AbbVie): Maria Hovenden, PhD, and Janet Matsuura, PhD, of CPS.

Disclosure of Interest: G. Burmester Grant/research support from: AbbVie, Bristol-Myers Squibb, Merck, Pfizer, Roche, and UCB, Consultant for: AbbVie, Bristol-Myers Squibb, Merck, Pfizer, Roche, and UCB, Speakers bureau: AbbVie, Bristol-Myers Squibb, Merck, Pfizer, Roche, and UCB, R. Panaccione Grant/research support from: Abbott Laboratories, Xcan, Bristol-Myers Squibb, Centocor, Elan, Millenium, and Procter and Gamble; honoraria from Abbott Laboratories, AstraZeneca, Byk Solvay, Centocor, Elan, Janssen, Procter and Gamble, Prometheus, Schering-Plough, and Shire, Consultant for: Abbott Laboratories, AstraZeneca, Bristol-Myers Squibb, Centocor, Elan, Ferring, GlaxoSmithKline, Procter and Gamble, Schering-Plough, Shire, and UCB, K. Gordon Grant/research support from: AbbVie, Amgen, Boehringer Ingelheim, Eli Lilly, and

Janssen, Consultant for: AbbVie, Amgen, Boehringer Ingelheim, Celgene, Eli Lilly, Janssen, Novartis, and Pfizer, J. Rosenbaum Shareholder of: Royalties: UpToDate, Grant/research support from: Alcon Research Institute, Consultant for: Abbvie, Gilead, Santen, Regeneron, UCB, Cavtherx, Portage, Eyeevensys, and Stem Cell Inc, Speakers bureau: Mallinckrodt, D. Arikian Shareholder of: AbbVie, Employee of: AbbVie, W. Lau Shareholder of: AbbVie, Employee of: AbbVie, R. Tarzynski-Potempa Shareholder of: AbbVie, Employee of: AbbVie
DOI: 10.1136/annrheumdis-2018-eular.3158

FRIDAY, 15 JUNE 2018

Clinical and therapeutic aspects of vasculitis_____

OP0234 INFLIXIMAB THERAPY IN PATIENTS WITH TAKAYASU ARTERITIS

E.H. Park¹, E.Y. Lee¹, Y.J. Lee², Y.J. Ha², B.Y. Choi³, W.-H. Yoo⁴, J.C. Paeng⁵, H. Y. Suh⁵, J.S. Park⁶, S. Shon⁶, Y.W. Song¹. ¹Department of Internal Medicine (Division of Rheumatology), Seoul National University Hospital, Seoul; ²Department of Internal Medicine (Division of Rheumatology), Seoul National University Bundang Hospital, Seongnam; ³Department of Internal Medicine (Division of Rheumatology), Seoul Medical Center, Seoul; ⁴Department of Internal Medicine (Division of Rheumatology), Chonbuk National University, Jeonju; ⁵Department of Nuclear Medicine, Seoul National University Hospital; ⁶Department of Molecular Medicine and Biopharmaceutical Sciences, Graduate School of Convergence Science and Technology, and College of Medicine, Seoul National University, Seoul, Korea, Republic Of

Background: Takayasu arteritis (TAK) is a chronic inflammatory disease that predominantly affects the aorta and its main branches. Glucocorticoids (GCs) are the cornerstones of the initial treatment of TAK. However, most patients relapse with steroid withdrawal.

Objectives: To evaluate the efficacy and safety of infliximab (IFX) in Korean patients with active TAK.

Methods: Patients with active TAK were enrolled in a single-centre prospective open label trial. Active disease was defined according to the National Institutes of Health (NIH) criteria. Concomitant GCs were tapered to prednisone ≤10 mg/day or equivalent at 2 weeks prior to the initiation of IFX. Patients received intravenous infusions of IFX, at a starting dose of 5 mg/kg at weeks 0, 2, 6, and then every 8 weeks, up to week 46, and were followed up to week 54. At week 30, patients with partial remission received increased dose of IFX by 1.5 mg/kg, and patients who failed with IFX terminated the study. At week 38 and 46, patients with symptom of active disease or high serum level of erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP), were instructed to increase the IFX dose by 1.5 mg/kg, up to 9.5 mg/kg, at each point. All the patients underwent Positron Emission Tomography-Computed Tomography (PET-CT) at baseline and week 30. The primary efficacy end point was the achievement of partial or complete remission at week 30.

Results: Twelve patients with TAK were enrolled and treated with IFX; 1 patient with study violation was excluded from analysis. At week 30, 3 patients (27.3%) achieved complete remission and 6 patients (54.5%) achieved partial remission. Statistically significant improvements were seen at week 30 for all of major secondary measures, including change from baseline in Indian Takayasu Clinical Activity Score 2010 ITAS 2010 (median 11.0, interquartile range [IQR] 10.0–11.8; 6.0, IQR 5.0–9.0, p=0.004), ITAS.A (14.0, IQR 12.0–14.0; 7.0, IQR 6.0–10.5, p=0.003) and serum levels of ESR (56.0, IQR 44.0–82.5; 26.0, IQR 20.0–56.5, p=0.031) and CRP (1.3, IQR 0.7–2.6; 0.2, IQR 0.1–2.1, p=0.019). PET parameters were significantly reduced, including maximum standardised uptake value (3.50, IQR 3.10–3.84; 3.10, IQR 2.49–3.27, p=0.023), target-to-vein ratio (1.34, IQR 1.13–1.95; 1.31, IQR 1.05–1.45, p=0.032), and target-to-liver ratio (2.38, IQR 1.47–3.05; 1.92, IQR 1.51–2.18, p=0.014) from baseline to week 30. Serum levels of pentraxin-3, soluble human leukocyte antigen-E (sHLA-E), interleukin-6 tended to decrease, while tumour necrosis factor-α level increased after IFX therapy. There were no serious adverse events (SAEs) or AEs necessitating discontinuation of IFX.

Conclusions: Treatment with IFX may lead to remission or improvement with lower glucocorticoid requirement in TAK (clinicaltrials.gov NCT02457585).

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3681