AB1111

ARTHRITIS IS NOT A PREREQUISITE DISEASE MANIFESTATION FOR THE DIAGNOSIS OF SYSTEMIC JIA: RESULTS OF A PROSPECTIVE COHORT TRIAL USING RIL-1RA AS FIRST LINE TREATMENT WITH LONG TERM FOLLOW-UP

N.M. Terhaar¹, A. Leek¹, A. El Idrissi¹, J.F. Swart¹, J. van Loosdregt², N. Wulfraat¹, S. de Rook¹, S. Vastert¹. ¹*Pediatric Rheumatology and Immunology*; ²*Laoratory for Translational Immunology, University Medical Center Utrecht, Utrecht, Netherlands*

Background: Systemic onset juvenile idiopathic arthritis (sJIA) is a multifactorial disease, characterised by arthritis, spiking fever, skin rash, lymphadenopathy, hepatosplenomegaly and/or serositis, in combination with increased inflammatory parameters as erythrocyte sedimentation rate (ESR), C-reactive protein (CRP) and ferritin. sJIA is nowadays seen as a complex autoinflammatory disorder. However, in the current ILAR classification, sJIA is still classified under the umbrella of JIA. The past decade has learned that the mechanisms underlying the systemic inflammation in sJIA differ in important aspects from the other subtypes like polyarticular JIA.

Objectives: Here we compare disease characteristics, manifestations and response to treatment of ILAR-criteria fulfilling sJIA (n=30) and 'sJIA without arthritis' (n=12), in order to evaluate whether arthritis should still be a prerequisite for the diagnosis of sJIA.

Methods: We included 30 consecutive diagnosed and prospectively followed new onset sJIA patients as well as 12 'sJIA without arthritis' from our paediatric rheumatology clinic from 2008 until 2017. The 'sJIA without arthritis' patients underwent extensive diagnostic procedures to exclude infections (PCR, blood cultures, serology etc), malignancies (bone marrow punctures, PET scans etc) and other diagnoses.

All patients followed a standardised treatment protocol, starting with rIL-1RA (2 mg/kg) as 1st line treatment (without steroids), as previously described.¹ In case of partial response, rIL-1RA dose was raised to 4 mg/kg with a maximum of 200 mg/day. If that failed, corticosteroids were added and/or patients switched to alternative biologicals as canakinumab or tocilizumab. If patients had inactive disease at 3 months after start of rIL-1RA treatment, rIL-1RA was tapered for a month (alternate day regimen) and subsequently stopped.

Peripheral blood samples were taken before initiation of rIL-1RA and at all follow-ups for routine lab measurements. For biomarker analyses, serum was isolated at disease onset and at three months after initiation of rIL-1RA.

Results: There were no differences in disease manifestations like skin rash, serositis, hepatosplenomegaly or symptoms like arthralgic (painful) joint count between sJIA and 'sJIA without arthritis' patients at diagnosis. Nor was there a difference in the levels of CRP, ESR, ferritin or IL-18 at start of therapy.

Importantly, also the response to rIL-1RA treatment did not differ between sJIA and 'sJIA without arthritis' patients in our cohort. At last follow-up (median 5.8 years, IQR 2.9–7.6 years), 95% of patients had inactive disease, of which 72% off medication.

Conclusions: Based upon disease manifestations and inflammatory parameters in patients with confirmed sJIA and 'sJIA without arthritis' at disease onset and on excellent treatment responses to a standardised treatment protocol with rIL-1RA as 1st line treatment, we conclude that arthritis should not be a prerequisite disease criterium in the next classification criteria of sJIA.

REFERENCE:

[1] : Vastert, SJ et al, Arthritis Rheumatology 2014. doi: 10.1002/art.38296.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7392

AB1112

AUTOANTIBODIES IN CHILDREN WITH JUVENILE ARTHRITIS ON BIOLOGICAL THERAPY

O.A. Oshlyanska. *connective tissue disorders in children, State Institute of Pediatrics, Obstetrics and Gynecology, Academy of Medical Sciences of Ukraine, Kyiv, Ukraine*

Background: Antinuclear antibodies (ANA) are described as initial serological findings in juvenile idiopathic arthritis (JIA). SLE-like syndromes with ANA seroconversion have been described in the patients on biological therapy, including JIA children. Biologicals have been used in Ukraine since 2011. In 459 of 2,542 JIA cases registered in 2016, patients were on biologicals. According to the data of the Ukrainian Registry of Patients Receiving Biologicals, autoimmune diseases as complications of biological therapy have not been registered in Ukraine so far.

Objectives: To study the ANA dynamics in JIA children on biologicals.

Methods: There was conducted a retrospective observational study which involved 138 JIA patients, including 44 children with the biologicals as follows: TOZ 12, ADA 29, ETA 3. JIA markers (HLA B27 (PCR), RF (ELISA), aCCP

(immunosorption), ANA (IFT and ELISA (10 subtypes) were determined in patients at the disease onset and after 1, 2, 5 years of biologics medication.

Results: Among all JIA patients, ANA were detected in 54% of oJIA, 47% of pJIA, 7% of all spondyloarthritis (spA) cases. At the beginning of biologics, only 22% children were ANA-positive, 66% of them being female, mean age at the disease debut was 6.9±3.81 y. The median of the time from the disease onset to the initiation of biologics was 4.2±3.2 y. The mean age at biologic therapy start was 10.3±4.18 y. At the time of the examination, the mean duration of biologics was 2.4±1.46 y. In pJIA onset ANA were detected in 62% of patients, in oJIA with uveitis in 75%, in spA in 14.2%; in sJIA ANA were not identified. RF and aCCP were found in 2 pJIA children; HLA B27 was detected in 1 child with uveitis and in 64.2% of patients with spA. The data obtained showed that ANA were found more often in patients on biologics than in general JIA population (pJIA, p=0.04, oJIA, p=0.02), RF was lower (4.5%) (p=0.001). The relationship with the sex was not seen (r=0.19). 21.4% of patients had been ANA positive before ADA therapy initiation. Reverse seroconversion was seen in most patients after a year of ADA therapy (3.5% remained ANA+, reduced ANA titer). After 2 year medication, ANA were detected by IFT in 3 seronegative at the disease debut patients (7.1%). They had negative antibodies to histones and aDNAII (ELISA). After 5 y, 2 patients on ADA remained ANA+. Among patients on ADA, who had ANA as a new phenomenon there were 2 female and 1 male teens in disease remission. They had no clinical manifestations of secondary autoimmune pathology. All the patients continued methotrexate, corticosteroids were discontinued in 2 of them. All patients were on isoniazid medication for over a year. In RF +patients on ADA, the RF was not detected a year later, after 2 y it was revealed in 1 pJIA patient who was RF-neg (aCCP neg). There was 1 RF +case at pJIA debut with TOZ, after the therapy it was not detected. Two paediatric patients on TOZ (16.6%) were ANA +at the debut, but the subsequent dynamics studies did not detect ANA (TOZ vs ADA, p=0.04). Patients treated with ETA were observed for less than 1 year.

Conclusions: ANA types which are detected at JIA debut and after initiation of biologic treatment may differ. It is suggestive of the biologics potential to modify the immune response, thus increasing the risk of overlap-syndromes. Therefore, it is advisable to monitor autoantibody titers in JIA children on biologic treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3094

AB1113

PREVALENCE OF GENERALISED JOINT HYPERMOBILITY IN THE CHILDREN POPULATION OF ORDU; TURKISH STUDY

Ö.F. Barçak^{1*}, S.S. Dedeoğlu². ¹*Physical Medicine and Rehabilitation, Özel Ünye Çakırtı Hastanesi, Ordu*; ²*Orthopedics and Traumatology, Okmeydanı Training and Research Hospital, İstanbul, Turkey*

Background: Generalised Joint hypermobility (GJH) is a clinical conditions that may cause common musculoskeletal pain during childhood. In our study, we aim to show the prevalence of GJH in children of 11–18 age group, and to provide guiding information for rheumatologists and paediatricians who are interested in this subject.

Objectives: Our study aimed to evaluate the frequency of GJH in children aged 11–18 years in the province, Ordu.

Methods: This cross-sectional study was performed with 410 students aged 11–18 years who receive education in the province, Ordu. Questionnaire forms were filled in, and each student was examined. The children, who reported to have any disease, were excluded from the study. GJH was diagnosed according to criteria of Beighton diagnosis.

Results: A total of 410 students, 210 of whom were girls (51.3%) and 200 (48.7%) of whom were boys, participated in the study. The subjects' mean age was 13.7±1.7 years for girls and 13.1±1.79 years for boys. The body mass index (BMI) of the girls was 21.5±3.4 kg/m² and of the boys were 22±3.8 kg/m². 160 (39%) of the students participated from the city centre and 250 (61%) from the district centres. The presentations of the students to the health institution due to any complaint in 1 year were examined. The students participating in the study were questioned in terms of presence and time of previous joint complaints. Accordingly, the number of participants who previously had a joint-related complaint was found to be 155 (37.8%). 40 (10.7%) of these participants had a joint-related complaint 3 months ago, 18 (4.3%) had it 6 months ago, 40 (9.7%) had it 1 year ago, and 47 (11.4%) had it more than one year ago. The frequency of GJH was 8.7%. 24 of 36 participants in whom GJH was detected and had a Beighton score of 5 and above consisted of girls; and this was 11.4% of the girls. The number of male participants in whom GJH was detected, was found to be 12; and this was 6% of the boys. There was a significant difference between female and male participants in terms of the frequency of GJH (p=0.021). A statistically significant and highly negative correlation was found between age and Beighton score (r=−0.182, p<0.001). A statistically significant and highly negative correlation was found between body mass index and Beighton score (r=−0.092, p<0.05).