AB1175

DEVELOPMENT AND PRELIMINARY VALIDATION OF AN OMERACT MRI ENTHESITIS SCORING SYSTEM FOR THE ANKLE IN SPONDYLOARTHRITIS

A.J. Mathew¹, S. Krabbe², F. Gandjbakhch³, R.G. Lambert⁴, K.-G. Hermann⁵, I. Eshed⁶, P. Bird⁷, W.P. Maksymowych⁸, J.-D. Laredo⁹, S.J. Pedersen², M. S. Stoeniou¹⁰, D. Glinatsi¹¹, I. Haugen¹¹, V. Foltz¹², J. Jaremko⁴, R.P. Poggenberg², P.G. Conaghan¹³, M. Østergaard¹⁴, on behalf of OMERACT MRI in arthritis working group. ¹Department of Clinical Immunology and Rheumatology, Christian Medical College, Vellore, India; ²Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Rigshospitalet, Copenhagen, Denmark; ³Service de rhumatologie, Hôpital Pitié-Salpêtrière, Paris, France; ⁴Department of Radiology and Diagnostic Imaging, University of Alberta, Edmonton, Canada; ⁵Department of Radiology-Arthritis Imaging Research Group, University Hospital Charité – Campus, Berlin, Germany; ⁶Department of Diagnostic Imaging, Sheba Medical Center, Affiliated to the Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel; ⁷Division of Medicine, University of New South Wales, Sydney, Australia; ⁸CaRE (Canadian Research Education) Arthritis and Department of Medicine, University of Alberta, Alberta, Canada; ⁹Service de Radiologie, Hôpital Lariboisière, APHP and Université Paris-Diderot, Paris, France; ¹⁰Cliniques Universitaires Saint-Luc, Institut de Recherche Expérimentale et Clinique (IREC), Université catholique de Louvain, Brussels, Belgium; ¹¹Department of Rheumatology, Diakonhjemmet Hospital, Oslo, Norway; ¹²Service de Rhumatologie, Hôpitaux Universitaires, Paris, France; ¹³Musculoskeletal Biomedical Research Unit, University of Leeds, Leeds, UK; ¹⁴Copenhagen Center for Arthritis Research (COPECARE) – Center for Rheumatology and Spine Diseases, University of Copenhagen, Rigshospitalet, Copenhagen, Denmark

Background: Enthesitis is regarded the primary lesion in spondyloarthritis (SpA) and MRI allows sensitive visualisation of enthesal inflammation/damage, but no validated, internationally accepted scoring system exists.

Objectives: To develop and perform preliminary validation of a novel OMERACT MRI scoring system for assessing ankle enthesitis in SpA patients, and to improve this through iterative multi-reader scoring exercises and calibration sessions.

Methods: A systematic literature review of MRI studies on enthesitis in SpA identified key inflammatory and structural pathologies. Definitions were agreed by consensus within the OMERACT MRI in arthritis working group. Then, in a first internet-based multireader exercise the Achilles tendon and plantar fascia entheses in 10 ankle MRIs (sagittal T1W, sagittal and Axial T2W fat suppressed) were scored by 15 readers (3 radiologists and 12 rheumatologists), with varying expertise in ankle MRI, for tendon/fascia thickness/signal change, tendon/peritendon signal alteration, retrocalcaneal bursitis, bone spur, erosion and bone marrow oedema, using semi-quantitative scores (0–3: no/mild/moderate/severe pathology). After a subsequent calibration session leading to minor modifications of assessed parameters, 16 ankle MRIs (specifications as above), were scored by 15 readers in exercise 2 with a modified score sheet. Rules were agreed for scoring pathologies. In both exercises, scores for each reader for individual variables were compared and discussed, and mean scores for each variable were determined using descriptive statistics, as were sum scores. Inter-reader agreement was calculated using two-way consistency single measures intra-class correlation coefficient (ICC 3,1) for inflammatory and structural lesions.

Results: Exercise 1: Mean pairwise inter-reader ICC for combined score of inflammatory and structural variables was 0.65 (range 0.10–0.94), with 75% of values being good/very good (≥ 0.50). Discussion of results led to minor modifications of parameters to be assessed.

Exercise 2: Inter-reader agreement (ICC) for all inflammatory variables combined ranged from 0.26–0.93 among reader pairs (mean 0.64; median 0.66; IQR 0.46–0.79). For structural variables combined ICC ranged from 0.05–0.91 among reader pairs (mean 0.45; median 0.45; IQR 0.2–0.6).

Conclusions: Initial steps in developing an OMERACT MRI heel enthesitis scoring system have demonstrated overall moderate reliability of the proposed variables. Further modification, refinement, calibration and validation (ongoing) are needed before this system is ready for use in SpA clinical trials.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5000

AB1176

ULTRASOUND EVALUATION OF ADHESIVE CAPSULITIS OF THE SHOULDER. DESCRIPTION OF A NEW AND SIMPLE DIAGNOSTIC SIGN

C. Moragues Pastor^{1,2}, X. Michelena², I. Morales², P. Juárez², J. Lluch², J. M. Nolla². ¹Rheumatology, Hospital Plató, Barcelona; ²Rheumatology, Hospital Universitari de Bellvitge, L'Hospitalet del Llobregat (Barcelona), Spain

Background: Adhesive capsulitis of the shoulder (ACS) is a common disease that is characterised by a global and progressive loss of mobility of the glenohumeral joint. It is the result of a capsular inflammation, thickening, and fibrosis with subsequent retraction of the joint capsule. Although it is a frequently self-limiting process, it may persist for years and be resolved with a permanent limitation of

glenohumeral mobility. Its diagnosis is based on clinical presentation because the glenohumeral capsule is not visible in simple radiology and the MRI does not offer specific diagnostic findings for ACS.

Objectives: To evaluate the specific ultrasound signs to diagnose ACS in patients with clinical suspicion of ACS.

Methods: Prospective, non-interventional observational study in consecutive patients with clinical suspicion of ACS referred for an ultrasound evaluation. All patients signed an informed consent before the study. Adult patients with a painful shoulder less than 2 years of evolution with clinical suspicion of ACS were included. We excluded patients with surgery or severe articular trauma in the symptomatic shoulder and patients with ultrasound findings that could explain the painful symptoms of the shoulder due to other causes such as rotator cuff tendinopathy, synovitis or glenohumeral osteoarthritis.

The ultrasound examination was systematically performed by an expert sonographer, using an Esaote MayLab 70 with a linear probe. In the ultrasound examination, the measurement of the joint capsule was performed in the axillary recess in the longitudinal plane (maximum passive abduction of the shoulder). The statistical analysis was carried out comparing the capsule means with the Mann-Whitney test (the variable did not follow a normal distribution verified with the Kolmogorov-Smirnov test).

Results: A total of 35 patients were included, with a mean age of 58.27 years, 46–77 17 men and 18 women. Nine patients had both shoulders affected, so a total of 70 shoulders (44 affected shoulders and 26 control shoulders) were studied. The mean time of evolution of the ACS clinic was 5.6 months. 1–18 The 28.57% of the patients were diabetic and 14% had a history of severe cardiovascular disease before the onset of symptoms. The 44% of bilateral ACS cases were diabetic. The totality of the affected shoulders presented in the clinical exploration limitation of the degrees of mobility of the glenohumeral joint; on the other hand, all control shoulders had a preserved mobility. Regarding the sonographic findings, the mean axillary capsular measurement of the affected shoulders was 4,414 mm (SD 0.177) compared to a mean measurement of 2,203 mm (SD 0.165) in the controls with a statistically significant difference between both groups ($p < 0.001$).

Conclusions: The glenohumeral capsule is easily visible in the axillary approach by ultrasound and allows thickness measurement, being a feasible test for the diagnosis of ACS.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5982

AB1177

A COHORT OF PATIENTS WITH ANTISYNTHEASE SYNDROME EVALUATED IN A MULTIDISCIPLINARY CONSULTATION

C. Tórnoro¹, L. Nuño¹, G. Bonilla¹, L. Gomez-Carrera², R. Álvarez-Sala², A. Balsa¹. ¹Rheumatology; ²Pulmonology, La Paz Hospital, Madrid, Spain

Background: Antisynthetase syndrome (AS) is an uncommon connective tissue disease characterised by the presence of antibodies to anti-aminoacyl t-RNA synthetase (anti-ARS) along with features of interstitial lung disease (ILD), myositis, arthritis and Mechanic's hands.

Objectives: To analyse the most common demographic, clinical, radiological, spirometry and capillaroscopic findings in a cohort of patients with AS. As a secondary objective, the association between capillaroscopic findings and diffusion lung capacity for carbon monoxide (DLCO) and the radiologic pattern in the high-resolution computed tomography of the chest (HRCT) was evaluated.

Methods: A ten-year (2007–2017) retrospective analysis of patients diagnosed with AS followed in a multidisciplinary consultation with a pulmonologist and a rheumatologist. The nailfolds from 2nd to 5th fingers in both hands were examined in all subjects by using videocapillaroscopy (Mediscope-Optilia).

Results: Twenty patients were included, 15 (75%) females and 50% (10/20) non-smokers. Mean age at the clinical debut was 47.5±16 years. Throughout their evolution, 90% of patients (18/20) presented Interstitial Lung Disease (ILD); 70% (14/20), arthritis and 75% (15/20), myositis. Furthermore, 40% (8/20) associated fever, 45% (9/20), Raynaud's Phenomenon and 55% (11/20), Mechanic's hands. Three patients only had ILD and Raynaud's Phenomenon at presentation. In the immunologic assessment, 73.7% (14/19) were positive for anti-Jo-1 antibodies and 26.3%, for anti-PL-12. As non-anti-ARS antibodies, 6 had positivity for rheumatoid factor; 3, for ACPA and 5, for anti-Ro-52 antibodies. Of those 18 diagnosed of ILD, spirometry tests at baseline were available for 13. It was remarkable that 69.23% (9/13) of patients presented a diminished value of DLCO (low in 23% (3/13), moderate in 46% (6/13) and normal in the rest of subjects). The most common ILD pattern was non-specific interstitial pneumonia (55%, 10/18) followed by usual interstitial pneumonia (33.3%, 6/18) and organising pneumonia (11.1%, 2/18). Nailfold capillaroscopy was performed in 16 patients: Scleroderma pattern was observed in 5 (all of them associating Raynaud's phenomenon) and 6 patients showed microangiopathic changes. The most frequent capillaroscopic findings were neoangiogenesis (93.8%) and microhaemorrhagias^{68,8} followed by avascular areas (37.5%) and megacapillaries (31%). An association between capillaroscopic findings and reduction of DLCO or the radiologic pattern was not observed,

possibly influenced by the number of patients included. One subject associated Sjögren Syndrome, another one cutaneous Lupus and another patient, morphea. All of the patients included, received steroids and immunosuppressants. Azathioprine was the most frequently used (18/20), followed by Methotrexate (7/20), Rituximab (5/20), Mycophenolate (4/20), Cyclophosphamide (3/20) and Abatacept (1/20).

Conclusions: In our cohort with AS, the most common clinical manifestation was ILD, and an important part of the patients showed a decreased DLCO. Capillaroscopy was pathogenic in most of them and could be useful for the diagnosis of AS, specially in those patients presenting solely with ILD at the clinical debut, although its prognostic value is still unknown. Multidisciplinary consultations are useful in the management of these truly challenging patients.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4977

AB1178 IS ANTI NUCLEAR ANTIBODY TEST REQUIRED CORRECTLY IN DAILY PRACTICE? A CHOOSING WISELY ANALYSIS

D. Üsküdar Cansu¹, H. Üsküdar Teke², E. Gündüz², C. Korkmaz¹. ¹Internal Medicine, Rheumatology; ²Internal Medicine, Hematology, Eskişehir Osmangazi University, Eskişehir, Turkey

Background: The most sensitive and gold-standard test used for the detection of antinuclear antibody (ANA) is accepted as the indirect immunofluorescence (IIF) test.¹ International proposals, including the choice of rational examination, referred to as "choosing wisely", include significant efforts to reduce overuse of examinations such as the ANA.²

Objectives: Our aim was to retrospectively analyse the distribution of the ANA test samples for clinical reasons according to the clinical departments and to determine the most frequently requested reasons and to draw attention to the rational ANA request.

Methods: Detection of ANA samples. Between January 2014 and August 2016, the results of the requested ANA test samples were screened for various reasons. The results of the first 4 divisions with the most common ANA test request and the reasons for the test were also analysed in detail (paediatric diseases were excluded). The ANA test was evaluated using the IIF method and the interpretation of the results was done according to the fluorescence intensity observed in the samples (+1 positive, +2 positive, +3 positive and +4 positive).

Results: 11407 ANA samples were evaluated. The first 5 sections requiring ANA test most were as follows: Neurology 1917 (24.7%), dermatology 1825 (23.6%), rheumatology 1574 (20.3%), paediatric nephrology 1404 (18.1%) and haematology 1027 (13.3%). In all the sections dealing with children, the ANA test number was 2461 and the mean age of the patients was 12.7 years, while the number of ANA samples required in the entire adult patient group was 8946 and the mean age of the patients was 46.8 years. ANA positivity rates for neurology, dermatology, rheumatology and haematology departments are 14.9%, 15.4%, 32.6% and 14.1%, respectively. The interclinical positivity rate was statistically significant ($p < 0.0001$). Positive +1, +2, +3 and +4 positivity rates were 33.5%, 27.8%, 22.2% and 16.3%, respectively, when all positive ANA results were evaluated in terms of titre. When positivity was evaluated in terms of ANA positivity rates/titre, +4 positivity was found to be the most common in the patients from the rheumatology department (135/262=51.5%) ($p < 0.0001$).

The reasons for ANA requests were determined according to ICD diagnostic codes. According to this; in neurology, headache, cerebrovascular event and multiple sclerosis; in dermatology urticaria, psoriasis and dermatitis; in rheumatology, SLE, arthritis and fibromyalgia; in haematology, anaemia and thrombocytopenia were the most common reasons for ANA requests.

Conclusions: Based on specific recommendations and "choosing wisely", the ANA test should be requested in patients with a high pre-test probability for autoimmune diseases. Raising awareness in terms of wisely choosing ANA test is needed in departments outside the rheumatology.

REFERENCES:

- [1] Meroni PL, Schur PH. ANA screening: an old test with new recommendations. *Ann Rheum Dis*. 2010;69:1420–2.
- [2] Lesuis N, Hulscher ME, Piek E, Demirel H, van der Laan-Baalbergen N, Meek I, van Vollenhoven RF, den Broeder AA. Choosing Wisely in Daily Practice: An Intervention 3. Study on Antinuclear Antibody Testing by Rheumatologists. *Arthritis Care Res (Hoboken)*. 2016;68:562–9.

Acknowledgements: None

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4168

AB1179 MUSCULOSKELETAL PATTERN OF 18F-FDG UPTAKE IN PATIENTS WITH POLYMYALGIA RHEUMATICA

D. Prieto-Peña¹, J. Loricera¹, I. Martínez-Rodríguez², J. Banzo², M. Calderón-Goercke¹, V. Calvo-Río¹, C. González-Vela³, M. González-Gay¹, J. Hernández⁴, R. Blanco¹. ¹Rheumatology; ²Nuclear Medicine; ³Pathology; ⁴Internal Medicine, Hospital Universitario Marqués de Valdecilla, IDIVAL, Santander, Spain

Background: Polymyalgia rheumatica (PMR) is a common inflammatory rheumatic disease of the elderly whose diagnosis is usually based on clinical and ultrasound findings. Recently, ¹⁸F-FDG PET/CT has been proposed as a promising one-step tool for assessing extent and severity of PMR. However, the pattern of ¹⁸F-FDG uptake in PMR is not well established and there is a lack of imaging guidelines.

Objectives: Our aim was to describe the musculoskeletal pattern of ¹⁸F-FDG uptake in PMR patients and assess if there were any differences between classic and atypical PMR.

Methods: Retrospective study of 75 patients with PMR and their respective PET/CT scans from a referral centre. We considered two groups: **a)** Classic PMR: patients who fulfilled the 2012 EULAR/ACR criteria; and **b)** Atypical PMR: patients with symptoms resembling PMR but did not fulfil the 2012 EULAR/ACR criteria. Distributions of categorical variables were compared by Pearson Chi2 or Fisher exact test as appropriate.

Results: We evaluated 75 patients (27 men and 48 women) with a mean age $\pm$ SD of 68.2 ± 10.7 years. A PET/CT was performed in all of them. Forty-two (56%) patients classic PMR and 33 (44%) atypical PMR. FDG-PET uptake was observed in the following musculoskeletal regions: in shoulders (n=45), sternoclavicular joints (n=33), hips (n=32), cervical interspinous bursae (n=8), lumbar interspinous bursae (n=29), pubic symphysis (n=4), subtrochanteric bursae (n=20), ischial tuberosities (n=19) and knees (n=33). The comparative study between both groups is shown in the **TABLE**, without observing any statistical significance.

Abstract AB1179 – Table 1

	PMR (N=75)	Classic PMR (n=42)	Atypical PMR (n=33)	P#
Sex (women), n (%)	48 (64.0)	29 (69.0)	19 (57.6)	0.30
Age (years), mean $\pm$ SD	68.2 $\pm$ 10.7	73.9 $\pm$ 8.5	61.6 $\pm$ 9.5	0.60
F-FDG uptake		25 (59.5)	20 (60.6)	0.92
Shoulders	45 (60.0)	20 (47.6)	13 (39.4)	0.48
Sternoclavicular joints	33 (44.0)	19 (45.2)	13 (39.4)	0.76
Hips	32 (43.7)	5 (12.0)	3 (9.0)	0.98
Cervical interspinous bursae	8 (10.7)	12 (28.6)	11 (33.3)	0.40
Lumbar interspinous bursae	29 (38.6)	1 (2.4)	3 (9.1)	0.31
Pubic symphysis	4 (5.3)	12 (28.6)	6 (18.2)	0.67
Subtrochanteric bursae	20 (26.7)	12 (28.6)	7 (21.3)	0.47
Ischial tuberosities	19 (25.4)	16 (38.1)	17 (51.5)	0.25
Knees	33 (44.0)	25 (59.5)	20 (60.6)	0.92

Comparisons between classic and atypical PMR.

Conclusions: In patients with PMR, ¹⁸F-FDG uptake seems to be more frequent in shoulders, sternoclavicular joints, hips and knees. In addition, ¹⁸F-FDG uptake can be also detected in lumbar interspinous bursae and less frequently in subtrochanteric bursae, ischial tuberosities, cervical interspinous bursae and pubic symphysis. No differences between classic and atypical PMR patients were seen.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4961

AB1180 THE INFLUENCE OF MECHANICAL STRESS ON THE HANDS ON ULTRASOUND RESULTS: A PROSPECTIVE STUDY WITH VOLLEYBALL PLAYERS (US-VOLLEY) – PRELIMINARY DATA

D. Scheicht¹, S. Zeglam¹, S. Frechen², J. Strunk¹. ¹Klinik für Rheumatologie, Krankenhaus Porz am Rhein; ²Independent researcher, Köln, Germany

Background: Various imaging studies have found an influence of physical exercise on joint and tendon pathology in ultrasound. It is not clear¹ how specific grey scale and power Doppler changes are in a context of mechanical stress and² if a distinction to findings in acute arthritis exists.

Objectives: To examine via grey scale and power Doppler ultrasound the acute physiological effects of increased physical stress on the fingers and wrists and their surrounding structures through volleyball training in amateur healthy subjects as a reference to the known pathological findings (effusion, increased perfusion) in acute arthritis.

Methods: Examination of 18 healthy amateur volleyball players via musculoskeletal ultrasound before and directly after their routine volleyball practice. Ultrasound assessment included a grey scale and power Doppler examination of the dominant hand (longitudinal planes of the dorsal wrist and dorsal and palmar