MExNvoxels.² Correlation between all clinical scores and all imaging parameters was done using Spearman rho, with significance levels of $p < 0.05$.

Results: The imaging markers of perfusion in the synovium of the knee (MExNvoxels and IREXNvoxels) were the only imaging measures, which showed a very high association with CRP in both RF + RA ($r = 0.92/0.97$, $p < 0.05$) and PsA patients ($0.93/0.99$, $p < 0.05$), whereas all other imaging markers of inflammation showed no statistical association with blood levels of CRP in these diseases. We found no association between CRP and any imaging assessed scores of inflammation in either RF- RA or OA. In addition, only RF + RA patients showed a positive moderate to high association between MExNvoxels and IL-6 ($r = 0.66$, $p < 0.05$) in the knee joint aspirate.

Conclusions: Quantitative imaging and blood biomarkers of inflammation, such as DCE-MRI parameters and CRP, appear to relate differently to each other in the four most common knee arthritic diseases, RF + RA, RF- RA, PsA and OA. DCE-MRI may have specific utility in differentiating these conditions and their disease activity.

REFERENCES:

- [1] Guermazi A, Roemer FW, Haugen IK, et al. MRI-based semiquantitative scoring of joint pathology in osteoarthritis. *Nat Rev Rheumatol*. 2013;9(4):236–5
- [2] Kubassova O et al; *Eur J Radiol*. 2010 Jun;74(3):e67–72.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6271

AB1207

CORRELATION BETWEEN CLINICAL FINDINGS AND ULTRASONOGRAPHY IN EVALUATING PAINFUL RHEUMATOID SHOULDER

M. Abdel Zaher¹, S. Tharwat¹, A. Abdel khalek², A. Abdel salam¹. ¹Internal Medicine (Rheumatology, Immunology unit); ²Radiodiagnosis department, Mansoura University Hospital, Mansoura, Egypt

Background: Shoulder pain is one of the most common complaints encountered in patients with rheumatoid arthritis (RA) ¹. In recent years ultrasonography (US) became an essential tool in diagnosing rheumatic diseases ². As clinical examination often does not allow an exact diagnosis, the agreement between both methods needs to be discussed to know how much the US add benefits to clinical examination.

Objectives: To determine the agreement between clinical examination and ultrasound in evaluating shoulder pain in rheumatoid.

Methods: A cross-sectional study including thirty RA patients, meeting the ACR/EULAR classification criteria for RA complaining of shoulder pain. They were recruited from rheumatology outpatient clinic in Mansoura University Hospitals. The sixty shoulders of the thirty patients were examined clinically by inspection, palpation and special tests, then fully examined by ultrasound including biceps tendon, subacromial bursa, rotator cuff tendons and acromioclavicular joint.

Results: Agreement among clinical examination and US was examined using Cohen's kappa. There was slight agreement between clinical examination and US regarding biceps tenosynovitis with $k = 0.206$, fair agreement regarding acromioclavicular osteoarthritis with $k = 0.392$ and SASD bursitis with $k = 0.233$. There was also moderate agreement between clinical examination and US examination of the shoulder in case of supraspinatus tendinopathy with $k = 0.464$. Data were statistically significant ($p < 0.001$). The overall agreement between clinical examination and ultrasound was poor.

Conclusions: Clinical examination of shoulder pain in rheumatoid arthritis is not accurate, insufficient. It should be confirmed with US examination during the initial evaluation of the shoulder to give reliable data and differentiate between different pathologies.

REFERENCES:

- [1] Amin M, Ismail F, El-shereef R. The Role of Ultrasonography in Early Detection and Monitoring of Shoulder Erosions and Disease Activity in Rheumatoid Arthritis Patients; Comparison with MRI Examination. *Acid Radiol* 2012;19(6):693–700.
- [2] Plaza M, Nowakowska-Plaza A., Pracon G., Sudol-szopinska I. Role of ultrasonography in the diagnosis of rheumatic diseases in light of ACR/EULAR guidelines. *Journal of ultrasonography* 2016;16:55.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4714

AB1208

THE ROLE OF ULTRASOUND IN THE DISCOVERY OF INTERSTITIAL CHANGES IN THE LUNGS IN PATIENTS WITH SYSTEMIC CONNECTIVE TISSUE DISEASE

N.L. Roganovic, M. Grujic, G. Radunovic. *Institute of rheumatology, Belgrade, Serbia*

Background: Interstitial lung disease (ILD) is one of the most serious lung complications in patients with connective tissue disease (CTD), most commonly in patients with systemic sclerosis (SSc). High-resolution Computed Tomography is a gold standard for assessing ILD. Ultrasound examination of the lungs has been increasingly used lately to evaluate the existence of interstitial changes (B lines) in the lungs.

Objectives: To determine whether there is a significant difference in the presence of B lines on lung ultrasound examination in patients with SSc compared to patients with other CTD and healthy controls. Also, to investigate if there is a significant difference in the presence of B lines on lung ultrasound examination in patients with diffuse SSc compared to patients with limited SSc.

Methods: The study included 150 people of both sexes, aged between 19 and 81, who were examined at the Institute of Rheumatology in Belgrade. In the first group there were 55 patients with SSc (28 with diffuse and 27 with limited form of SSc), in the second group 45 patients with other CTD (16 with rheumatoid arthritis, 16 with systemic lupus erythematosus and 13 with Sjogren syndrome) and in the third group 50 healthy subjects who were matched by gender and age with other two groups. At the ultrasound examination, the number of B lines was determined in all segments of the lungs. A positive ultrasound finding was considered to be one with 3 or more B-lines in at least two adjacent ultrasound scanning fields or one with more than 5 B-lines in any single field of ultrasound scanning.

Results: There was statistically significant difference in positive findings between the group of subjects with SSc and the group of healthy subjects (65,5% vs. 2%; $p < 0.001$) and between the group of subjects with SSc and the group with other CTD (65,5% vs. 13,3%; $p < 0.001$). Also, it was shown that there was statistically significant difference in positive findings between the group of subjects with diffuse SSc and the group with limited SSc (85,7% vs. 44,4%; $p < 0.001$).

Conclusions: The conducted study confirmed that the presence of B lines on lung ultrasound examination is significantly more frequent in patients with SSc, especially in the patients with diffuse SSc.

REFERENCES:

- [1] Shahin AA. Pulmonary involvement in systemic sclerosis. *Treat Respir Med* 2006;5:429–36.
- [2] Sperandeo M, Varirale A, et al. Transthoracic ultrasound in the evaluation of pulmonary fibrosis: our experience. *Ultrasound Med Biol* 2009; 35: 723–9.
- [3] Gargani L, Doveri M, et al. Ultrasound lung comets in systemic sclerosis: a chest sonography hallmark of pulmonary interstitial fibrosis. *Rheumatology (Oxford)* 2009;48:1382–7.
- [4] Barskova T, Gargani L, et al. Lung ultrasound for the screening of interstitial lung disease in very early systemic sclerosis. *Ann Rheum Dis* 2013; 72: 390–395.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7592

AB1209

USEFULNESS OF IMAGENOLGY TO DIFFERENTIAL DIAGNOSIS IN PATIENTS WITH PRESUMED SERONEGATIVE RHEUMATOID ARTHRITIS AND OTHER ARTHROPATHIES

P. Santos-Moreno¹, O. Valencia², E. Castillo¹, S. Farietta¹, L. Villareal¹, M. Cabrera¹. ¹Biomab Center of Rheumatoid Arthritis; ²Universidad de los Andes, Bogotá, Colombia

Background: It is difficult to make a differential diagnosis between seronegative RA and other inflammatory arthropathies. Many patients could be wrong diagnosed followed of expensive treatments.

Objectives: To assess the usefulness of X-rays of hands and feet (X-rays), Ultrasound (US) and Magnetic Resonance Imaging (MRI) to discard false positive diagnosis of seronegative RA from real-world evidence.

Methods: An analysis from medical records of patients with presumptive seronegative RA diagnosis reportedly seronegative for both rheumatoid factor and anti-cyclic citrullinated peptide antibodies and clinical criteria of RA, in the period between July 2016 and ; June of 2017 who were assessed by imagenology (X-rays, US or MRI) in a centre of rheumatoid arthritis to confirm diagnosis or discard it. Laboratory, and imagenology data was retrospectively analysed and multivariate analysis was performed to determinate the usefulness of imagenology.

Results: 360 patients were received in the centre with presumptive diagnosis of RA in the period, mean of age was 58 years, 80,9% females and 19,1% males. X-

rays shows only 3.59% of patients with erosions, 38.6% were positive for Osteoarthritis (OA), and 58.6% of patients without erosions. From patients without erosions, 36% were confirmed for RA and 13.4% for OA by US; on the other hand 37% was confirmed for RA and 20% for OA by MRI. A total of 94 patients (26.1%) had a final diagnosis of seronegative RA, while (261) patients 72.5% were confirmed for Osteoarthritis,⁵ 1.4% were classified in other arthropathies.

Conclusions: According with our findings the screening by imagenology is a useful tool with low cost to make a differential diagnosis of seronegative RA and other arthropathies. In order to improve outcomes through well-defined treatments and to save high cost treatments for pathologies with more precise diagnoses.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7273

AB1210

ANKLE BRACHIAL INDEX FOR THE DIAGNOSIS OF SUBCLINICAL ATHEROSCLEROSIS USING CAROTID INTIMA MEDIA THICKNESS IN SYSTEMIC LUPUS ERYTHEMATOSUS PATIENT IN SANGLAH HOSPITAL

P.K. Kurniari, G. Kambayana, E. Dharmawan. *Internal medicine, Udayana University/Sanglah General Hospital, Bali, Indonesia*

Background: Atherosclerosis and its complications in Systemic Lupus Erythematosus (SLE) patients occurred more rapidly than the general population. Early detection of atherosclerosis is currently a challenge for clinicians. Angiography as a gold standard diagnosis of atherosclerosis is invasive, has a limitation. The surrogate marker carotid intima media thickness (CIMT) examination with B-mode ultrasonography has been used widely and validated. The limitation of this examination is operator dependence. Ankle brachial index (ABI) examination is simpler, cheaper, objective and widely available and is expected to be used for the diagnosis of subclinical atherosclerosis.

Objectives: to determine the sensitivity, specificity, positive predictive value and negative predictive values of the ABI to establish the diagnosis of subclinical atherosclerosis in SLE patients.

Methods: A cross sectional study was enrolling 56 subjects and was conducted from September 2016 to July 2017 at Sanglah Hospital, Denpasar, Bali, Indonesia. We used 2 × 2 cross table to determine the sensitivity, specificity, positive predictive value and negative predictive values of ABI to establish the diagnosis of subclinical atherosclerosis.

Results: Of the 56 samples, 48 people (85.7%) were female. The area under ROC curve was 0.708 (70.8%), p=0.041. ABI examination to diagnose subclinical atherosclerosis in patients with SLE with a cutoff value of 0.95 has a sensitivity of 70%, specificity of 76.1%, 38.9% positive predictive value, and negative predictive value of 92.1%. The best cut-off value of ABI as a diagnostic tool for subclinical atherosclerosis in SLE patients is <0.95.

Conclusions: Examination with ABI can be considered as an alternative diagnostic when CIMT is not available. The diagnostic value of ABI is reliable enough for screening and diagnostic confirmation of subclinical atherosclerosis in patients with SLE.

REFERENCES:

- Westerweel PE, Luyten RK, Koomans HA, Derksen RH, Verhaar MC. Premature atherosclerotic cardiovascular disease in systemic lupus erythematosus. *Arthritis & Rheumatism* 2007; 56(5): 1384–1396.
- Cacciapaglia F, Zardi EM, Coppolino G, Buzzulini F, Margiotta D, Arcarese L, et al. Stiffness Parameter, intima-media thickness and early atherosclerosis in systemic lupus erythematosus patients. *Lupus* 2009;18:249–256.
- Wu GC, Liu HR, Leng RX, Li XP, Li XM, Pan HF, et al. Subclinical atherosclerosis in patient with systemic lupus erythematosus: A systemic review and meta-analysis. *Autoimmun Rev* 2016;15(1):22–37.
- Manzi S. Prevalence and risk factors of carotid plaque in women with systemic lupus erythematosus. *Arthritis Rheum* 2007;42:51–60.
- Theodoridou A, Bento L, D'Cruz DP. Prevalence and associations of an abnormal ankle-brachial index in systemic lupus erythematosus: a pilot study. *Ann Rheum Dis* 2003;62:1199–1203.
- Kadarnan JT, Anggriyani N, Wiryan W. Perbandingan sensitivitas dan spesifisitas ankle brachial index dengan carotid intima-media thickness dalam mendeteksi penyakit jantung koroner signifikan. *Jurnal Kedokteran Diponegoro* 2016; 5(4):1111–1124.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3595

AB1211

UTILITY OF PET/CT FOR THE DIAGNOSIS AND DISEASE MANAGEMENT: A STUDY OF 88 PATIENTS FROM AN AUTOIMMUNE DISEASES HOSPITAL UNIT IN A 2-YEAR PERIOD

P. Diéguez¹, A. Argibay, C. Vázquez Triñanes, B. Gimena, M. Estévez, R. Lorenzo, N. Val, M. Freire, A. Rivera. *Systemic Autoimmune diseases and Thrombosis Unit, University Hospital Complex of Vigo, Vigo, Spain*

Background: 18F-Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography (18F-FDG PET/CT) is a non-invasive imaging technique commonly used in clinical oncology. 18 F-FDG accumulation is recognised as useful for diagnosis and monitoring the response to therapy in patients with some inflammatory disorders, but the role of the PET/TC in the management of these diseases is debated.

Objectives: The aim of this study was to investigate the role of 18F-FDG PET/CT in the diagnosis of the disease and assessing disease activity in an autoimmune diseases unit, and to evaluate if the results of this image technique imply a change in clinical management.

Methods: We retrospectively reviewed all 18F-FDG PET/CT requested since August 2015 to August 2017 by our unit. Data collected were: patient demographics, reason for PET request, PET results and change in therapy.

Results: PET/CT were performed in 88 patients and were positive in 68 (77.3%) cases. Patients (49 women/39 men) had a mean age of 58.1±15.7 years (range, 27–92 years). The clinical diagnosis at the moment of ordering the PET/CT were: sarcoidosis (n=45), large-vessel vasculitis (LVV) (n=16), immunoglobulin G4-related disease (IgG4-RD) (n=9), collagen-vascular diseases (CVD) (n=7), mesenteric panniculitis (n=4), myopathy (n=4), polymyalgia rheumatica (PMR) (n=2) and ANCA-associated vasculitis (n=1). (See table 1).

Abstract AB1211 – Table 1

	Sarcoidosis	LVV	IgG4-RD	CVD	Mesenteric panniculitis	myopathy	PMR
n	45	16	9	7	4	4	2
women	27 (60%)	13 (81.2%)	1 (11.1%)	4 (57.1%)	1 (25%)	2 (50%)	1 (50%)
age	50.9 (27–83)	72.5 (29–92)	63.7 (48–80)	53 (30–80)	60.5	66.2	72.5
PET to the diagnosis	9 (20%)	2 (12.5%)	1 (11.1%)	3 (42.9%)	3 (75%)	2 (50%)	2 (100%)
PET to disease activity	36 (80%)	14 (87.5%)	8 (88.9%)	4 (57.1%)	1 (25%)	2 (50%)	0
Positive PET/TC	36 (80%)	11 (68.7%)	6 (66.7%)	6 (85.7%)	3 (75%)	3 (75%)	2 (100%)
Change of therapy	31 (68.9%)	11 (68.7%)	6 (66.7%)	7 (100%)	2 (50%)	3 (75%)	2 (100%)

PET results supported a change in therapeutic management in 71.6% of the cases. In the group of sarcoidosis there was a change in treatment in 68.9% of cases. PET/TC revealed extrapulmonary manifestations in 57.8%. All patients with extracranial manifestations of giant cell arteritis (GCA) showed uptake in PET/TC. That result supported the clinical diagnosis, although negative temporal artery biopsy. PET was useful in the diagnosis and treatment of IgG4-RD in 66.7% of cases. PET scan did not find malignancy in inflammatory myopathies nor mesenteric panniculitis.

Conclusions: PET/TC is an increasingly utilised in our patients with inflammatory disorders as a support for the diagnosis and management. PET/TC detection extrapulmonary sarcoidosis may have therapeutic and prognostic clinical implications. PET may be useful in patients with GCA suspicion with negative biopsy and extracranial symptoms. More studies will be necessary to establish the real role of PET/TC in autoimmune and inflammatory diseases.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6970

AB1212

ANALYTICAL VALIDATION OF AN INTERFERON-INDUCIBLE GENE EXPRESSION KIT AS A POTENTIAL DIAGNOSTIC TEST FOR ANIFROLUMAB

P.K. Brohawn¹, B.W. Higgs¹, S. Patel², A. Moody³, P. Cooper³, K. Ranade¹.

¹MedImmune LLC, Gaithersburg, MD, USA; ²AstraZeneca, Cambridge; ³QIAGEN Manchester Ltd, Manchester, UK

Background: Anifrolumab, a fully human monoclonal antibody that binds to the type I interferon receptor, is in Phase III development for moderate to severe systemic lupus erythematosus (SLE). Patients with greater interferon-inducible gene signatures (IFNGS) have enhanced response to anifrolumab treatment.¹

AstraZeneca and QIAGEN developed an *in-vitro* diagnostic test for interferon-inducible gene expression (IFIGx). Efficacy of anifrolumab will be evaluated in patients with high and low IFNGS.

Objectives: We aimed to analytically validate the IFIGx kit for use in a pivotal clinical study and potentially to support future regulatory submission.

Methods: The IFIGx kit measures expression of four interferon-inducible genes (IF127, IF114, IF114L and RSAD2) compared with three housekeeping genes. Measurements were performed on mRNA extracted from whole blood from adults with SLE. A score was generated that identified patients as "IFNGS test-high" or "IFNGS test-low." Analytical validation involved six studies measuring lot interchangeability, linearity, repeatability, reproducibility, cross contamination, and system verification.

Results: Repeatability was 100%. Reproducibility was >96%. No cross contamination was observed. Results of all studies validated the IFIGx kit (table 1).

Abstract AB1212 – Table 1

Summary of Analytical Validation of Anifrolumab IFIGx Test

Study	Description	Results
Lot interchangeability	Verification that scores and assay Ct values were robust when different lots of kit components were used Acceptance criterion was <0.58 Ct	Lot interchangeability verified Largest observed change=0.16 Ct
Linearity	Verification, using linear and quadratic regression analyses, that mRNA input concentration (10 ng/μL) is in assay's linear range	Linearity verified Change in score over the concentration range on either side of 10 ng/μL=0.0043 Ct
Repeatability	Verification of Dx result repeatability when the same operator tested 60 random samples using the same kit lot and instrument	Repeatability verified Observed repeatability=100%
Reproducibility	Verification of Dx result reproducibility when multiple operators tested 48 random samples using multiple kit lots and instruments at different sites For verification, overall rate of correct calls must be ≥95%	Reproducibility verified Overall rate=99.7% After six samples with values close to the cut-off were added for further confirmation, overall rate=96.5%
Cross contamination	Investigation of inter- and intra-run cross contamination	No cross contamination No cross contamination found in reverse transcription or PCR steps
System verification	Verification of functionality and utility of the IFIGx software and IFIGx assay package	System verification confirmed Software flags produced as expected

Ct, cycle threshold (PCR cycle at which fluorescence rises above background level); Dx, diagnostic; IFIGx, interferon-inducible gene expression; PCR, polymerase chain reaction; mRNA, messenger ribonucleic acid.

Conclusions: The IFIGx kit was shown to be a robust, reproducible diagnostic test for IFNGS. The IFIGx kit has demonstrated value in prior anifrolumab studies,¹ and will be used both for patient stratification in Phase III studies and to support anifrolumab regulatory filings.

REFERENCE:

[1] Furie, et al. *Arthritis Rheumatol.* 2017;69:376–86.

Disclosure of Interest: P. Brohawn Employee of: MedImmune LLC, B. Higgs Employee of: MedImmune LLC, S. Patel Employee of: AstraZeneca, A. Moody Employee of: QIAGEN Manchester Ltd, P. Cooper Employee of: QIAGEN Manchester Ltd, K. Ranade Employee of: MedImmune LLC

DOI: 10.1136/annrheumdis-2018-eular.2547

AB1213

ULTRASONOGRAPHIC EVALUATION OF THE HIP JOINTS IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS AND PSORIATIC ARTHRITIS: ASSOCIATED WITH CLINICAL AND LABORATORY MARKERS, IMAGING AND FUNCTIONAL OUTCOMES

R. Osipyan¹, M. Bogdanova², M. Bilinskaya³, D. Koshurnikov⁴, A. Baev⁵, T. Tamgina⁶. ¹RUDN University; ²FSBI "Polyclinic №1" of the ADP RF; ³PRNM University; ⁴Hospital for veterans of wars №3; ⁵ICDC of PAO "Gazprom"; ⁶JSC "Sogaz – Med", Moscow, Russian Federation

Background: The hip joints are frequently involved in spondyloarthritis (SpA). Should ultrasonography (US) be additionally used for clinical assessment of disease in patients (pts) with SpA.

Objectives: to detect by US synovitis of the hip joints and to analyse the correlation between US finding and magnetic resonance imaging (MRI) and/or X-ray sacroiliitis, clinical and laboratory markers and functional status in pts with axial SpA (axSpA) and psoriatic arthritis (PsA).

Methods: 13 pts (7M/6F) with definite axSpA (7 with ankylosing spondylitis (AS) and 6 with non-radiographic nrSpA:ASAS) and 7 pts (4M/3F) with PsA (CASPAR) of similar age [median (IQR), 44^{35–55} vs 42^{34–56} years] and disease duration [7 (1–10) vs 9^{7–10} years] were enrolled. US examination (Mindray DC-N6, C5–2 MHz probe) included bilateral of the hip joints according to the EULAR US Scanning App (www.eular.org; <http://ultrasound.eular.org>). MRI of sacroiliac joints (SIJ) was performed on Signa HDxT GE, 1.5 T. The bone marrow oedema on MRI (STIR) considered as active MRI sacroiliitis (MRI-SI).¹ X-ray of SIJs (modified New York criteria), functional status (BASFI), disease activity indices (ASDAS-CRP, DAPSA), laboratory tests (CRP, ESR, HLA-B27) were assessed.

Results: History of hip involvement was present in 7/20 (35%) pts who had ≥1 tender hip joint. 9/20 pts (45%) had US – synovitis of the hip joints. All pts were divided into two groups: g1 (n=9) – US synovitis "+" (at least one hip joint), g2 (n=11) – US – synovitis "–". Pts from g1 had higher CRP and BASFI compared to g2 (table 1). Elevated CRP ≥25 mg/L and ESR ≥30 mm/h, as well as MRI- and X-ray -SI were detected more often in g1pts than g2. But other parameters did not differ between the groups (table 1). US – synovitis of the hip joints was positively correlated with CRP (r=0.489, p<0.05), MRI-SI (r=0.688, p<0.05), X-ray-SI (r=0.634, p<0.05) and BASFI (r=0.541, p<0.05).

Abstract AB1213 – Table 1. Association between the US- synovitis of the hip joints and clinical and laboratory markers, imaging and functional outcomes

Parameters	US-synovitis "+" (n=9)	US-synovitis "–" (n=11)	P*
Age, years	44 (26–49)	46 (40–57)	.294
Duration since diagnosis, years	5 (1–10)	9 (7–10)	.551
AS/nrSpA/PsA	4 (57/3 (50)/2 (28.5)	3 (39/3 (50)/5 (71)	.411
Male sex	6 (67)	5 (45)	.456
HLA-B27	6 (67)	5 (45)	.456
CRP, mg/l	38 (6.7–51)	7.7 (3–19.2)	.031
Elevated CRP≥25	6 (67)	1 (9)	.031
ESR, mm/h	30 (16–39)	11 (6–25)	.080
Elevated ESR≥30	5 (55.5)	1 (9)	.027
ASDAS-CRP	3.4 (2.1–4.5)	2.7 (1.7–4.2)	.294
DAPSA	33 (19.6–46.4)	12.4 (12.3–12.4)	.190
BASFI (0–10)	5 (4–6)	3 (1.7–4)	.015
BASFI>4	6 (67)	2 (18)	.030
MRI-SI	9 (100)	5 (45)	<0.001
X-ray SI	9 (100)	5 (45)	<0.001

Numbers are medians (IQR) and n (%); * Mann-Whitney U-test.

Conclusions: US – synovitis of the hip joints was associated with CRP, MRI- and X-ray- SI and severe functional disability in SpA pts. US evaluation of the hip joints may be used for identifying SpA pts with high risk of the disease progression at an early stage.

REFERENCE:

[1] Mandl P. et al. *Ann Rheum Dis* 2015;0:1–13.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6258

AB1214

BONE MINERAL DENSITY AND T-SCORE EVALUATION IN A GROUP OF PATIENTS WITH RHEUMATOID ARTHRITIS USING A NEW ULTRASOUND METHOD: ECHOS – GENODYNAMIC - A PILOT STUDY –

R.D. Decianu¹, M. Bojinca², S. Daia-Iliescu¹, D. Mazilu¹, M.M. Negru¹, A. R. Balanescu¹, D. Opris- Belinski¹, R. Ionescu¹, V.C. Bojinca¹. ¹Internal Medicine and Rheumatology, Sf. Maria Clinical Hospital; ²Internal Medicine and Rheumatology, Dr. I. Cantacuzino Clinical Hospital, Bucharest, Romania

Background: There is a great need of a screening programme for osteoporosis in the general population and even more amongst the rheumatologic patients, with additional risk factors. Although DXA is the gold standard for diagnosing osteoporosis, it is not appropriate for screening, because of its' limitations regarding ionising radiation, but also the massive machine, that needs dedicated spaces and specialised operators, all in all, involving high costs.

Objectives: To apply a new method that integrates ultrasonography with radiofrequency signals from an echographic scan, in order to evaluate the bone mineral