disease activity assessment in RA, allowing visualisation of synovitis. However, few detailed studies or evaluations of clinical parameters related to synovitis in the foot have been performed.

Objectives: This study investigated the association between synovitis in the foot as observed on joint ultrasonography and the degree of disability on the health assessment questionnaire (HAQ) in RA patients.

Methods: The study included 79 outpatients (101 feet) with RA. Patients who had undergone surgery were excluded. The mean age was 66.0 years (24–92 years), and the mean disease duration was 13 years and 5 months (ranging from 1 month to 49 years). Joint ultrasonography was performed by the same examiner using the same diagnostic instrument in the same room. Synovitis was defined as a power Doppler score of $\geq$ Grade 1. The scanning sites were forefoot (metatarsophalangeal joints of toes 1–5), metatarsus (calcaneocuboid and talonavicular joints), and hindfoot (talocrural and subtalar joints, peroneal tendon, posterior tibial tendon). The presence or absence of synovitis was evaluated at each site, and the association with the HAQ score was examined.

Results: Synovitis was detected in the forefoot of 40 feet (39.6%), in the metatarsus of 34 feet (33.7%), and in the hindfoot of 63 feet (62.4%). Patients with synovitis in the forefoot (PD+: 0.41 ± 0.1 , PD–: 0.80 ± 0.1 , $p=0.018$) and hindfoot (PD+: 0.52 ± 0.1 , PD–: 0.86 ± 0.1 , $p=0.043$) had significantly lower HAQ scores.

Conclusions: Given that assessment of foot impairment is not included in the disease activity index for RA, there may be a delay in diagnosis of foot lesions. Our results showed that patients with synovitis in the forefoot and rearfoot had lower HAQ scores (disability measure), suggesting that increased physical activity may be associated with increasing incidence of synovitis in the foot. The ability of RA patients to perform activities of daily living may have improved with advances in pharmacotherapy, leading to increased physical stimulation, resulting in an increased incidence of synovitis of the foot.

REFERENCES:

- [1] Brown AK. Arthritis Rheum 58(10); 2958–67: 2008
- [2] Wakefield RJ. Ann Rheum Dis 63; 382–5: 2004

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6050

AB1222 COMPARISON OF QUANTITATIVE MRI FAT-FRACTION MEASUREMENT IN SIJ JOINT ON DIFFERENT SCANNER PLATFORMS

A. Bainbridge¹, T. Bray², J. Jones³, S. Tansley⁴, N. Fulstow⁴, R. Sengupta⁴, M. Hall-Craggs². ¹Medical Physics and Biomedical Engineering, University College London Hospitals NHS Foundation Trust; ²Centre for Medical Imaging, University College London, London; ³Radiology, Nuffield Health Hospital, Bristol; ⁴Royal National Hospital for Rheumatic Diseases, Bath, UK

Background: Quantitative MRI Proton Density Fat Fraction (PDF) measurement can objectively identify areas of active inflammation and structural damage in spondyloarthritis by identifying oedema and fat metaplasia in bone marrow.¹ PDF is typically implemented as a 3D multi-echo Gradient-echo protocol and all of the major MRI equipment manufacturers have PDF imaging products available that combine a complex multi-peak fat-spectrum model with simultaneous determination and correction for T2* signal decay. However, if the appropriate specialist imaging option has not been purchased, as is commonly the case, then the user may have to rely on simpler base-level protocols for fat-fraction measurement. An important step to implementing quantitative fat-fraction measurement as a clinical tool is to demonstrate that measurements are robust to both scanner platform and measurement technique.

Objectives: We aimed to compare quantitative FF measurements obtained using 3 scanners supplied by different manufacturers, and using a range of techniques.

Methods: The sacroiliac joints (SIJ) of 10 healthy volunteers (mean age 42 years, range 23–63) were scanned on 3 scanners: a) 3T Philips Ingenia, using MDixon Quant (MDQ) – a specialist PDF package; b) 1.5T GE Optima MR450W, using IDEAL and LAVA – both base-level options; and c) 3T Siemens Skyra, using DIXON – a base-level option. We also scanned a fat-water phantom constructed using tubes of emulsions of peanut oil and 3% agar. MDQ uses data from 6 different echo-times; IDEAL from 3; and LAVA and DIXON from 2. The excitation flip angle was set to 3 degrees for all acquisitions to minimise the T1 weighting. Each technique yields Fat-only (F) and Water-only (W) images. FF images were calculated using the formula: $FF = F/(W+F)$. Regions of interest (Rols) were drawn in the SIJ as shown in the figure 1. MDQ PDF was used as the 'gold-standard' measurement for comparison to IDEAL, LAVA and DIXON.

Results: The figure shows the results of the Rol analysis. For the phantom measurements, both LAVA and IDEAL correlated well with MDQ, but Siemens DIXON deviated away from the expected linear relationship. The *in vivo* data showed good linear relationships with MDQ for all 3 base-level techniques for FF in the range 0.4–0.8, although all 3 slightly underestimated FF compared to MDQ.

Abstract AB1222 – Figure 1

Conclusions: A clinically useful quantitative image-analysis tool should be applicable to results obtained from several platforms. FF measurement in the bone marrow of the SIJ is robust to both technique and platform. The slight underestimation of FF values on base-level FF measurements is probably due to the influence of T2* weighting. However, these variations are smaller than the FF changes observed in areas of oedema or fat metaplasia.¹ FF measurement is a quantitative and widely applicable method for the diagnosis and assessment of spondyloarthritis.

REFERENCE:

- [1] Bray TJP, Bainbridge A, Punwani S, Ioannou Y, Hall-Craggs MA. Simultaneous Quantification of Bone Edema/Adiposity and Structure in Inflamed Bone Using Chemical Shift-Encoded MRI in Spondyloarthritis. Magn Reson Med. 2017;

Acknowledgements: Funding: BANNAR; TB:ARUK grant 21369/MHC; UCLH BRC;

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5411

AB1223 ULTRASONOGRAPHY EXAMINATION OF ELBOWS IS USEFUL WHEN COMBINED WITH EXAMINATION OF PIP AND WRIST JOINTS

T. Nakajima¹, M. Shimizu², M. Hashimoto³, H. Yamamoto², R. Kawahara², S. Nakabo¹, R. Hiwa¹, Y. Gon¹, S. Funakoshi¹, Y. Nakamura¹, T. Sasai¹, T. Tomizawa⁴, K. Nishitani⁴, K. Murata⁴, M. Tanaka³, H. Ito⁴, Y. Fujii², T. Mimori¹. ¹Department of Rheumatology and Clinical Immunology, Kyoto University Hospital; ²Department of Human Health Sciences, Kyoto University Graduate School; ³Center for Rheumatic Disease; ⁴Department of Orthopedic Surgery, Kyoto University Hospital, Kyoto, Japan

Background: Because elbows are important joints for the function of upper limbs, the daily activity of the patients with rheumatoid arthritis (RA) may be impaired by synovitis of elbows. Although elbows are included into the 28 essential joints for evaluation of the disease activity of RA, ultrasonography (US) examination of elbows is not routinely performed and there have been a few surveys on the correlation between US scores of elbows and disease activity.

Objectives: To reveal whether there is a correlation between US scores of elbows and disease activity in the patients with RA.

Methods: We recruited 198 patients with RA from KURAMA cohort, which is the database of the patients with RA treated in the Kyoto University Hospital, and performed US examination of proximal interphalangeal (PIP) joints, metacarpophalangeal (MP) joints, wrist joints and elbow joints. Synovial hypertrophy is scored semiquantitatively on a scale of 0–3. We analysed correlations between the total scores of each component (PIP, MP, wrist and elbow) and clinical parameters.

Results: We found weak correlations between DAS28-ESR and the US scores of PIPs, MPs and wrists ($\rho=0.25$, 0.21 and 0.29 , respectively) whereas there was little correlation between DAS28-ESR and the US score of elbows ($\rho=0.16$). However, multivariable analysis showed that the US scores of PIPs, wrists and elbows were independently associated with DAS28-ESR ($p=0.021$, 0.0027 and 0.025 , respectively). US scores of elbows showed little or weak correlations with US scores of PIPs, MPs and wrists ($\rho=0.23$, 0.16 and 0.11 , respectively).

Conclusions: US examination of elbows may be useful to assess the disease activity of patients when combined with US examination of PIPs and wrists although US scores of elbows alone has a weak correlation with DAS28-ESR. US examination of elbows may detect the disease activity which is overlooked by the US examination of joints of hand.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2018-eular.5338

AB1224 CT PET SCANS IN SUSPECTED LARGE VESSEL VASCULITIS AND GIANT CELL ARTERITIS – AN AUDIT IN THE BELFAST HEALTH AND SOCIAL CARE TRUST (BHSC)

U.A. Lavery, E. Banks, M. McHenry. NHS, Belfast, Ireland

Background: British Society of Rheumatology (BSR) guidelines, due to be updated in April 2018, recommend consideration of CT PET when there is suspicion of large-vessel GCA (LV-GCA) in patients with prominent systemic symptoms, limb claudication or persistently high-inflammatory markers despite adequate glucocorticosteroid therapy. Vascular Ultrasound is unhelpful in assessment of the aorta.

Objectives: We investigated the use of CT PET in suspected cases of LV-GCA and its impact on management of patients in the BHSC from August 2016 to August 2017.

Methods: The IT support team in Royal Victoria Hospital provided a list of CTPET scans requested under the specific code for vasculitis and/or Pyrexia/infection and another code for general. Of the 250 scans identified under these codes, 34 scans were requested by Rheumatology for possible vasculitis following a review of the electronic care records. A proforma was used to aid data collection.

Results: Female:Male ratio was 3.25:1, with a mean age of 65. 88% of the scans were requested due to a suspected diagnosis of vasculitis and 12% were for follow up of known vasculitis. 24% of CT PET scans were positive for large vessel vasculitis (LVV). The ESR was greater than 50 mm/hr in 75% of positive scans. Of those patients with a positive CT PET scan, 88% were treated with steroids. Of those patients with a negative CT PET scan, 42% were treated with steroids. It is noteworthy that 29% of patients were on steroids at the time of CT PET which may impact results. 60% of patients who were on steroids at the time of CT PET were on 60 mg of prednisolone daily. 31% of patients with negative scans were on steroids at the time of CT PET. 46% of patients with negative CT PET scans remained on steroid treatment. Steroid treatment was continued in patients with negative scans on basis of active aortic valve histology $\pm$ clinical criteria for diagnosis of GCA $\pm$ cerebral vasculitis on neuroimaging $\pm$ polymyalgia rheumatica evident on CTPET. CT PET changed management in 65% of patients with positive results supporting steroid treatment and negative results guiding withdrawal of steroids.

Conclusions: We are fortunate to have access to CT PET in Northern Ireland. CT PET scans changed management in 65% of our patients, despite 29% of patients being on steroid treatment prior to CT PET. We wish to increase awareness of the role CTPET in the diagnosis and management of LVV. We are liaising with radiology colleagues to refine and maximise appropriate referrals for CT PET scans for patients with suspected vasculitis.

REFERENCES:

- [1] BSR and BHPR guidelines for the management of giant cell arteritis. Bhasakar Dasgupta, et al. Rheumatology, Volume 49, Issue 8, 1 August 2010, Pages1594–1597, Published: 05 April 2010
- [2] EULAR recommendations for the use of imaging in large vessel vasculitis in clinical practice. Christian Dejaco, et al. Ann Rheum Dis published online January 22, 2018

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2018-eular.7390

AB1225 ANTIBODIES AGAINST HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEINS (RA33) MAY HAVE A DIAGNOSTIC AND PROGNOSTIC VALUE IN RHEUMATOID ARTHRITIS, PARTICULARLY WHEN OTHER SEROLOGICAL TESTS ARE NEGATIVE

V.A. Aleksandrov¹, L.N. Shilova², A.V. Alexandrov¹. ¹Federal State Budgetary Science Institution Research Institute for clinical and experimental rheumatology; ²The Department of Hospital Therapy, The Volgograd State Medical University, Volgograd, Russian Federation

Background: The high frequency of detection of antinuclear antibodies in rheumatoid arthritis (RA), although predominantly in low titres, allows us to consider the possibility of using varieties of these antibodies (primarily antibodies to RA-33) markers of RA especially in the early stages of the disease.

Objectives: To study the frequency of occurrence and determine the prognostic significance of antibodies to RA-33 in RA patients.

Methods: 57 RA patients were examined (mean age 50.5 $\pm$ 10.1 years).

The patients with the developed stage of the disease (47.4%) prevailed, the average activity (DAS28=3.2–5.1) of the pathological process (86%); with the second radiographic stage (42%) and functional class 2 (77%).

The antinuclear factor was determined in the indirect immunofluorescence reaction on the HEp-2 cell line (norm <1:80), IgM-rheumatoid factor by the latex agglutination method (up to 20 IU/ml), antibodies (norm up to 20 U/ml) and antibodies of IgG class to RA33 antigen (norm up to 25 U/ml) by ELISA test.

Results: ANF was detected in 18 patients with RA (32%), and in 94.4% of cases (17 patients) a diagnostic titer of 1:80 and a diffuse type of glow of the nucleus were noted. The ANF titres did not correlate either with the activity of the disease or with extra-articular manifestations of RA ($p>0.1$). Anti-RA33 was detected in 20 (35%) RA patients: 18 positive people (90%) had low positive anti-RA33 values (25 to 75 U/ml). In 35 (61.4%) of RA patients, anti-CCP was detected: 19 (33.3%) had low positive values (20 to 60 U/ml), 16 (28%) had highly positive values (more 60 U/ml). IgM-RF was detected in 26 (45.6%) patients in the values >48 IU/ml, as well as in 17 (30%) patients with RA in the values from 24 to 48 IU/ml. 14 people (54%) of patients with high IgM-RF levels had systemic manifestations of RA. It should be noted that anti-RA33 was detected in 9 patients with seropositive for anti-CCP and in 11 patients with seronegative both for anti-CCP and IgM-RF. Thus, when the results of standard serological tests are negative, an additional study of anti-RA33 is recommended to diagnose seronegative RA.

When assessing the prognostic significance of the available clinical and laboratory and instrumental data we have analysed the results of the study of anti-CCP, anti-RA33, as well as all data on the presence of X-ray erosions in patients with RA with magnetic resonance imaging or ultrasound examination of affected joints. The presence of erosion was noted in 23 (40.4%) RA patients. The frequency of detection of anti-CCP in RA patients was significantly higher in the presence of erosive lesions of joints (19 of 35 patients were positive by anti-CCP, compared with 4 of 22 patients, negative for anti-CCP, $\chi^2=5.89$, $p=0.015$). In the RA group of patients positive for a wide range of antibodies (IgM-RF, anti-CCP, anti-RA33), the signs of joint erosion were identified in 22 of 46 patients, and in isolated increase only anti-RA33 – in 1 patient out of 11 people in this group $\chi^2=4.04$, $p=0.044$.

Conclusions: In the presence of highly positive anti-CCP values, RA patients have a more unfavourable prognosis, while an isolated increase in anti-RA33 is associated with a "milder progression" of the disease and inversely proportional to erosive processes in the joints.

Disclosure of Interest: None declared
DOI: 10.1136/annrheumdis-2018-eular.6309

AB1226 THE DIAGNOSTIC VALUE OF SERUM KL-6 IN CONNECTIVE TISSUE DISEASE ASSOCIATED INTERSTITIAL LUNG DISEASE

X. Wu, C.N. Luo, Y.M. Shi, L.J. Wu. Department of Rheumatology and Immunology, People's Hospital of Xinjiang Uygur Autonomous Region, Urumqi, China

Background: The connective tissue diseases is a group of inflammatory, immune-mediated diseases. CTD often leads to autoimmunity and subsequent tissue injury. It is an important contribute to thoracic changes, particularly interstitial lung disease, is are the main causes of mortality and morbidity among patients with connective tissue diseases. Prognosis and response to therapy are the most pressing challenges for connective tissue disease-associated ILD (CTD-ILD).

At present, the basic methods for the diagnosis of various types of ILD includes high-resolution computed tomography, bronchoscopy, and surgical lung biopsy. In addition, continuous lung function tests are commonly used to monitor disease activity and predict the outcome of patients with ILDs. But these tests require specific inspection machines and Repeatability is not good. At present, many biomarkers have been developed to detect ILDs, and the most important biomarkers is KL-6 and lung surface active protein A (SP- A) and surfactant protein D (SP – D) witch secreted by alveolar epithelial type II cells. But Relevant studies have shown that the sensitivity, specificity and accuracy of KL-6 are higher than SP-A and SP-D.

Objectives: To evaluate the diagnosis of the serum Krebs von den Lungen-6 (KL-6) for the interstitial lung disease(ILD) associated with connective tissue diseases(CTD).

Methods: We retrospectively analysed the medical records of 50 patients with CTD associated ILD,46 CTD patients without ILD. Measurement of serum KL-6 levels and pulmonary function tests performed in parallel were reviewed.T test, X^2 test, non-parametric test, SPSS and correlation analysis were used for data analysis.

Results: The significantly higher levels of KL-6 were determined in the CTD-ILD group than in either the CTD without pulmonary involvement group($P<0.05$). Serum KL-6 correlated negatively with forced vital capacity(FVC%) (%predicted), forced expiratory volume in one second (FEV1)(%predicted) and diffusing capacity of the lung for carbon monoxide (DLCO) (% predicted). By the ROC curves of serum KL-6 levels in 96 patients. The optimal cutoff value of serum KL-6