Abstract AB1252 – Table 1. Characteristics of patients with RA and comorbidities

Pain medication	n	%
Non-opioids	2234	52
Opioids	990	23
NSAID	67	2
COMBINATIONS		
Non opioids +Opioids	430	10
Non opioids +NSAID	309	7
Opioids+NSAID	209	5
All	70	2

Conclusions: Rheumatoid arthritis is a pain associated condition; two thirds of patients are using of pain medications mainly women; the most prescribed medication was paracetamol or opioids, coinciding with other studies.² This descriptive study is useful for further studies to develop in Latin America.

REFERENCES:

- [1] Parker JC, Frank RG, Beck NC, Smarr KL, Buescher KL, Phillips LR, et al. Pain management in rheumatoid arthritis patients. *Arthritis & Rheumatism*. 1988;31(5):593–601.
- [2] Samison LH, Randriatsarafa FM, Ralandison S. Joint pain epidemiology and analgesic usage in Madagascar. *The Pan African medical journal*. 2017;26:77.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7201

AB1253

ANALGESIC DRUGS AND RISK OF ISCHAEMIC STROKE IN PATIENTS WITH OSTEOARTRITIS: A REAL WORLD DATA CASE-CONTROL STUDY

A. Gomez-Lumbreras¹, R. Vives^{2,3}, M. Giner-Soriano^{1,3,4}, H. Pera¹, M. Fradera², J. R. Marsal⁵, R. Morros^{1,3,4}. ¹Medicines Research Unit, Institut Universitari d'Investigació en Atenció Primària (IDIAPJGol), Barcelona; ²Institut d'Investigació i Innovació Parc Taulí, Corporació Sanitària Parc Taulí, Sabadell; ³Universitat Autònoma de Barcelona; ⁴Institut Català de la Salut, Barcelona; ⁵Unitat de Suport a la Recerca, Institut Universitari d'Investigació en Atenció Primària (IDIAPJGol), Lleida, Spain

Background: Pharmacological treatment of osteoarthritis (OA) usually include analgesics, non-steroidal anti-inflammatory drugs (NSAID) and symptomatic slow-acting drugs in OA (SYSADOA). The association between these groups of drugs and the risk of ischaemic stroke has not been properly addressed.

Objectives: To analyse the risk of stroke in patients using analgesics, NSAID and SYSADOA drugs.

Methods: We used a population-based patient hospital registry to identify all patients with a first-ever stroke discharge diagnosis between 2009–2015. Cases were matched to controls obtained from the Information System for Research in Primary Care (SIDIAP) database. Information on drug exposure was obtained from invoice data from pharmacies. Crude and adjusted odds ratios (OR, OR_{adj}) and their 95% confidence interval (95% CI) were calculated using multivariate models of conditional logistic regression for the next pharmacological groups and individual agents of each group: acetic acid derivatives, oxicams, propionic acid and derivatives, coxibs, SYSADOA and analgesics (opioids, metamizole and paracetamol). A cardiovascular risk score was calculated for each subject based on comorbidities.

Results: 12 616 cases were matched to 1 25 264 controls by gender, age and geographic area. Among cases, 43% were women. The mean age was 72.6 (IQR 65–82) years and more cases were classified as high cardiovascular risk patients (n=2,511, 19.9%) than controls (n=12,467, 10.0%). Mortality in the following year after the index date was higher for cases (n=2,633, 20.9%) than for controls (n=8,168, 6.5%). A higher percentage of cases had a previous diagnosis of ischaemic heart disease, 13.8% (n=1,745) vs 7.7% (n=9,656) of the controls. OA diagnosis was present in 2823 (22.4%) cases and in 29 098 (23.2%) controls. Paracetamol was the most used drug (n=106,515, 77.3%) followed by ibuprofen (n=84,790, 61.5%).

Current users of acetic acid derivatives showed an increased risk of stroke [OR_{adj} (95% CI) 1.10 (1.01–1.19)], so did diclofenac [OR_{adj}1.14 (1.04–1.25)]. Among the propionic acid derivatives, current users showed an increased risk [OR_{adj}1.24 (1.27–1.32)], dextetopfen showed the highest risk [OR_{adj}1.42 (1.25–1.60)] and naproxen the lowest [OR_{adj}1.18 (1.02–1.37)]. The SYSADOA group did not show any increased risk for any type of exposure, with a decreased risk of 17% [OR_{adj}0.83 (0.77–0.91)] for lifetime glucosamine exposure and a 22% [OR_{adj}0.84 (0.78–0.90)] for chondroitin sulfate. Analgesics were the most consumed drugs,

and cases were more exposed to all subgroups of analgesics than controls. While the non-adjusted OR showed an increased risk of stroke for lifetime exposure for all agents in this group, this association was not observed with the adjusted OR_{adj} for fentanyl, neither buprenorphine, but it showed a risk for current users of metamizole [OR_{adj} of 1.67 (1.56–1.80)], tramadol [OR_{adj}1.15 (1.06–1.24)] and paracetamol [OR_{adj}1.43 (1.37–1.51)].

Conclusions: Current exposure to NSAIDs, tramadol, metamizole and paracetamol is a risk factor for ischaemic stroke. Exposure to chondroitin sulphate and glucosamine are associated with a lower risk of ischaemic stroke.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1428

AB1254

IMPROVING RHEUMATOLOGIC CARE AND EDUCATION IN THE REPUBLIC OF MACEDONIA: A MODEL FOR PROMOTING RHEUMATOLOGIC EDUCATION IN A DEVELOPING COUNTRY

V. Ogneovski¹, M. Arsovska-Nalbanti². ¹Division Rheumatology, University of Michigan, Ann Arbor, USA; ²Division Rheumatology, Clinical Center Dr. Trifun Panovski, Bitola, Macedonia, The Former Yugoslav Republic of Ireland

Background: In 2011 ILAR supported a pilot project in training a rheumatologist in an underserved area in Macedonia. The pilot took place in the clinical centre of Bitola. Previously, rheumatologic care was provided by visiting rheumatologists from the rheumatology clinic in the capital city.¹

Objectives: The primary goal of this project was to pilot a model in rheumatologic training and improve rheumatologic care in an underserved area in Macedonia.

Methods: An internist from the clinical centre in Bitola was enrolled in a two-year training by the Rheumatology clinic at Ss Cyril and Methodius University in Macedonia, and the rheumatology division at the University of Michigan. Pre intervention metrics of quality of care as measured by access, standard therapy, and DAS 28 scores were compared with post intervention metrics.

Results: The primary goal of the project enabled training of a rheumatologist. A local internist completed her training in rheumatology (2014), and established a rheumatology clinic within the clinical centre in the city of Bitola in 2014, providing daily access to patients with rheumatic conditions in a region with a population of 3 00 000. Patient access changed from 120 visits/month to 800 visits/month. The service provided consultations to hospitalised patients.

The secondary endpoints: quality of care as measured by standard therapy for rheumatoid arthritis and DAS-28 scores are as shown on table 1.

Abstract AB1254 – Table 1

Rheumatoid Arthritis patients	Pre-intervention 2011 (n=115)	Post-intervention 2017 (n=86)	Statistical Significance
Patients taking methotrexate	44%	53%	
Methotrexate dose	11.5 mg (10–15 mg)	13.9 mg (10–25 mg)	p<0.00001
Combination therapy*	39%	38.5%	NS
Dual therapy*	31%	35%	NS
Triple therapy*	3.5%	5%	NS
DAS-28 average	4.8	4.41	NS

MTX=methotrexate; *methotrexate, sulfasalazine, leflunomide, antimalarial (chloroquine, hydroxychloroquine)

Post intervention, more patients were taking methotrexate and at higher doses. Despite this trend, its average dose was less than 50% of its maximal dose (25 mg/week) commonly used in standard practice. The frequency of combination therapy remained unchanged. Likewise, no significant change in the DAS-28 scored was observed.

Conclusions: Prior to launching the pilot, the rheumatologic care in this region was provided by visiting rheumatologists from the university clinic in the capital city. Initial assessment pointed to several obstacles: poor access and substandard therapy that likely contributed to the prevalence of high disease activities. Our pilot succeeded in training a rheumatologist, thus reaching our primary goal of improving local access to rheumatic care. The secondary goals of improving the quality of care as measured by the DAS-28, standard of therapy for rheumatoid arthritis- reflected by appropriate use of DMARDs (use of methotrexate, combination therapy) showed a modest improvement While higher doses of methotrexate were used, combination therapy and better control of rheumatoid arthritis (DAS-28) remained unchanged, thus posing a challenge for ongoing need and future intervention goals.

REFERENCE:

- [1] Ogneovski, VM, Arsovska- Nalbanti, M, Chichikj D, Calovski, J. Improving care and education in the Republic of Macedonia: model for rheumatologic