DOI: 10.1136/annrheumdis-2018-eular.4202

AB1325

THE FREQUENCY OF ANTI-DFS70 AUTOANTIBODIES IN JAPANESE WOMEN WITH PERI- AND POST-MENOPAUSAL ARTHRALGIA

K. Miyachi^{1,2}, A. Ihar³, B. Sasse⁴, M.Y. Choi⁵, M.J. Fritzler⁵. ¹Rheumatology and Women's Health, Keigu Clinic; ²The Technological Development, Health Sciences Research Institute; ³Rheumatology, Keigu Clinic, Yokohama, Japan; ⁴Rheumatology, Monash Health, Victoria, Australia; ⁵Cumming School of Medicine, University of Calgary, Calgary, Canada

Background: The antinuclear antibody (ANA) test is often used as a screening test to aid in the diagnosis of systemic lupus erythematosus (SLE) and other systemic autoimmune rheumatic diseases (SARD). However, recently it was reported that a certain autoantibody referred to as anti-dense fine speckled 70 (DFS70) are much more common (up to 25%) of non-SARD and healthy individuals but rare (<5%) in SLE and certain SARD. The frequency of anti-DFS70 is even lower (i.e. <1%) if monospecific anti-DFS70 antibodies are found. Some reports suggested that anti-DFS70 are more common in younger females, suggesting that hormonal factors may be responsible for the cognate B cell response.

Objectives: The primary objective of our study was to determine the frequency of anti-DFS70 in menopausal women who were referred for evaluation of undifferentiated arthritis (UA).

Methods: 282 women including 105 with UA and an age range of 27–91 years (mean=60.3) were enrolled. Menopausal women were divided into pre-, peri-, and post-menopausal stages according to menstrual regularity. (E2), Follicle stimulating hormone (FSH), rheumatoid factor (RF), ANA (by indirect immunofluorescence (IIF) (MBL) anti-cyclic citrullinated peptide antibody (ACPA: MBL ELISA) and C-reactive protein (CRP) were included in the serology workup. Postmenopausal arthralgia was designated PoMA and perimenopausal arthralgia PeMA.

Results: In PoMA women who received HRT for two months estradiol levels increased, FSH levels decreased and the joint pain visual analogue scale was reduced by 70%, as compared to baseline. Similarly in PeMA women, administration of 600 mg tocopherol N daily had the same efficacy as that observed in PoMA. The overall frequency of anti-DFS70 of 26.7% (28/105) in PoMA and PeMA women was significantly higher than that in UA females who were diagnosed with rheumatoid arthritis (RA) (7/65:10.8%) or primary Sjogren's syndrome (SjS) (3/31: 9.7%) (p<0.05). In addition, anti-DFS70 Ab was observed primarily in low titer (1:40 – 1:160) ANA positive sera. On the other hand, higher titer ANAs (titer >1:320) were observed in the females that were diagnosed as systemic lupus erythematosus (SLE) (30%) or primary SjS (48.4%), the majority of whom had lower titers of anti-DFS70, although several sera contained both high titer ANA and anti-DFS70 (see figure).

Abstract AB1325 – Figure 1. Relationship of ANA titers and anti-DFS70 levels in peri- and post-menopausal women. The majority of anti-DFS70 positive sera had only low to modest ANA titers. ANA titer in vertical axis and anti-DFS70 titers by ELISA on the horizontal axis.

Conclusions: Anti-DFS70 was found in higher frequency in PeMA and PoMA women than in women who developed a defined systemic autoimmune rheumatic disease such as SLE, SjS or RA. This is the first study to suggest that the presence of this autoantibody may reflect oestrogen fluctuations or deficiency. A negative association between ANA titer and anti-DFS70 levels by ELISA remains to be confirmed in larger studies.

REFERENCES:

- M. Mahler, P. L. Meroni, L. E. Andrade, M. Khamashta, N. Bizzaro, C. A. Casiano, and M. J. Fritzler. Towards a better understanding of the clinical association of anti-DFS70 autoantibodies. *Autoimmun. Rev.* 15:198–201, 2016.
- K. Conrad, N. Rober, L. E. Andrade, and M. Mahler. The Clinical Relevance of Anti-DFS70 Autoantibodies. *Clin. Rev. Allergy Immunol.* 52:202–216, 2016.

Disclosure of Interest: K. Miyachi Shareholder of: none, Grant/research support from: none, Consultant for: none, A. Ihar: None declared, B. Sasse: None declared, M. Y. Choi: None declared, M. J. Fritzler: None declared
DOI: 10.1136/annrheumdis-2018-eular.2013

AB1326

NICOTINAMIDE PHOSPHORIBOSYLTRANSFERASE MAY BE NEW FACTORS CONTRIBUTING TO OSTEOARTHRITIS

L. Sivordova¹, J. Polyakova¹, Y. Akhverdyan¹, V. Kravtsov¹, N. Fofanova², B. Zavodovsky¹. ¹Federal State Budgetary Institution "Research Institute of Clinical and Experimental Rheumatology named after A.B. Zborovskiy"; ²Federal State Budgetary Volgograd Medical University, Volgograd, Russian Federation

Background: Obesity is a condition that prolongs chronic inflammation and promotes synthesis and secretion of pro-inflammatory factors by adipose tissue including adipokines¹ and can be one of the significant factors in the progression of osteoarthritis (OA).² Nowadays many studies have shown that level nicotinamide phosphoribosyltransferase (Namt) increases in obesity and can also affect the inflammation in local tissues and together with other adipokines influences the development of OA.³

Objectives: To study the effect of weight loss over 5 kg on the clinical manifestations of OA, indicators of water, lipid metabolism and Nampt serum levels in patients with OA.

Methods: We observed 110 people: 80 patients with OA and 30 healthy individuals in the control group with body mass index (BMI) of 25 to 35 kg/m², aged 18 to 79 years participated in the study.

Namt level in serum was determined by indirect solid phase ELISA using a commercial test systems (RaiBiotech, cat. № EIA – VIS – 1).

Results: As overweight patients were recruited in the study, hypocaloric diet low in animal fats and physiotherapy has been recommended to all participants. The first group consisted of patients who were able to reduce body weight by 5 kg and more (18 pers.), the second group – patients with weight reduction of less than 5 kg and patients without any weight loss (62 pers.). Nampt concentration were evaluated in these patients I groups 4,33±0,39 before treatment to 2,40±0,23 ng/ml (t=5,85, p<0,001). In analysing the parameters before and after treatment, it should be noted that we observed significant decrease in the severity of the clinical manifestations of OA, visfatin level, CRP, and glucose levels and lipid profile in the 1st group of patients. These data proves that obesity may be an important risk factor for OA progression. As a result, weight loss results in decreasing metabolic disorder severity. In the second group of patients we have seen a decrease in all the parameters, but a significant difference has been observed only in the level of CRP, level of pain at rest and during walking according to VAS scale and total index on the WOMAC.

Conclusions: As a result of our study patients with OA with weight loss of more than 5 kg had more obvious pain relief than patients with the original weight. At the same time a significant improvement has been seen in carbohydrate and lipid metabolism. These findings suggest that there is a possible role of Nampt in the pathogenesis of OA.

REFERENCES:

- Akhverdyan Y, Zavodovsky B, Seewordova L, et al. Adipokines as new laboratory markers in osteoarthritis. *Ann Rheum Dis* 2013;72 (Suppl3):702.
- Polyakova Y, Seewordova L, Akhverdyan Y. et al. Some pathophysiological aspects of pain syndrome in osteoarthritis. *Osteoporosis International* 2017;28(Suppl 1):154.
- Sivordova LE, Simakova ES, Polyakova JV, et al. Overweight and high level of nicotinamide phosphoribosyltransferase as factors contributing to osteoarthritis progression and metabolic syndrome development. *Bulgarian medicine* 2016;Tom VI, 1:11–13.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3064

AB1327

CARDIAC MANIFESTATIONS IN SPONDYLOARTHRITIS: PREVALENCE AND PREDICTIVE FACTORS

J. Eddarami, L. Ichchou. Department of Rheumatology, Mohamed VI University Hospital, Mohammed I University, Oujda, Morocco

Background: Cardiac involvement is a well-recognised complication of spondyloarthritis (SpA). The spectrum of this condition is large and includes mitral valve disease, conduction disorders and pericarditis, but the aortic disease remains the most characteristic lesion.