Results: A total of 166 SLE patients participated to the study (93.4% female, 97% Caucasian); mean age ($\pm$ standard deviation) at enrollment was 45.4 years $\pm$ 13, mean disease duration resulted 14.3 $\pm$ 10 years. Overall, disease activity at enrollment as expressed by the SLEDAI score resulted quite low (median SLEDAI 2, IQR 0–4) and 27.3% of patients presented an active disease (SLEDAI >4). At least one organ damage according with the SLICC/DI was present in 111 patients (56.35%) with a median score of 2 (IQR1–4). In the pilot test, 80% of patients answered to all the questions; 93% of the patient declared no difficulty in completing the questionnaire and the median time required resulted 5 min.^{3–5} BILDit showed very high reliability (test-retest α >0.8). The BILDit scores in our sample showed a strong positive correlation with SLICC/DI (ρ =0.69; p <0.001); by comparing the single BILDit and SLICC/DI items, we found a significant concordance for all but retinopathy and cerebrovascular disease that resulted underestimate by patients. BILDit scores showed a positive linear relation with age (p <0.01) and disease duration (p <0.01); in a multivariate analysis, a significant inverse correlation was found between BILDit scores and FACIT scores (p <0.01) and with both physical and mental components of the SF-36 (p <0.01).

Conclusions: BILDit demonstrated to be acceptable, comprehensible, feasible and reliable in our routine clinical setting; it also showed good correlation with physician's driven instruments and quality of life measures. The BILDit can be considered a useful screening tool for the assessment of organ damage as perceived from the patient's point of view before the standard visit.

REFERENCES:

- [1] Yazdany, J. et al. The Brief Index of Lupus Damage (BILD): A patient-reported measure of damage in SLE. *Arthritis Care Res.* 2011; 63, 1170–1177.
- [2] Guillemin F. et al. Cross-cultural adaptation of health-related quality of life measures: Literature review and proposed guidelines. *Journal of Clinical Epidemiology* 1993; 46: 1417–1432

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6519

AB1360

RANGE AND CONSISTENCY OF OUTCOME MEASURES REPORTED IN RANDOMISED TRIALS IN DERMATOMYOSITIS: A SYSTEMATIC REVIEW

A.H. Kelly^{1,2,3}, G. Hassett³, D. Singh-Grewal², A. Tong¹. ¹Sydney School of Public Health, University of Sydney; ²Paediatric Rheumatology, Sydney Children's Hospital Network; ³Rheumatology, Liverpool Hospital, Sydney, Australia

Background: Dermatomyositis – an idiopathic inflammatory myopathy- can be resistant and unresponsive to initial treatments and require chronic therapy. Symptoms including muscle weakness and skin rash can lead to severe complications and impair quality of life. There are a paucity of randomised trials in dermatomyositis, which may not consistently report outcomes that are important to patients and their clinicians.

Objectives: To assess the scope and consistency of outcomes reported in randomised trials in dermatomyositis.

Methods: MEDLINE, Embase, CINAHL, PsycINFO were searched to February 2017 for randomised trials in children and adults with dermatomyositis. The frequency and characteristics of the outcome domains and measures reported were then analysed. Similar outcomes were grouped into outcome domains and classified as surrogate, clinical or patient-reported outcomes. The frequency and characteristics of the outcome domains and measures reported were then analysed.

Results: We included 18 trials (n=844 participants), reporting 680 outcome measures. These were grouped into 27 different outcome domains; including 12 (44%) clinical outcomes, 9 (33%) surrogate/biochemical outcomes, and 6 (22%) patient-reported outcomes. The 5 most frequently reported outcome domains were: physical function (13 trials, 72%), muscle strength (12, 67%), muscle inflammation (12, 67%), biomarkers (11, 61%), and composite i.e. myositis disease activity (10, 56%).

Conclusions: The majority of outcomes reported in trials in dermatomyositis are clinical outcomes and few are patient-reported. Establishing a core set of patient-important outcomes may improve the consistency and relevance of outcomes reported in trials in dermatomyositis to inform decision-making.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3153

AB1361

SEASONAL SENSITIVITIES OF DISEASE ACTIVITY INDICES IN RHEUMATOID ARTHRITIS

A. Şahin Eşkara¹, N. Taştekin¹, M. Birtane¹, N. Süt². ¹Trakya University Faculty of Medicine, Department of Physical Therapy and Rehabilitation, Edirne; ²Trakya University Faculty of Medicine, Department of Biostatistics and Medical Informatics, Edirne, Edirne, Turkey

Background: Rheumatoid arthritis (RA) is a chronic and progressive autoimmune disease which is primarily characterised by the inflammation of synovial joints. In rheumatoid arthritis, evaluation of the disease activity is very important in monitoring the disease progression and effectiveness of the treatment

Objectives: The aim of this study was to evaluate the seasonal sensitivity of disease activity indices (DAS28-ESR, DAS28-CRP, SDAI and CDAI) in rheumatoid arthritis. In addition, we investigated the presence of seasonal variation in disease activity in rheumatoid arthritis and its relation to vitamin D levels.

Methods: This is a prospective study. This study included female RA patients, who were diagnosed/verified according to 2010 ACR/EULAR criteria, with a diagnostic history of 5 years or older. The seasons were identified as spring (March, April, May), summer (June, July, August), autumn (September, October, November) and winter (December, January, February), and the study was completed with 85 patients. We assessed the disease activation with DAS28-ESR, DAS28-CRP, SDAI and CDAI scales, functional status with HAQ questionnaire, quality of life with the 36-Item Short Form Health Survey (SF-36), and presence of depression with beck depression inventory (BDI). Blood samples from patients were kept at appropriate storage conditions at –80 °C, and serum levels of 25-OH vitamin D were analysed by the chemoluminescence assay method at the end of each season.

Results: The assessment of mean of DAS28-ESR and DAS28-CRP scores found that the disease activity showed statistically significant seasonal differences (p <0.05). Although the assessment of mean of SDAI and CDAI scores showed that the disease activity was higher in the autumn season, this difference was not statistically significant (p >0.05). There was no statistically significant correlation between the mean vitamin D levels and mean of DAS28-ESR, DAS28-CRP, SDAI, CDAI scores in any season (p >0.05).

Conclusions: These results suggest that there may be seasonal variation in disease activity in RA and that DAS28-ESR and DAS28-CRP may be more sensitive than CDAI and SDAI in establishing this variation. There was no correlation between seasonal changes in disease activity and changes in serum 25-OH Vitamin D levels

REFERENCE:

- [1] Anderson J, Caplan L, Yazdany J, Robbins ML, Neogi T, Michaud K, et al. Rheumatoid arthritis disease activity measures: American college of rheumatology recommendations for use in clinical practice. *Arthritis Care Res* 2012;64(5):640–7.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2965

AB1362

SPECIFIC FEATURES OF THE CLINICAL COURSE AND OUTCOMES OF ERYTHEMA NODOSUM, ASSOCIATED WITH INFECTION

B. Belov, N. Savushkina, O. Egorova, V. A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: Erythema nodosum (EN) is the most common type of septal panniculitis (Pn) without vasculitis. The prevailing identifiable cause of EN in paediatric population is bacterial infection (*Streptococcus*), while in adults sarcoidosis and streptococcal infection are considered to be the most common etiological factors.

Objectives: The aim of the current study is to evaluate specific features of the clinical course and outcomes of EN, associated with infection, in rheumatology practice.

Methods: The study included 121pts (112 females—92,6%, 9 males—7,4%, mean age 38,9 $\pm$ 12,6 y) who were on the record at VA Nasonova Research Institute of Rheumatology during 7 years with referral diagnosis «erythema nodosum». The disease duration varied from 1 week to 20 years. Patient's assessment included general clinical examination, immunological and serological tests for *Chlamidia* (*Chl. Pneumoniae*, *Chl. trachomatis*), *Mycoplasma* (*M. pneumoniae*, *M. hominis*), *Ureaplasma*, Herpes viruses type I and II, Cytomegalovirus, Epstein-Barr virus, *Yersinia*, HBV and HCV, HIV, anti-streptolysin O (ASO) using ELISA and immunoturbidimetry. The outcomes were assessed in 1–6 years.

Results: EN association with infection was identified in 32 pts (30 females, 2 males, mean age 39,6 $\pm$ 13 y). Chronic conditions prior to EN onset were found in 12 pts (37%): tonsillitis—in 8 pts (25%), endometriosis—in 1 (3%), bronchitis—in 1 (3%), colpitis—in 1 (3%), toxoplasmosis—in 1 (3%). Elevated ASO titers were documented in more than half cases (18 (56,25%) pts). "Blooming bruise" symptom was characteristic for this population (p <0,0001). Clear clinical and laboratory