Results: A total of 166 SLE patients participated to the study (93.4% female, 97% Caucasian); mean age ($\pm$ standard deviation) at enrollment was 45.4 years $\pm$ 13, mean disease duration resulted 14.3 $\pm$ 10 years. Overall, disease activity at enrollment as expressed by the SLEDAI score resulted quite low (median SLEDAI 2, IQR 0–4) and 27.3% of patients presented an active disease (SLEDAI >4). At least one organ damage according with the SLICC/DI was present in 111 patients (56.35%) with a median score of 2 (IQR1–4). In the pilot test, 80% of patients answered to all the questions; 93% of the patient declared no difficulty in completing the questionnaire and the median time required resulted 5 min.^{3–5} BILDit showed very high reliability (test-retest α >0.8). The BILDit scores in our sample showed a strong positive correlation with SLICC/DI (ρ =0.69; p <0.001); by comparing the single BILDit and SLICC/DI items, we found a significant concordance for all but retinopathy and cerebrovascular disease that resulted underestimate by patients. BILDit scores showed a positive linear relation with age (p <0.01) and disease duration (p <0.01); in a multivariate analysis, a significant inverse correlation was found between BILDit scores and FACIT scores (p <0.01) and with both physical and mental components of the SF-36 (p <0.01).

Conclusions: BILDit demonstrated to be acceptable, comprehensible, feasible and reliable in our routine clinical setting; it also showed good correlation with physician's driven instruments and quality of life measures. The BILDit can be considered a useful screening tool for the assessment of organ damage as perceived from the patient's point of view before the standard visit.

REFERENCES:

- [1] Yazdany, J. et al. The Brief Index of Lupus Damage (BILD): A patient-reported measure of damage in SLE. *Arthritis Care Res.* 2011; 63, 1170–1177.
- [2] Guillemin F. et al. Cross-cultural adaptation of health-related quality of life measures: Literature review and proposed guidelines. *Journal of Clinical Epidemiology* 1993; 46: 1417–1432

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6519

AB1360

RANGE AND CONSISTENCY OF OUTCOME MEASURES REPORTED IN RANDOMISED TRIALS IN DERMATOMYOSITIS: A SYSTEMATIC REVIEW

A.H. Kelly^{1,2,3}, G. Hassett³, D. Singh-Grewal², A. Tong¹. ¹Sydney School of Public Health, University of Sydney; ²Paediatric Rheumatology, Sydney Children's Hospital Network; ³Rheumatology, Liverpool Hospital, Sydney, Australia

Background: Dermatomyositis – an idiopathic inflammatory myopathy- can be resistant and unresponsive to initial treatments and require chronic therapy. Symptoms including muscle weakness and skin rash can lead to severe complications and impair quality of life. There are a paucity of randomised trials in dermatomyositis, which may not consistently report outcomes that are important to patients and their clinicians.

Objectives: To assess the scope and consistency of outcomes reported in randomised trials in dermatomyositis.

Methods: MEDLINE, Embase, CINAHL, PsycINFO were searched to February 2017 for randomised trials in children and adults with dermatomyositis. The frequency and characteristics of the outcome domains and measures reported were then analysed. Similar outcomes were grouped into outcome domains and classified as surrogate, clinical or patient-reported outcomes. The frequency and characteristics of the outcome domains and measures reported were then analysed.

Results: We included 18 trials (n=844 participants), reporting 680 outcome measures. These were grouped into 27 different outcome domains; including 12 (44%) clinical outcomes, 9 (33%) surrogate/biochemical outcomes, and 6 (22%) patient-reported outcomes. The 5 most frequently reported outcome domains were: physical function (13 trials, 72%), muscle strength (12, 67%), muscle inflammation (12, 67%), biomarkers (11, 61%), and composite i.e. myositis disease activity (10, 56%).

Conclusions: The majority of outcomes reported in trials in dermatomyositis are clinical outcomes and few are patient-reported. Establishing a core set of patient-important outcomes may improve the consistency and relevance of outcomes reported in trials in dermatomyositis to inform decision-making.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3153

AB1361

SEASONAL SENSITIVITIES OF DISEASE ACTIVITY INDICES IN RHEUMATOID ARTHRITIS

A. Şahin Eşkara¹, N. Taştekin¹, M. Birtane¹, N. Süt². ¹Trakya University Faculty of Medicine, Department of Physical Therapy and Rehabilitation, Edirne; ²Trakya University Faculty of Medicine, Department of Biostatistics and Medical Informatics, Edirne, Edirne, Turkey

Background: Rheumatoid arthritis (RA) is a chronic and progressive autoimmune disease which is primarily characterised by the inflammation of synovial joints. In rheumatoid arthritis, evaluation of the disease activity is very important in monitoring the disease progression and effectiveness of the treatment

Objectives: The aim of this study was to evaluate the seasonal sensitivity of disease activity indices (DAS28-ESR, DAS28-CRP, SDAI and CDAI) in rheumatoid arthritis. In addition, we investigated the presence of seasonal variation in disease activity in rheumatoid arthritis and its relation to vitamin D levels.

Methods: This is a prospective study. This study included female RA patients, who were diagnosed/verified according to 2010 ACR/EULAR criteria, with a diagnostic history of 5 years or older. The seasons were identified as spring (March, April, May), summer (June, July, August), autumn (September, October, November) and winter (December, January, February), and the study was completed with 85 patients. We assessed the disease activation with DAS28-ESR, DAS28-CRP, SDAI and CDAI scales, functional status with HAQ questionnaire, quality of life with the 36-Item Short Form Health Survey (SF-36), and presence of depression with beck depression inventory (BDI). Blood samples from patients were kept at appropriate storage conditions at –80 °C, and serum levels of 25-OH vitamin D were analysed by the chemoluminescence assay method at the end of each season.

Results: The assessment of mean of DAS28-ESR and DAS28-CRP scores found that the disease activity showed statistically significant seasonal differences (p <0.05). Although the assessment of mean of SDAI and CDAI scores showed that the disease activity was higher in the autumn season, this difference was not statistically significant (p >0.05). There was no statistically significant correlation between the mean vitamin D levels and mean of DAS28-ESR, DAS28-CRP, SDAI, CDAI scores in any season (p >0.05).

Conclusions: These results suggest that there may be seasonal variation in disease activity in RA and that DAS28-ESR and DAS28-CRP may be more sensitive than CDAI and SDAI in establishing this variation. There was no correlation between seasonal changes in disease activity and changes in serum 25-OH Vitamin D levels

REFERENCE:

- [1] Anderson J, Caplan L, Yazdany J, Robbins ML, Neogi T, Michaud K, et al. Rheumatoid arthritis disease activity measures: American college of rheumatology recommendations for use in clinical practice. *Arthritis Care Res* 2012;64(5):640–7.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2965

AB1362

SPECIFIC FEATURES OF THE CLINICAL COURSE AND OUTCOMES OF ERYTHEMA NODOSUM, ASSOCIATED WITH INFECTION

B. Belov, N. Savushkina, O. Egorova. V. A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: Erythema nodosum (EN) is the most common type of septal panniculitis (Pn) without vasculitis. The prevailing identifiable cause of EN in paediatric population is bacterial infection (*Streptococcus*), while in adults sarcoidosis and streptococcal infection are considered to be the most common etiological factors.

Objectives: The aim of the current study is to evaluate specific features of the clinical course and outcomes of EN, associated with infection, in rheumatology practice.

Methods: The study included 121pts (112 females—92,6%, 9 males—7,4%, mean age 38,9 $\pm$ 12,6 y) who were on the record at VA Nasonova Research Institute of Rheumatology during 7 years with referral diagnosis «erythema nodosum». The disease duration varied from 1 week to 20 years. Patient's assessment included general clinical examination, immunological and serological tests for *Chlamidia* (*Chl. Pneumoniae*, *Chl. trachomatis*), *Mycoplasma* (*M. pneumoniae*, *M. hominis*), *Ureaplasma*, Herpes viruses type I and II, Cytomegalovirus, Epstein-Barr virus, *Yersinia*, HBV and HCV, HIV, anti-streptolysin O (ASO) using ELISA and immunoturbidimetry. The outcomes were assessed in 1–6 years.

Results: EN association with infection was identified in 32 pts (30 females, 2 males, mean age 39,6 $\pm$ 13 y). Chronic conditions prior to EN onset were found in 12 pts (37%): tonsillitis—in 8 pts (25%), endometriosis—in 1 (3%), bronchitis—in 1 (3%), colpitis—in 1 (3%), toxoplasmosis—in 1 (3%). Elevated ASO titers were documented in more than half cases (18 (56,25%) pts). "Blooming bruise" symptom was characteristic for this population (p <0,0001). Clear clinical and laboratory

signs of active inflammation were not found. Based on etiological agent all pts were divided into 3 groups: Group I—with bacterial infection (5 pts (15%)), Group II—with viral infection (11 pts (35%)), Group III—with mixed bacterial and viral infection (16 pts (50%)). Rapid nodule regression was strongly associated with elevated ASO titers at baseline ($p=0.02$) and presence of bacterial agents ($p=0.0007$). Recurrences were documented in 13 pts (40.6%), among them 7 (54.1%) cases were triggered by ARVI/common frigoris, 2 (15.3%) cases—by stress, 2 (15.3%) cases—by non-compliance or treatment failure, 2 (15.3%)—by exacerbation of chronic tonsillitis. There was no statistically significant association between intake of individual medications and full reversal of the disease. There was 1 (8% from total number) recurrence episode in Group I, 7 (54%) episodes—in Group II, and 5 (38%)—in Group III. Recurrent disease inversely correlated with affected surface area (affected leg surfaces) ($p=0.03$). Pts who achieved nodular regression had elevated ASO at EN onset ($p=0.00008$), in contrast to pts with recurrent disease.

Conclusions: Streptococcus spp (56.3%) seem to be the leading cause of EN associated with infection. Lab verification of streptococcal infection with subsequent adequate antibacterial therapy facilitates the favourable clinical course of EN.

Acknowledgements: The study had no sponsorship.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2278

AB1363 ARE YOU ABLE TO CUT YOUR MEAT? – EXPLORING THE CHALLENGES DURING THE CULTURAL ADAPTATION OF THE HEALTH ASSESSMENT QUESTIONNAIRE DISABILITY INDEX (HAQ-DI) INTO 130 LANGUAGES

C. Acquadro¹, C. Anfray², P. Bindi², V. Uzun³. ¹Thought Leadership, Mapi, an ICON plc Company; ²Author Collaboration Unit, Mapi Research Trust; ³Language Services, Mapi, an ICON plc Company, Lyon, France

Background: The disability assessment component of the Health Assessment Questionnaire, the HAQ-DI,¹ developed in US English, assesses a patient's level of functional ability and includes questions of fine movements of the upper extremity, locomotor activities of the lower extremity, and activities that involve both upper and lower extremities. There are 20 questions in eight categories of functioning which represent a comprehensive set of functional activities – dressing, rising, eating, walking, hygiene, reach, grip, and usual activities. The stem of each item asks over the past week "Are you able to ..." perform a particular task. The patient's responses are made on a scale from zero (no disability) to three (completely disabled).

Objectives: To explore the challenges faced during the cultural adaptation of the HAQ-DI, focusing on one simple task, i.e., "Are you able to cut your meat?" in the eating category.

Methods: The archives of Mapi Language Services were searched and 130 translations were retrieved, representing 13 language families. The translation methods followed either the standard linguistic validation process [i.e., conceptual analysis, dual translation process (forward, backward translation into English), test with patients and clinician review] or the adjusted process in case of countries using national variants of the same language (i.e., Australian English or English used in India).²

Results: In most of the target languages, cutting a whole piece of meat presented in one plate was not a cultural issue. However, in vegetarian-driven cultures or countries where cutting meat was performed while cooking, the item had to be adapted and changed to comply both with the local culture and the concept explored by the item (i.e., fine movement of the upper extremity). For instance, an equivalent of "to cut meat while cooking" was chosen in Korean; "to cut meat (when eating or preparing food)," in Bahasa (Indonesia); "to cut raw vegetables," in Gujarati (India); or "make bite size pieces of your food (e.g., chapatis)" in English for India.

Conclusions: The close collaboration between the developer and the translation teams was essential in finding appropriate conceptual equivalents of the simple task of cutting meat in 13 different language families.

REFERENCES:

- [1] Bruce B, Fries JF. The Stanford Health Assessment Questionnaire: dimensions and practical applications. *Health Qual Life Outcomes*. 2003 Jun 9;1:20.
- [2] Mear I, Giroudet C. Chapter 1: Linguistic Validation Procedures. In: Acquadro C, Conway K, Giroudet C, Mear I, editors. *Linguistic Validation Manual for Health Outcome Assessments*. Lyon, France: Mapi Institute; 2012. p. 15–117.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4591

AB1364 ADAPTATION AND VALIDATION OF THE RHEUMATOID ARTHRITIS QUALITY OF LIFE (RAQOL) SCALE FOR PORTUGAL

F. Fernandes¹, C. Duarte^{1,2}, N. Madeira³, P. Pinto⁴, C. Furtado⁵, C. Ribeiro⁶, M. Couto⁷, D. Santos Faria⁸, T. Rocha⁹, R. Freitas¹⁰, A.R. Cunha¹¹, A. Heaney¹², S. McKenna¹², J.A. Pereira da Silva^{1,2}. ¹Faculdade de Medicina da Universidade de Coimbra; ²Centro Hospitalar Universitário de Coimbra, Coimbra; ³Instituto Português de Reumatologia, Lisboa; ⁴Centro Hospitalar Vila Nova de Gaia, Vila Nova de Gaia; ⁵Hospital do Divino Espírito Santo, S. Miguel; ⁶Centro Hospitalar Universitário do Algarve, Faro; ⁷Centro Hospitalar Tondela-Viseu, Viseu; ⁸Unidade Local de Saúde Alto Minho, Ponte de Lima; ⁹Centro Hospitalar São João, Porto; ¹⁰Hospital Garcia de Orta, Almada; ¹¹Centro Hospitalar Baixo Vouga, Aveiro, Portugal; ¹²Galen Research, London, UK

Background: Rheumatoid Arthritis (RA) is a chronic inflammatory disease that has a major impact on patients' quality of life. The Rheumatoid Arthritis Quality of Life Questionnaire (RAQoL) is a patient-centric outcome measure, specific to RA. The measure has not previously been available for use with Portuguese RA patients.

Objectives: To produce a Portuguese version of the RAQoL that is acceptable to Portuguese patients and demonstrates sound psychometric properties.

Methods: The dual panel methodology was used to translate the UK RAQoL into Portuguese. This involved conducting a bilingual panel (providing the initial translation into Portuguese) followed by a lay panel (where items are assessed for comprehension and acceptability). Cognitive debriefing interviews were conducted with Portuguese RA patients to determine the face and content validity of the translated scale. A large-scale postal validation survey was carried out to establish the psychometric properties of the Portuguese RAQoL. The measure was administered on two occasions to RA patients, alongside a comparator instrument – the Nottingham Health Profile (NHP).

Internal consistency was assessed using Cronbach's alpha coefficient. Spearman's Rank correlation coefficient was employed to assess test-retest reliability. Convergent validity was tested by correlating RAQoL scores with those on the NHP sections. Known group validity was assessed using non-parametric tests for independent samples. This involved determining the ability of the RAQoL to distinguish between patients that differed according to their self-perceived severity of RA and general health.

Results: The translation panels produced a Portuguese version of the RAQoL that was easily understood and considered natural by native speakers. Interviewees considered the new language version to be relevant and appropriate. One hundred and seventy-eight RA patients (82% female) took part in the postal validation survey with a mean age of 56.6 (range 25 to 79) years. The Portuguese RAQoL demonstrated excellent internal consistency (Cronbach's $\alpha=0.95$) and test-retest reliability ($r=0.92$), indicating that the measure produces low levels of random measurement error.

RAQoL scores correlated most strongly with scores on the NHP Physical mobility scale ($r=0.77$) and showed moderately strong correlations with the Emotional reactions, Pain and Energy level section scores.

Non-parametric tests for independent samples demonstrated significant differences in RAQoL scores between patients who differed according to their self-perceived RA severity ($p<0.001$) and general health ($p<0.001$).

Conclusions: The Portuguese version of the RAQoL was found to be a comprehensive, reliable and valid questionnaire. The new language version is recommended for use in routine clinical practice and for research purposes, to assess quality of life in Portuguese RA patients.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5989

AB1365 PROPROTEIN CONVERTASE SUBTILISIN/KEXIN TYPE 9 (PCSK9) IN PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS/LUPUS NEPHRITIS

C. Fang, T. Luo, L. Lin. Second Affiliated Hospital of Fujian Medical University, Quanzhou, China

Background: SLE patients have a tendency of accelerated atherosclerosis (AS) which can only partly be explained by traditional risk factors for cardiovascular disease. Proprotein convertase subtilisin/kexin type 9 (PCSK9), which is a protease associated with cardiovascular risk that regulates both cholesterol metabolism and inflammatory reaction, had been regarded as a highly promising therapeutic target for cardiovascular disease.¹ Recent study had demonstrated that SLE patients with lupus nephritis (LN) had much higher risk of atherosclerosis.²

Objectives: To assess serum PCSK9 concentrations and the possible factors linked with PCSK9 variation in SLE/LN patients.

Methods: 47 SLE patients and 30 healthy controls were included. Traditional cardiovascular risk factors were compared. According to cIMT, SLE patients were divided into two subgroups (SLE-AS subgroup and SLE-NonAS subgroup, cut-off point is 1.0 mm). PCSK9 concentrations were compared between: ¹SLE patients

and controls;²SLE-AS subgroup and SLE-NonAS subgroup;³SLE patients with and without lupus nephritis(LN). The correlational analyses between PCSK9 levels and disease parameters were undergone.

Results: The differences of lipids parameters, body mass index (BMI), uric acid (UA) between SLE group and controls had no statistical significance. Even so, the ratio of cIMT thickening were higher in SLE patients, when compared with healthy controls (23.40% versus 6.67%, $p=0.05$). Serum PCSK9 levels were also significantly elevated in SLE patients than controls (median of PCSK9 level: 390.53 ng/ml versus 292.44 ng/ml, $p<0.05$). Patients in SLE-AS subgroup had even higher PCSK9 and C-reactive protein(CRP) levels than those in SLE-NonAS subgroup (median of PCSK9 level: 516.41 ng/ml versus 364.47 ng/ml, $p<0.05$; median of CRP level: 4.56 mg/L versus 1.05 mg/L, $p<0.01$) (figure 1). PCSK9 levels correlated with CRP levels, but not with age, disease activity, lipids characteristics, BMI or UA levels, particularly in female patients (table 1). Despite of no statistical significance, the ratio of lupus nephritis were higher in SLE-AS subgroup than those in SLE-NonAS subgroup (63.64% versus 44.44%, $p>0.05$). SLE patients with LN had higher PCSK9 concentrations than those without LN evidence(509.53 $\pm$ 131.69 ng/ml versus 332.02 $\pm$ 92.72 ng/ml, $p<0.001$).

Abstract AB1365 – Table 1. Correlational analysis of PCSK9 levels and disease parameters in SLE patients.

Variables	r_s	p -value
Age(years)	-0.014	0.927
SLEDAI	0.092	0.539
BMI(kg/m ²)	-0.120	0.420
Cholesterol(mmol/l)	-0.012	0.935
LDL cholesterol(mmol/l)	0.002	0.989
ApoA1(g/l)	0.011	0.943
ApoB(g/l)	0.181	0.223
Triglycerides(mmol/l)	-0.020	0.896
HDL cholesterol(mmol/l)	-0.112	0.453
Fasting blood glucose(mmol/l)	-0.193	0.193
Uric acid(μ mol/l)	-0.079	0.599
CRP(mg/l)	0.351	0.016
CRP(mg/l) in female patients	0.487	0.001

Abstract AB1365 – Figure 1

Conclusions: Elevation of PCSK9 concentrations can be observed in SLE patients, especially in those with LN or atherosclerosis. PCSK9. PCSK9 is probably linked with low-grade inflammation participating in the pathogenesis of atherosclerosis in SLE/LN patients.

REFERENCES:

- [1] Robinson JG, Farnier M, Krempf M, et al. Efficacy and safety of aliroc-mab in reducing lipids and cardiovascular events[J]. *N Engl J Med*. 2015 Apr 16;372(16):1489–99.

- [2] McHugh J. Systemic lupus erythematosus: Atherosclerosis confined to patients with nephritis[J]. *Nat Rev Rheumatol*. 2017 Jun;13(6):322.

Acknowledgements: The authors thank Qiulan Li and Hongzhi Gao for the excellent technical assistance.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1622

AB1366 RADIOISOTOPIC SYNOVIORTHESIS IN THE CONTROL OF REFRACTORY SYNOVITIS IN CASTILLA-LA MANCHA (A REGION OF SPAIN). AN EXPERIENCE OF 10 YEARS

D. Castro Corredor. RHEUMATOLOGY SERVICE, HOSPITAL GENERAL UNIVERSITARIO DE CIUDAD REAL, CIUDAD REAL, Spain

Background: Our hospital is the reference hospital in Nuclear Medicine for the realisation of radioisotopic synoviorthesis for all of Castilla-La Mancha (a region of Spain).

Objectives: To describe the experience in the performance of radioisotopic synoviorthesis in arthritis refractory to other treatments in our hospital.

Methods: Observational, descriptive and cross-sectional study protocolised through the review of the database of radioisotopic synoviorthesis performed between 2007 and 2017. Previous clinical data were collected (age, sex, pathology, previous treatments, previous infiltration and affected joint) and evolutive at 6 months after administering the isotope. An Excel database was created for a frequency analysis with SPSS 21.

Results: 30 radiosynovitis were performed with the most frequent pathologies, in this order: pigmented villonodular synovitis (40%), rheumatoid arthritis (23.3%), spondyloarthritis (13.3%), osteoarthritis (10%) and nonspecific arthritis (6.7%), followed by systemic lupus erythematosus and gout. After 6 months, 56.7% of patients improved compared to 36.7% who remained the same. Likewise, none of them presented complications related to the procedure. A 6.6% of patients were lost to follow-up.

Conclusions: In patients with episodes of recurrent arthritis with associated joint effusion in one or two joints, refractory to systemic treatments, to local infiltrations with corticosteroids and in those patients in whom other treatments may be contraindicated, we must consider the possibility of performing an isotope radiosynoviorthesis, as it is a simple, safe technique with a success rate of more than 50%.

REFERENCES:

- [1] Delbarre f, Cayla, Menkes C. La synoviorthese par les radioisotopes. *Presse Med* 1968; 76:1045
- [2] J. Román Ivorra, J. Chismol Abad, M. Morales Suárez Varela, M. Ureña, J. Ribes Iborra, R. Sopena Monforte. La sinoviortesis radioisotópica en la artritis reumatoide. Estudio de 108 casos. *Revista Española de Medicina Nuclear*. 2000;19(4):275–278.
- [3] Taylor W, Corkill M, Rajapasko C. A retrospective review of yttrium-90 synovectomy in the treatment of knee arthritis. *Br J Rheumatol* 1997; 36:1100–1105.
- [4] Harbert C. Radiocolloid joint therapy in joint diseases. In Habert JC. Edition. Nuclear medicine therapy. New York. Thieme Medical; 1987:169–186.
- [5] Stevenson AC, Bedford J, Hill AGS, Hill H. Chromosome damage in patients who have had intraarticular injections of radioactive gold. *Lancet* 1937b; 2:837839.
- [6] Román J, Chismol J, Morales M, Ureña M, Ribes J, Sopena R. La sinoviortesis radioisotópica en la artritis reumatoide. Estudio de 108 casos. *Rev Esp Med Nucl* 2000; 19:275–278.
- [7] Siegel M, Siegel H, Luck J. Radiosynovectomy's clinical applications and cost effectiveness: A review. *Sem Nucl Med* 1997;27:364–371.
- [8] Mulero J. Sinovectomía quirúrgica y sinovectomía isotópica. *Med Clin (Barc)* 1985; 85:835–837.
- [9] Jones G. Yttrium synovectomy: a meta-analysis of the literature. *Aust NZ J Med* 1993; 23:272–275.
- [10] Heuft-Dorenbosch L, de Vet H, van der Linden S. Yttrium radiosynoviorthesis in the treatment of knee arthritis in rheumatoid arthritis: a systematic review. *Ann Rheum Dis* 2000;59:583–586.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2785

AB1367

DIAGNOSTIC ASSOCIATION BETWEEN ELASTIN AND ELASTASE ANTIBODIES WITH CARDIOVASCULAR SYSTEM INVOLVEMENT IN SYSTEMIC SCLERODERMA PATIENTS

I.P. Gontar¹, O.A. Rusanova², O.I. Emelyanova¹, L.A. Maslakova¹, I. A. Zborovskaya¹. ¹Research Institute of Clinical and Experimental Rheumatology named after A.B. Zborovsky; ²Research Institute of Clinical and Experimental Rheumatology named after A.B. Zborovsky, Volgograd, Russian Federation

Background: Large amounts of elastin are contained in vascular walls and in the cardiac valves. Elastin and elastase antibodies are predictors, *sui generis*, of vascular disease development in systemic scleroderma. The mechanism of direct proatherogenic effect of antibodies consists in stimulation of adhesion molecules and cytokine synthesis, oxLDL capture, induction of serine proteinase cascade by the coagulation system. In systemic rheumatoid disease, autoimmune inflammation is nowadays regarded as one of risk factors for the development of early atherosclerosis, disturbance of the structure and elasticity of vascular wall and of related cardiovascular conditions.

Objectives: Studying the effect of elastin and elastase autoantibody production on the cardiovascular system in systemic scleroderma (SSD) patients.

Methods: 42 patients hospitalised at municipal hospital 25 with diagnosis of SSD verified by ^{ARA} diagnostic criteria (1980) the diagnosis was considered firm if the patient showed one major and two minor criteria in any combination, simultaneously or consecutively, regardless of the time of their onset. The systemic scleroderma patients included 11 men and 31 women aged 22–72. The mean age of patients was 44.1±15.4 years. 30 healthy donors from the Volgograd hemotransfusion centre served as controls. Antibodies to elastin and elastase were determined in the blood serum using indirect enzyme immunoassay with magnetocontrollable adsorbents based on polyacrylamide granules according to the original technique by Gontar et al (1990).

Results: SSD patients showed considerable increase in the rate of elastase (52%) and elastin (38%) antibodies formation, in comparison with the controls. The upper normal limit of elastin antibody was within the range of 0.131 optical density units, elastase antibodies – 0.131 optical density units. In SSD patients elastin antibodies amounted to 0.125±0.068 optical density units. The titer of elastase antibodies was 0.143±0.071 optical density units. Healthy individuals did not show any elastin or elastase antibodies. An elevated antibody titer was associated with heart and vessels lesion in 47% of patients with SSD. In 20 patients of the studied group we revealed cardiovascular disease (IHD, macrofocal cardiovascular sclerosis with false infarction changes, chronic cardiac failure, and aorta atherosclerosis).

Conclusions: Among the patients we examined, 47% showed cardiovascular lesion associated with elevated elastin and elastase antibodies. This fact indicates that in systemic rheumatic disease, autoimmune inflammation is a risk factor for the development of early atherosclerosis, and of related cardiovascular conditions. Elastin and elastase antibodies are predictors, *sui generis*, of the development of vascular disease in patients with SSD.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4470

AB1368

THE COMPARATIVE RESPONSIVENESS OF HOSPITAL UNIVERSITARIO PRINCESA INDEX AND OTHER COMPOSITE INDICES FOR ASSESSING RHEUMATOID ARTHRITIS ACTIVITY

I. Gonzalez-Alvaro¹, I. Castrejon², L. Carmona³. ¹Rheumatology, Hospital Universitario La Princesa. IIS-IP, Madrid, Spain; ²Rheumatology, Rush University, Chicago, USA; ³Rheumatology, InMusc, Madrid, Spain

Background: HUPI was developed with data from the PEARL study (Princess Early Arthritis Register Longitudinal study) as an easy to calculate index, which avoided the gender bias affecting DAS28 and SDAI. In addition, it can be calculated either with erythrocyte sedimentation rate (ESR), C reactive protein (CRP) or both. Response criteria based on HUPI have also been developed¹.

Objectives: To evaluate the responsiveness of the Hospital Universitario La Princesa Index (HUPI) comparatively to the traditional composite indices to assess disease activity in rheumatoid arthritis (RA), and to compare the performance of HUPI-based response criteria with that of the EULAR response criteria.

Methods: Post-hoc analyses were performed using data from the following studies: ACT-RAY (clinical trial), PROAR (early RA cohort) and EMECAR (pre-biologic era long term RA cohort). Responsiveness was evaluated by: 1) comparing change from baseline (Δ) of HUPI with Δ in other scores by calculating correlation coefficients; 2) calculating standardised effect sizes. The accuracy of response by HUPI and by EULAR criteria was analysed using linear regressions in which the dependent variable was change in global assessment by physician (Δ GDA-Phy).

Results: Δ HUPI correlation with change in all other indices ranged from 0.387 to 0.791; HUPI's standardised effect size was larger than those from the other indices in each database used. In ACT-RAY, depending on visit between 65% and

80% of patients were equally classified by HUPI and EULAR response criteria. However, HUPI criteria were slightly more stringent, with higher percentage of patients classified as non-responder, especially at early visits. HUPI response criteria showed a slightly higher accuracy than EULAR response criteria when using Δ GDA-Phy as gold standard.

Table 1

Comparison of sensitivity to change HUPI with other composite indices.

	Standardised size effect					Response (% EULAR vs HUPI)		
	HUPI	DAS28-VSG	DAS28-PCR	SDAI	CDAI	None	Moderate	Good
3 months	2'3	2'18	2'2	1'99*	1'67*	9 vs 22	53 vs 42	38 vs 36
6 months	3'13	2'93	2'85	2'56*	2'27*	3 vs 9'5	34 vs 29	63 vs 60'5
12 months	3'69	3'45	3'28*	2'87*	2'65*	2 vs 6'5	23 vs 20	75 vs 73'5

*p<0.05 respect HUPI

Conclusions: HUPI shows good responsiveness in each studied scenario (clinical trial, early RA cohort, and established RA cohort). Response criteria by HUPI seem more stringent than EULAR's.

REFERENCE:

- Gonzalez-Alvaro I, Castrejon I, Ortiz AM, Toledano E, Castaneda S, Garcia-Vadillo A, et al. Cut-Offs and Response Criteria for the Hospital Universitario La Princesa Index (HUPI) and Their Comparison to Widely-Used Indices of Disease Activity in Rheumatoid Arthritis. PLoS One. 2016;11(9): e0161727.

Acknowledgements: We are indebted to the Roche laboratory for the transfer of the ACT-RAY trial data and to the Spanish Rheumatology Society for the transfer of data from the EMECAR and PROAR projects. Without the work of all the researchers who participated in these three projects, this work could not have been developed. This work has been supported by project PI14/00442 to IG-A from the Ministry of Economy and Competitiveness (Carlos III Health Institute) and co-financed with the European Fund for Regional Development (FEDER).

Disclosure of Interest: I. Gonzalez-Alvaro Grant/research support from: UCB, Roche., Consultant for: Lilly, Pfizer, BMS, Sanofi, Speakers bureau: Abbvie, Lilly, UCB, SER, I. Castrejon: None declared, L. Carmona: None declared

DOI: 10.1136/annrheumdis-2018-eular.4479

AB1369

ELEVATED CA-125 IN IGG4-RELATED MESENTERIC DISEASE: A RED HERRING? A SYSTEMATIC LITERATURE REVIEW

I. Panayotidis¹, M. Michaud Maturana¹, S. Psarelis², E. Nikiphorou³. ¹Medical School, University College London (UCL), London, UK; ²Rheumatology Department, Nicosia General Hospital, Nicosia, Cyprus; ³Department of Academic Rheumatology, King's College London, London, UK

Background: Mesenteric panniculitis (MP), a rare fibrotic inflammatory disease of the bowel mesentery, can be a rare manifestation of IgG4-related disease (IgG4-RD). IgG4-RD is a chronic inflammatory disease most commonly affecting the pancreas, characterised by infiltration of IgG4-positive plasma cells and lymphocytes into various organs.¹

We recently encountered a male patient with IgG4-related MP who was incidentally found to have a very high level of CA-125, which correlated with CRP levels and normalised after steroid treatment. This prompted a systematic literature review (SLR) to better understand this unexpected phenomenon.

Objectives: To investigate for associations between CA-125, MP and/or IgG4-RD, understand possible common pathophysiological mechanisms and explore potential clinical implications.

Methods: The SLR was performed using MEDLINE, EMBASE, Web of Science, and Scopus, looking for literature on either MP and CA-125 or IgG4-RD and CA-125 up to January 8 2018, using a comprehensive search strategy with relevant mesh terms and keywords linked to the above broad categories. Literature screening was performed by two independent reviewers.

Results: 24 unique citations were found, of which 13 were unanimously identified as relevant by the two reviewers. The final selected articles included: 8 case reports, 3 conference abstracts of case reports, 1 cohort study of 22 patients, and a retrospective study of 7 patients (table 1). CA-125 was raised in 22/40 patients in the identified reports (shown in red), including males, and was often the only elevated tumour marker (yellow). We also report on the presence of effusions (blue), as this may be linked to the causal mechanism.

Table 1 The clinical and laboratory characteristics of cases of IgG4-RD and MP

Article	Patient demographics	Main diagnosis	Serum CA-125 (U/mL)	Other tumour markers	Presence of effusions
Tong et al. (2017)	43 F	IgG4-RD	↑ (1065)	CA 19-9 ↑, AFP ↑, CEA ↔	Ascites, pleural and pericardial effusions
Khosla et al. (2016)	72 M	AIP	↑ (196.4)	CA 19-9 ↔, CEA ↔	N/A
Thomas et al. (2015)	46 F	AIH, AI myositis	↑ (417)	CA 19-9 ↑	Ascites
Zhou, Zeng (2014)	24 F	IgG4-RD (lung)	↔	CA 19-9 ↔, CEA ↔, CA-153 ↔	Pleural effusion
Fengjing et al. (2014)	Cohort study of 22 patients	IgG4-RD	↑ in 12/22	CA 19-9 ↑ in 6/22	N/A
Cheng et al. (2013)	45 M	AIP	↑ (77.5)	CA 19-9 ↔	Ascites, pleural effusions
Liu et al. (2012)	41 M	IgG4-RD (lung)	↔	CA 19-9 ↔, CEA ↔, PSA ↔, NSE ↔	No
Graham, Harvin (2016)	66 F	IgG4-RD (SM)	↔	CA 19-9 ↔, CEA ↔	Ascites
Ghadir et al. (2012)	70 M	AIP	↔	CA 19-9 ↔, CEA ↔	N/A
Dogaru et al. (2015)	60 F	SM	↑ (295.5)	CA 19-9 ↔, AFP ↔, CEA ↔	N/A
Buyukbayrak et al. (2011)	38 F	SM	↑ (44.5)	CA 19-9 ↔, AFP ↔, CEA ↔, CA 153 ↔	Ascites
Malter et al. (2017)	57 F	MP	↑ (630)	N/A	Ascites
Xiaoting et al. (2017)	Retrospective study of 7 patients (3 F, 4 M)	IgG4-RD (lung)	↑ in 3/3 F (46.54, 156, 85.58)	CA 19-9 and CEA ↑ in 1/3 F	No
Current report	56 M	IgG4-related MP	↑ (1358)	CA 19-9 ↔, AFP ↔, CEA ↔	Pleural effusion in 1/4 M

Abstract AB1369 – Figure 1

AI: Autoimmune, **AIH:** Autoimmune Hepatitis, **AIP:** Autoimmune Pancreatitis, **SM:** Sclerosing Mesenteritis, **↑:** Raised, **↔:** Normal

Conclusions: To our knowledge, this is the first SLR exploring the association between CA-125 and IgG4-RD and MP. Despite the small sample, our results do indicate that CA-125 was raised in more than 50% of reported cases, the majority of which also had some kind of effusion. Although traditionally a marker of ovarian cancer, this report highlights that a raised CA-125 can be found in other, non-malignant, inflammatory conditions, and potentially correlates with inflammatory burden. CA-125 in this setting may thus represent a useful biomarker and have a role in monitoring treatment response.

REFERENCE:

- [1] Minato H, Shimizu J, Arano Y, et al. IgG4-related sclerosing mesenteritis: a rare mesenteric disease of unknown etiology. *Pathol Int.* 2012;62:281–6.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2221

AB1370 PATIENT REPORTED OUTCOMES MEASURES FOR FIBROMYALGIA: A REVIEW

J.B. Negrón, L. Carmona. *Instituto de Salud Musculoesquelética, Madrid, Spain*

Background: Persons with fibromyalgia (FM) suffer from numerous symptoms with various levels of intensity, such as widespread pain, fatigue, cognitive dysfunction, non-restorative sleep, depression, and anxiety among others. The patient's perspective has been recognised by Outcome Measure in Rheumatology (OMERACT) as a key assessment in FM. For that reason, some patient reported outcomes measures (PROMs) have been developed. PROMs allow the comparison of a specific component of the disease in a patient through time, and also provide the opportunity to health professionals to compare patients between them in clinical trials. However, some PROMs are used over others despite a limited validation process. In addition, different versions of the same PROMs can co-exist, what can lead to uncertainty when selecting the adequate instrument.

Objectives: To identify the existing PROMs for FM and analyse their psychometric properties.

Methods: The authors performed a comprehensive search in electronic databases (Medline, Embase, and Cochrane) in order to identify validation studies of PROMs for FM. Studies published between January 1990 and November 2017 were included. Generic PROMs and validation of diagnostic criteria and related screening tools were not considered as PROMs and were excluded. Information was gathered based on the Consensus-based Standards for the selection of health Measurement Instruments (COSMIN) checklist manual.

Results: The electronic search produced 1832 records. After screening, a total of 48 studies containing 16 PROMs for FM were included. The PROMs included address different constructs of the disease, Fibromyalgia Impact Questionnaire (FIQ), Revised Fibromyalgia Impact Questionnaire (FIQR), Combined Index of Symptom Severity, Combined Index of Severity of Fibromyalgia, Comprehensive Rating Scale for Fibromyalgia Symptomatology, Fibromyalgia Assessment Status, Fibromyalgia Bladder Index, Fibromyalgia Burden Assessment, Fibromyalgia Health Assessment Questionnaire, Fibromyalgia Impact Questionnaire+Visual Analogue Scales, Fibromyalgia Participation Questionnaire, Fibromyalgia Sleep Diary, Multidimensional Inventory of Subjective Cognitive Impairment, Multidimensional Patient Reported Outcome Measures Questionnaire, PROMIS Fatigue FM Profile, and Multidimensional daily diary of fatigue-fibromyalgia-17 items.

Almost all PROMs have adequate content validity. Three PROMs do not report construct validity; seven do not report reliability, and six do not report internal consistency. Only three PROMs evaluate criterion validity and three responsiveness. The FIQ and the FIQR are the PROMs more widely cross-cultural validated with 18 and 13 adaptations respectively.

Conclusions: PROMs for FM have, in general, only partial validation of their psychometric properties. Validation of an instrument is a continuous process in which quality is more important than quantity. Instead of creating new PROMs for FM, future works should focus on completing missing parts of the validation process of existing ones. In addition, cultural adaptations and translations of the available PROMs should be prioritised in order to offer researchers across the globe a tool-box of options in where they can choose the best PROMs to address their objectives in a highly subjective syndrome as FM.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2175

AB1371 CLINICAL SIGNIFICANCE OF KL-6 AND SP-D AS SERUM MARKERS FOR INTERSTITIAL LUNG DISEASE IN PATIENTS WITH CONNECTIVE TISSUE DISEASE

K.-E. Lee, J.-H. Kang, D.-J. Park, S.-S. Lee. *CHONNAM NATIONAL UNIVERSITY MEDICAL SCHOOL AND HOSPITAL, Gwangju, Korea, Republic of Ireland*

Objectives: To evaluate the association between serum levels of Krebs von den Lungen-6 (KL-6) and surfactant protein-D (SP-D) and the clinical manifestations and severity of interstitial lung disease (ILD) in patients with connective tissue disease (CTD).

Methods: Eighty patients with various CTDs were included, as follows: 33 with rheumatoid arthritis (RA), 19 with systemic lupus erythematosus (SLE), 10 with systemic sclerosis (SSc), 9 with Sjögren's syndrome (SS), and 9 with inflammatory myositis. KL-6 and SP-D levels were measured using an enzyme-linked immunosorbent assay and defined as abnormal if KL-6 ≥ 500 U/mL and SP-D ≥ 110 mg/mL. All patients were simultaneously evaluated for parameters related to disease activity using laboratory tests and a pulmonary function test, and interstitial lung abnormalities (ILA) using chest computed tomography (CT). Patients were subclassified according to ILA score: 0 for no ILD, 1 for indeterminate ILD, 2 for mild ILD, and 3 for advanced ILD based on chest CT scans.

Results: In all, 29 patients had radiologically advanced ILD, 18 had mild ILD, 18 had indeterminate ILD, and 15 had no ILD. A higher ILA score was associated with more severe dyspnea, and decreased volume and percent of functional vital capacity, forced expiratory volume in 1 s, and diffusion capacity of carbon monoxide. As clinical manifestations, a higher ILA score was associated with a higher GAP index but not with the parameters of disease activity. A higher ILA score was associated with higher levels of KL-6 and SP-D and a higher percentage of subjects with abnormal levels, and this was more pronounced in SLE, SSc, and SS than in RA.

Conclusions: Serum levels of KL-6 and SP-D are associated with the radiological severity of ILD. Hence, these can serve as markers for ILD severity, especially in SLE, SSc, and SS.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2742

AB1372 TOWARDS REFORMING THE TAXONOMY OF HUMAN DISEASE: THE PRECISESADS CROSS SECTIONAL STUDY

L. LAIGLE¹, L. BERETTA², J. WOJICK³, J. MAROVAC⁴, J.-O. PERS⁵, B. LAUWERYS⁶, J. FROSTEGARD⁷, F. MAC DONALD⁸, M. JUAREZ⁴, R. BENSCHOP⁹, E. DOW⁹, S. RAO¹⁰, C. CHAMBERLAIN⁴, J. MARTIN¹¹, M. E. ALARCON-RIQUELME¹², on behalf of the Clinical Consortium. ¹Center for Therapeutic Innovation – Immuno-Inflammatory Disease, I.R.I.S., SURESNES CEDEX, France; ²Scleroderma Unit, Fondazione IRCCS Policlinico, Milan, Italy; ³QuartzBio SA, Geneva, Switzerland; ⁴UCB Biopharma, Slough, UK; ⁵UMR1227, INSERM, Brest, France; ⁶Cliniques Universitaires Saint-Luc and Université Catholique de Louvain, Brussels, Belgium; ⁷Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; ⁸Bayer AG, Berlin, Germany; ⁹Eli Lilly and Company, Indianapolis; ¹⁰Sanofi-Aventis Research and Development, Bridgewater, USA; ¹¹Instituto de Parasitología y Biomedicina López-Neyra; ¹²GENYO, Pfizer-University, Granada, Spain

Objectives: The PRECISESADS project aims at using OMICs, and bioinformatics to identify new classifications for systemic autoimmune diseases (SADs) known to share common pathophysiological mechanisms in view of personalised treatments. Multi OMICs parameters collected in addition to routine clinical data in a cross-sectional study involving patients suffering from systemic lupus erythematosus (SLE), systemic sclerosis (SSc), Sjögren's syndrome (Sjs), rheumatoid arthritis (RA), primary antiphospholipid syndrome (PAPs), mixed connective