analyse the association between these physical functioning measures and the assessment of affect and life satisfaction.

Results: We found independent associations of the objective measures but not the subjective assessments of physical activity with positive affect and satisfaction with life (both, $p < 0.01$ and adjusted $R^2 > 0.06$) and of sedentary time with positive affect ($p < 0.02$ and adjusted $R^2 > 0.03$). Moreover, we observed consistent and independent associations of both the objective and subjective dimensions of physical fitness with all the components of subjective well-being (all, $p < 0.01$ and adjusted R^2 ranged from 0.02 to 0.05).

Conclusions: Both objective and subjective measures of physical activity and sedentary behaviour independently impact affect and subjective well-being. Strategies to enhance both dimensions of physical fitness may be a promising approach for improving the subjective well-being in fibromyalgia.

REFERENCES:

1. Estévez-López F, et al. *Disabil Rehabil* 2018;40(3):329–337.
2. Estévez-López F, et al. *Rheumatology (Oxford)* 2017;56(11):2015–2024.

Acknowledgements: This study was funded by the Spanish Ministries of Economy and Competitiveness [I+D+I DEP2013–40908-R, I+D+I DEP2010–15639, BES-2014–067612] and Education [FPU15/00002].

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6370

FRI0711-HPR KINESIOPHOBIA IN ADULT PATIENTS WITH FAMILIAL MEDITERRANEAN FEVER: ASSOCIATION WITH PHYSICAL ACTIVITY, DEPRESSION AND ANXIETY

I. Karanki, E.I. Günaydin, A. Çetinkaya, S. Saka, M.G. Yavuzer. *Haliç University, İstanbul, Turkey*

Background: Familial Mediterranean fever (FMF) characterized by pain and inflammation with fever which requires a lifelong treatment¹. All of this causes psychosocial influences and also causes kinesiophobia, which is defined as avoidance of physical activity and fear of repetition of the pain. As it may affect the levels of physical activity and functional capacity, it is important to identify factors related to kinesiophobia in patients with FMF.

Objectives: The aim of this study was to investigate the existence of kinesiophobia and related factors such as depression, anxiety and physical activity in adult patients with FMF.

Methods: 38 subjects with FMF (15 male) from Society of Behçet & FMF Diseases included in the study. The demographic characteristics of the participants were recorded. The 'Tampa Scale for Kinesiophobia (TSK)²', 'International Physical Activity Questionnaire (IPAQ)³', 'Hospital Anxiety and Depression Scale (HADS)⁴' and 'Fatigue Severity Scale (FSS)⁵' were used to assess associated factors.

Results: Demographic and disease characteristics of the participants were presented in the table 1. Only 9 subjects reported regular exercise. 86% of the subjects had a score over 37 in TSK representing high kinesiophobia⁶. TSK scores were positively correlated with HAD-D ($r=0.530$; $p=0.001$) and FSS ($r=0.340$; $p=0.035$). On the other hand, age, disease duration, body mass index, HAD-A and IPAQ scores were not correlated with TSK. Gender or exercise routine had no effect on TSK.

Table 1 Demographic and disease characteristics

	Mean±Std	Min.	Max.
Age (year)	37.05±8.71	19	57
Disease duration (years)	13.11±11.14	1	46
BMI (kg/cm ²)	24.5±4.43	16.65	38.05
IPAQ	6942.5 ±2520.4	3018	14580
TSK	44.16±5.06	33	54
HAD-A	9.84±4.43	2	20
HAD-D	8.82±4.70	1	20
FSS	56.45±13.05	23	70

Conclusions: Kinesiophobia is very common and associated with depression and fatigue in patients with FMF. Limited number of participants reported regular exercise habit which should be added to treatment programs to increase physical activity and functional capacity in patients with FMF.

REFERENCES:

1. Makay, Emiroğlu N, Unsal E. Depression and anxiety in children and adolescents with familial Mediterranean fever. *Clin Rheumatol* 2010 Apr;29(4):375–9.
2. Yılmaz OT, Yakut Y, Uygur F, Uluğ N. Turkish version of the Tampa Scale for Kinesiophobia and its test-retest reliability *Fizyoterapi Rehabilitasyon* 2011;22(1):44–49.
3. Sağlam M, Arıkan H, Savcı S, Inal-Ince D, Boşnak-Güçlü M, Karabulut E, et al. International physical activity questionnaire: reliability and validity of the Turkish version. *Perceptual and motor skills* 2010;111(1):278–84.
4. Aydemir O, Güvenir T, Küey L, Kültür S. Reliability and Validity of the Turkish version of Hospital Anxiety and Depression Scale. *Turkish journal of psychiatry* 1997;8(4):280–287.
5. Armutlu K, Korkmaz NC, Keser I, Sumbuloglu V, Akbiyik DI, Guney Z, Karabudak R. The validity and reliability of the Fatigue Severity Scale in Turkish multiple sclerosis patients. *Int J Rehabil Res*. 2007 Mar;30(1):81–5.
6. Vlaeyen J, Kole-Snijders AMJ, Rotteveel AM, Ruesnik R, Heuts PHTG. The role of fear of movement((re)injury in pain disability. *J Occup Rehab* 1995;5(4):235–51.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3695

FRI0712-HPR THE EXPERIENCE OF PATIENTS WITH DIFFERENT CHRONIC DISEASES WITH HEALTH CARE. A SURVEY WITH THE IEXPAC INSTRUMENT

J. de Toro¹, L. Cea-Calvo², M.L. García-Vivar³, L. Pantoja⁴, C. Lerín⁵, S. García-Díaz⁶, M.J. Galindo⁷, I. Marín-Jiménez⁸, M. Mosquera², M.J. Arteaga², D. Orozco-Beltrán⁹. ¹Rheumatology Department, H. U. A Coruña, Universidad A Coruña, INIBIC, A Coruña; ²Merck Sharp & Dohme, Madrid; ³Rheumatology Department, Basurto University Hospital, Bilbao; ⁴Rheumatology Department, H. El Bierzo, Ponferrada; ⁵Rheumatology Department, Manacor Hospital, Palma de Mallorca; ⁶Rheumatology Department, Moisés Broggi Hospital, Barcelona; ⁷Internal Medicine Department, H. Clínico de Valencia, Valencia; ⁸Gastroenterology Department, H. U. Gregorio Marañón, Madrid; ⁹Department of Medicine, Miguel Hernández University, Sant Joan, Alicante, Spain

Background: Insights from patients are important for health care planning

Objectives: In this study we describe the perception of patients with 4 different chronic conditions with health care in Spain through the IEXPAC scale ("Instrument to Evaluate the Experience of Patients with Chronic diseases"), a scale developed and validated in Spain.

Methods: The IEXPAC scale (<http://www.iemac.es/iepac/>) was developed and validated in Spain by health care professional and social organizations, experts in quality of health care and chronic patients. The scale is structured in 12 items with Likert responses from "always" to "never", yields a score from 0 (worst experience) to 10 (best experience), and identifies aspects of health care needing improvement. A survey was handed to patients needing care in at least two different levels (i.e. primary care and hospital) and with one of the following chronic conditions A) Rheumatic diseases, B) Inflammatory bowel disease (IBD), C) Human immunodeficiency virus infection (HIV) and D) Diabetes mellitus (DM) plus cardiovascular or renal chronic disease. Patients completed the survey at home and responded by pre-paid mail.

Results: 2474 patients received the survey, 1618 (65.4%) returned it (359 with rheumatic disease, 341 with IBD, 467 with HIV infection, 451 with DM, mean age 56 years, 41% women). Only 6.1% were affiliated to a patients association. Patients declared a median of 8 visits (IQR 25–75: 4–15) to primary care or specialty clinics in the last year and 29% had visited an emergency room. In the last 3 years 48% had been hospitalized. Up to 61% reported to search for information on diseases, therapies, lifestyle or diet in webpages, general or social media. Responses to the IEXPAC items (percentages that responded "mostly" + "always" to each item) are displayed in the table. In general, these % were higher in HIV patients, which represents a better experience with health care. In some items, patients with rheumatic diseases scored lower (table 1). Mean IEXPAC score was 6.0 (SD1.8) and was higher in HIV patients (table 1). Worst scores were seen in items related to access or guidance for getting reliable information on health and on social resources, contact with other patients and follow-up after hospital discharge.

Percentage (%) who responded "mostly" + "always"						
IEXPAC item	All patients	Rheumatoid Disease	IBD	HIV	DM	P
They respect my lifestyle	81.5%	76.5%	75.1%	89.6%	81.9%	<0.001
They are coordinated to offer good health care to me	69.3%	60.6%	69.1%	76.8%	73.3%	<0.001
They help me to get information from the Internet	15.0%	12.8%	19.0%	19.8%	8.3%	<0.001
Now I can take care of me better	81.0%	74.3%	79.3%	89.7%	78.3%	<0.001
They ask me and help me to follow my treatment plan	79.8%	73.5%	77.8%	87.6%	78.2%	<0.001
We set goals for a healthy life and better control my	70.1%	63.4%	62.6%	74.7%	76.1%	<0.001
I can use Internet and my mobile phone to consult my medical records	7.2%	7.3%	5.5%	8.6%	7.1%	0.529
They make sure that I take medication correctly	76.0%	72.4%	73.7%	83.5%	72.9%	<0.001
They worry about my welfare	84.3%	79.1%	80.4%	91.5%	83.8%	<0.001
I have been informed on the health and social	41.3%	33.8%	32.3%	52.6%	42.3%	<0.001
They encourage me to talk to other patients	14.9%	10.3%	15.7%	20.4%	12.0%	<0.001
Respond to the following statement only if you have been admitted to the hospital in the last 3 years,						
They care about me when I come home after being in the hospital	30.6%	25.7%	28.9%	33.0%	32.8%	0.205
Global IEXPAC score, mean (SD)	6.0 (1.9)	5.5 (2.0)	5.9 (2.0)	6.6 (1.7)	5.9 (1.9)	<0.001

Conclusions: The IEXPAC questionnaire identified areas of improvement in chronic patients health care in Spain, especially those related with access to reliable information and services, interaction with other patients and continuity of health care after hospital discharge. Patients with HIV infection scored higher, maybe consequence of a more personalized care. In several items, patients with rheumatic diseases scored lower

Acknowledgements: Funded by Merck Sharp & Dohme, Spain, with endorsement of 4 patients associations: CONARTITIS (patients with rheumatoid diseases), ACCU (patients with IBD), SEISIDA (HIV multidisciplinary group), FEDE (patients with DM)

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1588

FRI0713-HPR THE PATIENTS' PERSPECTIVES TOWARDS THE PROVISION OF INFORMATION DURING TRANSITION TO A BIOSIMILAR

J. Kreuk, A. Twisk, J. Meilink, R. Alblas, W. van der Weele, R.C. Hebing, J. Ruwaard, M. l'Ami, Z. Layegh, G. Wolbink. *Amsterdam Rheumatology and Immunology Center Reade, Amsterdam, The Netherlands*

Background: In July 2015, at the Amsterdam Rheumatology and Immunology Centre, location Reade, 93% of the patients with rheumatoid arthritis, treated with reference product infliximab switched to the biosimilar infliximab. This transition was needed to enable cost savings. Patients gained information via several ways, where the rheumatology nurses had a central role in informing the patients. Patients' perspectives are warranted to investigate the effect of the transition to the biosimilar.

Objectives: To investigate patients' perspectives towards the provided information regarding the switch to the biosimilar.

Methods: Consecutive patients treated with reference product infliximab were switched to the biosimilar in the period July 2015 to June 2016 at the Amsterdam Rheumatology and immunology Center, Reade. Patients were informed by a letter about the transition to the biosimilar and were subsequently contacted by a nurse or the pharmacist for additional questions and whether they agreed upon the switch. All patients who were switched from the reference product to the biosimilar were approached at the day care to fill in a questionnaire. In this qualitative questionnaire, patients were asked to evaluate the information provision process and how they gained information about the transition to the biosimilar initially. This was done by the nurses of the day care.

Results: All patients who switched to the biosimilar (n=46) filled in the questionnaire, of which 15 patients scored the information provision as excellent (33%), 25 patients as good (54%), 4 patients as reasonable (9%) and 2 patients found the information sufficient (4%). Furthermore, the majority of patients was initially informed by nurses and rheumatologist prior to the letter that was sent to all patients. In total, 12 patients were initially informed by rheumatology nurse (26%), 12 patients by the rheumatologist (26%). Four patients were informed via the letter that was sent at first (9%) and 3 patients gained the information about the transition otherwise (7%). Fifteen patients gave more than one answer to the question by whom they were informed initially.

A. Patient's perspective

B. Initially informed by:

Conclusions: Patients were satisfied about the information provision process, there were no patients who experienced the information provision insufficient. Next to the rheumatologist, rheumatology nurses played an important role in informing the patients about the transition.

Disclosure of Interest: J. Kreuk: None declared, A. Twisk: None declared, J. Meilink: None declared, R. Alblas: None declared, W. van der Weele: None declared, R. Hebing: None declared, J. Ruwaard: None declared, M. l'Ami: None declared, Z. Layegh: None declared, G. Wolbink Grant/research support from: Pfizer, Speakers bureau: Abbvie, UCB, BMS, Pfizer

DOI: 10.1136/annrheumdis-2018-eular.6884

FRI0714-HPR THE PATIENT'S JOURNEY FROM ONSET OF BACK PAIN TO DIAGNOSIS OF AXIAL SPONDYLOARTHRITIS: RESULTS FROM A QUALITATIVE STUDY

L. Passalent^{1,2,3}, L. Soever^{1,2}, K. Bednis¹, C. Hawke^{1,2}, A. Bidos⁴, J. Bloom^{1,2}, Y. R. Rampersaud^{1,2,3}, N. Haroon^{1,2,3}, R. Inman^{1,2,3}. ¹Toronto Western Hospital; ²University of Toronto; ³Krembil Research Institute; ⁴University Health Network, Toronto, Canada

Background: People with axial spondyloarthritis (SpA) wait an average of 9 years between onset of back pain and time to diagnosis by a rheumatologist (Feldkeller 2000). During this time, patients often experience significant pain and dysfunction, while waiting for appropriate diagnosis and intervention. Understanding patients' perspectives of this diagnostic process is essential in order to optimize early detection and promote appropriate management for axial SpA.

Objectives: The purpose of this study was to understand patients' perspectives of referral and screening practices for axial SpA from onset of back pain to diagnosis by a rheumatologist.

Methods: Semi-structured key informant interviews were conducted with patients diagnosed with axial SpA, based on ASAS criteria, attending a tertiary academic spondylitis clinic. Interviews addressed patients' experiences with screening and referral practices for adults with chronic back pain and suspected axial SpA from symptom onset to diagnosis. All interviews were recorded, transcribed verbatim and evaluated using a compare and contrast analysis by coding groups of words that addressed the research objectives. Two members of the research team undertook this exercise independently and then met to reconcile emergent overarching categories and their respective themes. NVIVO V9 was used to assist with organization of codes.

Results: A total of 10 patient interviews were conducted. 90% of participants were male, mean age 42.8 years ($\pm$ 12.6). The mean duration of back pain prior to diagnosis of axial SpA was 8.9 years ($\pm$ 6.0). The majority of patients (90%) had post-secondary education. Three overarching categories were identified regarding patients' experiences with screening and referral practices for axial SpA from symptom onset to diagnosis and included: 1) "system factors"; 2) "healthcare provider factors" and 3) "patient factors". Themes related to "system factors" included timely and appropriate access to care. Perceived lack of clinical skills and healthcare provider interpersonal skills were allocated to the category of "healthcare provider factors". Themes identified under the category of "patient-related factors" included: coping with uncertainty; the role of health literacy; the notion of hope, and the belief of stoicism.

Conclusions: The results of this study indicate that care provided by knowledgeable, caring, empathetic and receptive healthcare providers is critical to patients with axial SpA as they navigate the healthcare system from symptom onset to