FRI0717-HPR ASSESSING THE BURDEN OF TREATED AND UNTREATED CHRONIC LOW BACK PAIN IN EUROPE

M.J. Doane¹, D.H. Jaffe², E. Dragon³, L. Abraham⁴, L. Viktrup⁵, A.G. Bushmakina⁶, J.C. Cappelleri⁶. ¹Kantar Health, Horsham, United States; ²Kantar Health, Tel Aviv, Israel; ³Pfizer Ltd., Budapest, Hungary; ⁴Pfizer Ltd., Surrey, United Kingdom; ⁵Eli Lilly & Company, Indianapolis; ⁶Pfizer Ltd., Groton, United States

Background: Chronic low back pain (CLBP) is estimated to affect about 20% of adults¹ and is the greatest contributor to disability globally.² Current pharmacological treatment options may provide limited pain relief or may not be appropriate for long-term use in all patients because of adverse events.³ Despite the large burden of CLBP, few studies have evaluated the humanistic and economic impact of this condition in Europe, or whether this burden varies for those being treated with prescription (Rx) treatments compared with those who are not Rx treated.

Objectives: To assess the burden of CLBP in Europe and to determine whether burden differs by pain severity and treatment status.

Methods: A retrospective, cross-sectional study was conducted using data from the 2016 and 2017 National Health and Wellness Survey (NHWS) from five European Union countries (5EU): France, Germany, Italy, Spain, and United Kingdom. NHWS respondents with a self-reported CLBP diagnosis, current pain lasting ≥ 3 months, and who completed the pain module were identified. Neuropathic and phantom limb pains were excluded. CLBP respondents were categorized into 4 groups by severity of pain and treatment: moderate/severe Rx treated [M/S-Treated]; moderate/severe Rx untreated [M/S-Untreated]; mild Rx treated; and mild Rx untreated (reference). Outcomes of interest included health-related quality of life (HRQoL) (SF-12v2: mental and physical component summary [MCS, PCS]), health status (EQ-5D), productivity loss (Work Productivity and Activity Impairment [WPAI] questionnaire), and health care professional (HCP) visits in past 6 months. Multivariable analyses adjusted for baseline differences between groups (e.g., demographic and health characteristics).

Results: A total of 2,086 CLBP patients were identified from the NHWS. CLBP patients reported an average age of 56.4 (SD=13.2) years and most were female (61.2%). Two-thirds of CLBP patients had M/S pain (n=1,403). Stratification by pain and treatment groups resulted in the following groups: M/S-Treated=29.8%, M/S-Untreated=37.4%, Mild-Treated=19.5%, and Mild-Untreated=13.2%. Increased pain severity among both treated and untreated patients showed significantly worse HRQoL, health status, work and activity impairment, and greater HCP visits compared with the reference group (Mild-Untreated) (table 1).

Table 1 Adjusted mean levels per outcome according to disease severity and prescription treatment status^a

Outcome	M/S-Treated (A) (n=622)	M/S-Untreated (B) (n=781)	Mild-Treated (C) (n=407)	Mild-Untreated (Ref) (n=276)	Mild-Untreated (Reference) p-value		
					vs. Group A	vs. Group B	vs. Group C
Mental Component Score (MCS)	39.9	42.1	42.4	44.2	<0.001	0.003	0.024
Physical Component Score (PCS)	38.5	45.8	42.5	49.9	<0.001	<0.001	<0.001
Health Status (EQ-5D)	0.51	0.71	0.66	0.79	<0.001	<0.001	<0.001
Absenteeism ^b	5.6	1.6	3.3	0.9	0.053	0.138	0.075
Presenteeism ^c	31.6	21.4	26.4	12.3	<0.001	<0.001	<0.001
Overall Work Impairment ^d	33.7	22.2	28.5	12.6	<0.001	<0.001	<0.001
Activity Impairment	56.8	38.5	45.9	25.5	<0.001	<0.001	<0.001
General Practitioner Visits	3.1	1.9	2.6	1.6	<0.001	0.043	<0.001
Total Traditional Healthcare Provider Visits	7.1	4.6	6.0	3.7	<0.001	0.009	<0.001

^aAll models controlled for age, sex, marital status, education, income, employment status, smoking status, alcohol use, exercise, body mass index, Charlson Comorbidity index, and country using generalized linear regression models.

^bAmong employed respondents only: M/S-Treated n=192; M/S-Untreated n=321; Mild-Treated n=144; Mild-Untreated n=135.

^cAmong employed respondents only: M/S-Treated n=163; M/S-Untreated n=304; Mild-Treated n=131; Mild-Untreated n=132.

Conclusions: Results suggest the majority of European patients with CLBP in this study had moderate to severe pain irrespective of treatment. Whether treated or untreated, those with moderate to severe pain demonstrated a substantial burden related to HRQoL, health status, overall work and activity impairment, and HCP visits compared with those with mild pain.

REFERENCES:

- Meucci RD, et al. Rev Saude Publica 2015;49:1–10.
- GBD. Lancet 2017;390(10100):1211–59.
- Noble, M, et al. Cochrane Database Syst Rev. 2010;20(1):1–68.

Disclosure of Interest: M. Doane Grant/research support from: Pfizer Ltd., D. Jaffe Grant/research support from: Pfizer Ltd., E. Dragon Employee of: Pfizer Ltd., L. Abraham Employee of: Pfizer Ltd., L. Viktrup Employee of: Eli Lilly & Company, A. Bushmakina Employee of: Pfizer Ltd., J. Cappelleri Employee of: Pfizer Ltd.

DOI: 10.1136/annrheumdis-2018-eular.7497

FRI0718-HPR ARE PAIN, ACTIVITY LIMITATIONS AND QUALITY OF LIFE ASSOCIATED WITH OBJECTIVELY MEASURED PHYSICAL ACTIVITY IN PATIENTS WITH END-STAGE OSTEOARTHRITIS OF THE HIP OR KNEE?

C. Leichtenberg¹, F. van Tol¹, T. Krom¹, C. Tilbury¹, H. Horemans², H. Bussmann², R. Nelissen¹, M. Gademian^{1,3}, T. Vliet Vlieland¹ on behalf of the LOAS Study group. ¹Department of Orthopaedics, LUMC, Leiden; ²Department of Rehabilitation, ErasmusMC, Rotterdam; ³Department of Epidemiology, LUMC, Leiden, Netherlands

Background: Hip and knee osteoarthritis (OA) are a major cause of disability, yet relatively little is known on the relationship between OA and the total amount of physical activity.

Objectives: We investigated if OA-associated pain, functional disability and quality of life (QoL) are associated with objectively measured physical activity in patients with end-stage hip/ knee

Methods: Preoperative data from the Longitudinal Leiden Orthopaedics Outcomes of Osteoarthritis Study (LOAS) were used. All patients were scheduled for primary total hip or knee arthroplasty. Patients wore an accelerometer (Activ8 Remedy Distribution Ltd) which assessed levels of physical activity (LPA) in activity counts and %time spent on physical activity (%PA, i.e. time spent on walking, cycling or running) and sedentary behaviour (%SB, i.e. time spent in lying/sitting). Pain, functional disability and health related QoL were assessed with the Hip Disability and Knee Injury Osteoarthritis Outcome Scores (HOOS/KOOS) and Short-Form 12 (SF12). Multivariate linear regressions models adjusted for confounding were conducted. The physical activity outcomes were transformed to Z-scores.

Results: 9 hip OA and 48 knee OA patients were included. When awake, mean LPA was 18.787 $\pm$ 7.247 for hip OA patients and 21.193 $\pm$ 6.164 for knee OA patients. In hip OA patients %PA was on average 14 $\pm$ 6.4, while %SB was 66 $\pm$ 10.5. In knee OA patients these percentages were 15 $\pm$ 5.0 and 68 $\pm$ 8.7, respectively. In hip OA, the HOOS-QoL and SF12-PCS were positively associated with more LPA, (β 0.028; 95%CI:0.007 – 0.048, β 0.041; 95%CI:0.010 – 0.071) and better SF12-PCS scores also with PA (β 0.040, 95%CI:0.007 – 0.073). No other associations were found.

Conclusions: Whereas QoL was associated with physical activity in end-stage hip OA patients, no associations between pain or functional disability and physical activity were observed in either hip or knee OA.

Acknowledgements: This study was supported by the Dutch Arthritis Association [grant number LLP13].

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3478

FRI0719-HPR SELF-ASSESSMENT OF QUALITY OF LIFE OF PATIENTS WITH RHEUMATIC DISEASES AND OTHER CHRONIC DISEASES IN THE IEXPAC PROJECT

M.L. García-Vivar¹, L. Cea-Calvo², J. de Toro³, L. Pantoja⁴, C. Lerín⁵, S. García-Díaz⁶, M.J. Galindo⁷, I. Marín-Jiménez⁸, S. Fernández², Y. Mestre², D. Orozco-Beltrán⁹. ¹Rheumatology Department, Basurto University Hospital, Bilbao; ²Medical Affairs Department, Merck Sharp & Dohme, Madrid; ³Rheumatology Department, H. U. A Coruña, A Coruña; ⁴Rheumatology Department, H. El Bierzo, Ponferrada; ⁵Rheumatology Department, Manacor Hospital, Palma de Mallorca; ⁶Rheumatology Department, H. de Sant Joan Despi Moisès Broggi, Barcelona; ⁷Internal Medicine Department, H. Clínico de Valencia, Valencia; ⁸Gastroenterology Department, H. U. Gregorio Marañón, Madrid; ⁹Department of Medicine, Miguel Hernández University, Sant Joan, Alicante, Spain

Background: Improving quality of life is a goal in the treatment of patients with rheumatic diseases.

Objectives: In this work, we describe the self-assessment of quality of life made by patients with rheumatic diseases and with other chronic diseases through an anonymous survey in the context of a wider project on quality of care.

Methods: In the context of a quality of care project, focused in the perceptions of chronic patients with health care in Spain (assessed with the IEXPAC scale ("Instrument to Evaluate the EXperience of PATients with Chronic diseases", <http://www.iemac.es/iexpac/>), a survey was handed patients with 4 different profiles of chronic diseases needing care in at least two different levels (i.e. hospital clinic and primary care): A) Patients with rheumatic diseases (rheumatoid arthritis or spondyloarthritis) from hospital clinics, B) Inflammatory bowel disease (IBD) patients from hospital clinics, C) Patients with human immunodeficiency virus (HIV) infection from HIV units and D) Patients with diabetes mellitus (DM) plus