most of the patients are moderately disabled, and about 10% of the patients suffered from severe disablement which brings the necessity of a caregiver to become the patient's companion due to its chronic disease.² The caregiving role can have an impact in the psychological and physical spheres of the caregiver's life.³

Objectives: The aim of this study was to explore demographical characteristics and caregiver burden through the Zarit Scale.

Methods: We conducted a cross sectional study in a meeting where caregivers in a rheumatoid arthritis specialised setting. We collected sociodemographic information, and applied the Zarit caregiver burden interview (ZBI) adapted to Spanish. The ZBI includes 22 questions which has 5 responses from 0 (never) to 4 (nearly always), where scores lower than 47 indicated little to no burden, 47 to 55 low burden and >55 intense burden. We calculated means, and standard deviations for continuous variables and categorical variables were presented as rates. We categorised age of caregivers and compared to it to ZBI score, we used X² to perform bivariate analysis.

Results: 115 caregivers were included in the study, 63% were female and 37% were male. Mean age was 49 years±18 and 35% were single. Regarding educational level 44% had college degree, most of caregivers 30% had a full time job, and also 45% had other family members as their responsibility. See table 1. Zarit mean score was 44±14, additionally 71% had a score lower than 47. See table 1. Regarding age groups 35% of patients were older than 60 years and 15% of them were considered with intense burden disease, there was no statistical association between age and disease burden, see table 2.

Abstract AB1451HPR – Table 1. Zarit score

Zarit Score		
<47	82	71
47–55	14	12
>55	19	17

Abstract AB1451HPR – Table 2. Zarit scale classification according to age groups

Variable	15–39 years n=36	40–60 years n=38	Older than 60 years n=41	P value
No burden	20	28	30	0.108
Low burden	8	5	5	0.375
Intense burden	8	5	6	0.436

Conclusions: Although a high proportion of caregivers reported to have not burden, it is important to develop strategies and activities in order to attend the necessities of the caregivers in order to reduce the overload of responsibilities among them. Also further research is needed in order to identify the risk factors or protector factors that can prevent disease burden in caregivers.

REFERENCES:

- [1] Prothero L, Georgopoulou S, de Souza S. Patient involvement in the development of a handbook for moderate rheumatoid arthritis. 2016.
- [2] Jacobi CE, van den Berg B, Boshuizen HC, Rupp I, Dinant HJ, van den Bos GA. Dimension-specific burden of caregiving among partners of rheumatoid arthritis patients. *Rheumatology (Oxford, England)*. 2003;42(10):1226–33.
- [3] Win KK, Chong MS, Ali N, Chan M, Lim WS. Burden among Family Caregivers of Dementia in the Oldest-Old: An Exploratory Study. *Frontiers in medicine*. 2017;4:205.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6957

AB1452-HPR WHICH ONE HAS A GREATER EFFECT ON FUNCTION AND THE PSYCHOSOCIAL STATUS IN JIA?: DISEASE TYPE OR THE PRESENCE OF PAIN

G. Arın¹, H.E. Sönmez², N.B. Karaca¹, E. Sağ², A. Özçadırcı¹, S. Demir², F. B. Oflaz¹, Y. Bilginer², D. Aydın Hakkı³, R. Alpar², E. Ünal¹, S. Özen². ¹Department of Physiotherapy and Rehabilitation, Hacettepe University Faculty of Health Sciences; ²Department of Pediatric Rheumatology; ³Department of Biostatistics, Hacettepe University Faculty of Medicine, Ankara, Turkey

Background: Juvenile idiopathic arthritis (JIA) is a chronic inflammatory childhood disease with symptoms such as joint inflammation, pain and

loss of quality of life.¹ Types of disease and the presence of pain can impact the child psychosocially, as well as affecting functional activity.²

Objectives: The aim of this study is to examine the results of functional and psychosocial status according to the disease type and the presence of pain symptoms in children with JIA.

Methods: The study included 71 children diagnosed with JIA who applied to the Hacettepe University İhsan Doğramacı Children's Hospital Rheumatology Department. Following the collection of demographic information, functional status was assessed with the Child Health Assessment Questionnaire (CHAQ) and psychosocial and functional status was assessed with the scale developed in Hacettepe University Faculty of Health Sciences Department of Physiotherapy and Rehabilitation for functional and psychosocial status of children with rheumatism by Edibe Ünal.³ Children were divided into groups according to disease type as oligoarthritis or polyarthritis and the presence or absence of pain.

Results: Table 1 shows the mean age and numbers of children. There was no difference between the groups according to disease type (p>0,05). On the other hand, comparing scores for the CHAQ total, CHAQ general VAS assessment, functional and psychosocial status according to the presence or absence of pain revealed significant differences (p<0,05).

Abstract AB1452HPR – Table 1. Assessment values and comparison statistics

	Disease Type			Pain		
	Oligoarthritis (n=51)	Polyarthritis (n=20)	p	Present (n=21)	Absent (n=50)	p
Age (years)	10,88±3,81	13,50±3,92	0016	11,71 ±3,77	11,58 ±4,12	0885
CHAQ Total	0,28±0,29	0,46±0,41	0127	0,51±0,4 ±0,27	0,26 ±0,27	0012
CHAQ (General VAS)	2,49±2,43	3,93±3	0068	4,46 ±2,86	2,24±2,3	0002
Function (range 0–30)	4,72±4,85	5,05±6,32	0766	7,85±6,6 ±4,02	3,54 ±4,02	0004
Psychosocial (range 0–30)	23±5,67	14,5±5,82	0363	16±5,74 ±5,4	12,34 ±5,4	0012

Function; Psychosocial; Functional and Psychosocial subscales of Ünal's scale.³

Conclusions: We conclude that pain has a greater effect on functional, psychosocial and overall disease assessment in children with JIA when compared to the disease type. Thus, it must be taken into consideration that child's ability to cope with pain should be improved.

REFERENCES:

- [1] Angelo Ravelli, Alberto Martini. Juvenile idiopathic arthritis. *The Lancet* 2007, 369(9563); 767–778.
- [2] Laura E Schanberga, John C Lefebvre, et al. Pain coping and the pain experience in children with juvenile chronic arthritis. *Pain* 1997, 73(2); 181–189.
- [3] Kısacık Pınar, Ünal Edibe, et al. Juvenil İdiyopatik Artritli Hastalarda Çok Yönlü Bir Değerlendirme Sistemi Olusturulması Delphi Çalışması. 2016, *Annals Of Paediatric Rheumatology*

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5860

AB1453-HPR ARE DMARD THERAPIES EFFECTIVE ON QUALITY OF LIFE, ANXIETY AND DEPRESSION LEVELS IN PATIENTS WITH RHEUMATOID ARTHRITIS?

G.İ. Kınıklı¹, F. Akar², S. Demirtas², M.A. Duman², S. Ergin², M.E. Gebes², Y. Guneser², G. Kınıklı². ¹Department of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Hacettepe University; ²Department of Rheumatology, Faculty of Medicine, Ankara University, Ankara, Turkey

Background: Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease with persistent joint inflammation manifesting as joint pain, stiffness, and swelling. Treatment of rheumatoid arthritis (RA) should be aimed at achieving the lowest possible disease activity. Conventional DMARDs (eg, methotrexate [MTX], hydroxychloroquine, and sulfasalazine) are still widely used in newly diagnosed RA patients. On the other hand, the rate of biologic therapy use in clinical practice is rising as more agents become available in spite of efficacy of these treatments is broadly comparable. Depression is a common under-recognised co-morbidity in patients with RA accompanying with substantial disability, reduced quality of life.