OP0319

LOW AND MODERATE PHYSICAL ACTIVITY REDUCES LOCALISED IL-1B IN AN ACUTE MOUSE MODEL OF GOUT BY DOWN-REGULATING TLR2 EXPRESSION ON CIRCULATING NEUTROPHILS

K. Jablonski¹*, B. Sandoval¹, P. Harb¹, A. Kalyanasundaram², J.M. Hampton¹, W. N. Jarjour¹, N. Schlesinger³, N.A. Young¹. ¹Rheumatology and Immunology; ²Physiology and Cell Biology, The Ohio State University Wexner Medical Center, Columbus; ³Rheumatology, Rutgers Robert Wood Johnson Medical School, New Brunswick, USA

Background: While physical activity was originally believed to exacerbate inflammation in rheumatic disease, recent studies have shown significant reductions in inflammation with regular exercise. It has been previously shown that down-regulation of toll-like receptor (TLR)2 and TLR4 expression correlates with increased physical activity in humans. Furthermore, both TLR2 and TLR4 knockout mice are resistant to monosodium urate (MSU) crystal-induced gout. Additionally, mesenchymal stem cells (MSCs) can be immunosuppressive by secreting IL-1 receptor antagonist (IL-1RA) and have also been shown to be up-regulated with exercise.

Objectives: The aim of this study was to investigate the mechanism by which exercise suppresses gouty inflammation and to define the potential roles of TLR2, TLR4, and MSCs in the process.

Methods: NFκB reporter mice [BALB/C-Tg(NFκB-RE-luc)-Xen] were exercised daily by treadmill walking (45 min/day for 2 weeks) at low intensity (35% VO₂max; 8 m/min), moderate intensity (55% VO₂max; 11 m/min), and high intensity (75% VO₂max; 15 m/min). Mice were then injected with MSU crystals (0.5 mg) into the tibio-tarsal joint (ankle). Localised NFκB activity was measured 16 hours later in the injected ankle by bioluminescent imaging. Tissue was collected and processed for immunohistochemical (IHC) analysis and whole blood was collected for both flow cytometry and serum analysis.

Results: Mice in the low/moderate intensity exercise groups had decreased inflammation, F4/80⁺ macrophages, and MPO⁺ neutrophils at the site of MSU injection compared to high intensity and non-exercised controls. Similarly, bioluminescent imaging of NFκB activity was significantly reduced locally in both low/moderate intensity groups compared to high-intensity and non-exercised controls. Surface expression of TLR4 on peripheral monocytes or neutrophils showed little difference by flow cytometry, while TLR2 expression on peripheral neutrophils was significantly reduced. In concordance, localised IL-1β expression via IHC was reduced in low/moderate intensity exercise conditions. IL-1RA expression correlated with IL-1β induction locally by IHC and was elevated in serum. Also, bone marrow-derived MSCs were significantly reduced in low/moderate intensity exercise compared to high-intensity or non-exercised controls.

Conclusions: These data show that while low/moderate intensity exercise regimens can reduce the localised MSU crystal-induced inflammation, high intensity training negates this response. Moreover, the exercise-mediated suppression of NFκB activity and IL-1β expression locally can be partially explained by a reduction in peripheral neutrophil recruitment via downregulation of TLR2 expression in the peripheral blood. Although not clearly defined mechanistically in this study, our results also suggest that MSCs may contribute to this immunosuppressive response and are mobilised out of the bone marrow with low/moderate intensity exercise.

REFERENCES:

- [1] Khanna D, et al. Arthritis Care Res 2012;64:1447.
- [2] Agudelo CA, et al. Arthritis Rheum 1972;15:609.
- [3] Liu-Bryan R, et al. Arthritis Rheum 2005;52:2936.

Acknowledgements: Support provided by Ironwood Pharmaceuticals, Cambridge MA 02142

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4764

OP0320

S100A9 HAMPERS OSTEOCLAST DIFFERENTIATION FROM CIRCULATING PRECURSORS BY REDUCING THE EXPRESSION OF RANK

M.H. Van Den Bosch¹, I. Di Ceglie¹, T. Vogt², J. Roth², C.S. Goodyear³, P.M. van der Kraan¹, A.B. Blom¹, P.L. van Lent¹. ¹Experimental Rheumatology, Radboud university medical center, Nijmegen, Netherlands; ²Institut für Immunologie, Münster, Germany; ³Institute of Infection, Immunity and Inflammation, Glasgow, UK

Background: High levels of the damage-associated molecular patterns (DAMPs) S100A8 and S100A9 are produced in the synovium during both experimental and human rheumatoid arthritis (RA). These alarmins have been implicated in inflammation-induced bone resorption. We and others have previously shown that stimulation of mature osteoclasts with S100A8/A9 results in increased numbers and bone resorptive activity. In agreement, reduced bone destruction was observed after induction of experimental RA models in S100a9^{-/-} mice. However, the effects of S100A8/A9 on monocyte-to-osteoclast differentiation remain elusive.

Objectives: Here, we investigated the effects of S100A9 on CD14⁺ monocytes and their potential to differentiate into osteoclasts.

Methods: CD14⁺ monocytes were isolated from buffy coats of healthy donors using density gradient centrifugation and magnetic cell sorting. Cells were differentiated into osteoclasts with macrophage colony-stimulating factor (M-CSF) and Receptor activator of nuclear factor kappa-B (RANK) ligand (RANKL) in the presence or absence of S100A9. mRNA expression was determined by RT-qPCR and protein expression was determined using Luminex analysis. Moreover, osteoclast differentiation was assessed using Tartrate-resistant acid phosphatase (TRAP) staining and the resorptive capacity was determined using mineral-coated plates. RANK protein expression was assessed using FACS.

Results: We observed that S100A9 stimulation of monocytes resulted in a strong induction of various pro-inflammatory factors, such as interleukin (IL)1β, IL6, IL8, and tumour necrosis factor (TNF)α after 24 hour, both on the mRNA and protein level. Interestingly, we observed a strong decrease in the number of multinucleated osteoclasts as determined by TRAP staining, at day 6 and 8 after start of the cultures. In agreement with this, the cells showed a strongly reduced resorptive capacity after 10 days of culture. We demonstrated that already a 24 hour stimulation with S100A9 strongly reduced the osteoclastogenic potential of the CD14⁺ monocytes. Finally, to determine the mechanism of how this short S100A9 stimulation might reduce the osteoclast development, we determined the protein expression of the RANK receptor, which is crucial for osteoclast differentiation. We observed that S100A9 stimulation hampered the M-CSF-induced upregulation of RANK after 24 hours, suggesting that this underlies the hampered osteoclast differentiation. Interestingly, this S100A9-induced decreased RANK expression could be reversed by addition of the TNFα-inhibitor etanercept, but not by addition of IL1 receptor antagonist.

Conclusions: Whereas S100A8/A9 have been previously shown to stimulate the numbers and resorptive capacity of mature osteoclasts, we here show that stimulation of monocytes with S100A9 strongly inhibits their osteoclastogenic potential, possibly via TNFα-induced reduction of RANK expression. This suggests that S100A8/A9 does not solely stimulate osteoclast formation and function but rather that the timing of exposure to S100A8/A9 is an important determinant for monocyte-to-osteoclast differentiation.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3689

FRIDAY, 15 JUNE 2018

Pathophysiology and biomarkers in PsA: what impact?

OP0321

PRECISION MEDICINE USING DIFFERENT BIOLOGICAL DMARDS BASED ON CHARACTERISTIC PHENOTYPES OF PERIPHERAL T HELPER CELLS IN PSORIATIC ARTHRITIS

I. Miyagawa, S. Nakayama, K. Nakano, S. Kubo, S. Iwata, Y. Miyazaki, M. Yoshikawa, H. Yoshinari, Y. Tanaka. The First Department of Internal Medicine, University of Occupational and Environmental Health, Japan, Kitakyusyu, Japan

Background: Biological DMARDs targeting TNF-α, IL-17, and IL-12/23 (p40) are available. The high efficacy of these drugs has been proven in numerous clinical trials. However, there are some cases in which a change from one bDMARDs to another one is necessary because of the refractory nature of the disease, and there is no established method to select the optimal bDMARDs according to the individual case, despite the fact that various drugs are available.

Objectives: We sought to investigate the selection of specific biological DMARDs based on characteristic lymphocyte phenotypes for treating PsA.

Methods: we performed this study to evaluate the efficacy of biologics therapy in 64 patients with PsA after 6 months of therapy, and to compare the results of