- [3] Wallace ZS, Mattoo H, Mahajan VS, et al. Predictors of disease relapse in IgG4-related disease following rituximab. *Rheumatology (Oxford)*. 2016 Jun;55(6):1000–8.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3895

THU0039 INSULIN-LIKE GROWTH FACTOR 1 RECEPTOR REGULATES THE PHENOTYPE AND FUNCTION OF CD21+ B CELLS

M. Erlandsson, C. Wasen, G. Gravina, M.I. Bokarewa. *Rheumatology and Inflammation Research, GOTHENBURG UNIVERSITY, Gothenburg, Sweden*

Background: Ligand to the inducible T cell costimulator (ICOSL) on B cells is essential for the ICOS-dependent follicular recruitment of activated T cells. In patients with rheumatoid arthritis (RA) the IGF1-IGF1R axis is altered. Inhibition of IGF1R alleviated arthritis by reducing IL-6-dependent formation of Th17 cells.¹ Here we study the role of IGF1R on CD21+ cells in experimental arthritis.

Methods: Female Balb/c mice were immunised with methylated BSA or with CII. Consequences of the IGF-1R inhibition for arthritis were studied in mBSA and CII-immunised mice treated with NT157 compound promoting degradation of insulin receptor substrates or using shRNA producing construct (shIGF1R). At termination three sub-populations of CD21+ cells were analysed: follicular dendritic cells (FDC, CD21+CD19-CXCR5-); marginal zone B cells (MZBc, CD21+CD19+CXCR5-); follicular B cells (FBc, CD21+CD19+CXCR5+). Supernatants of LPS-stimulated splenocytes were analysed for production of cytokines, chemokines using Cytokine Array. Serum levels of antigen specific and autoantibodies were measured in an ELISA.

Results: In spleen of mBSA-immunised mice, ICOSL expression on CD21+ cells correlated to IGF1R ($r=0.70$, $p=0.007$). Inhibition of IGF1R induced a 20% reduction in ICOSL expression in all CD21+ subsets ($p=0.007$) followed by an increase in the number of MZBc ($p=0.003$), while FDC and FBc were unchanged. Inhibition of IGF1R had no effect on the expression of ICOSL on CD4 T cells or the subset of CXCR5+ follicular T cells. Reduction of the ICOSL+CD21+B cells was associated with lower production of IL-13. Inhibition of IGF1R signalling by NT157 and by shRNA, reduced production of CXCL13 and CXCL12, the chemokines essential for B cell migration towards follicles. In contrast, the production of chemokines CCL5 and CXCL12 preventing intra-follicular migration was increased, which explains the increase of MZBc. Additionally, the insufficient ICOSL signalling significantly reduced the production of IL-7 and IL-4, regulating class switching of B cells in germinal centres and differentiation of B cells into plasma cells. The described disbalance in the cytokines aiding B cell development led to the reduced production anti-inflammatory IL-10 and of mBSA-specific IgM ($p=0.005$) and increased production of autoreactive RF-IgM levels ($p=0.001$).

Conclusions: The study shows that IGF1R controls B cell development through the expression of ICOSL on CD21+ cells. Insufficient ICOSL signalling disturbs a balance between antigen-specific response and autoantibody production in experimental arthritis.

REFERENCE:

- [1] Erlandsson MC, Toyra Silfversward S, Nadali M, Turkkila M, Svensson MND, Jonsson IM, et al. IGF-1R signalling contributes to IL-6 production and T cell dependent inflammation in rheumatoid arthritis. *Biochim Biophys Acta*. 2017;1863:2158–70.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5300

THU0040 ENHANCEMENT OF EARLY HUMAN B CELL DEVELOPMENT BY JAK INHIBITION

J. Thiel¹, N. Venhoff¹, I. Janowska¹, A. Troilo¹, N. Frede¹, M. Erlacher², M. Kunze³, R. Lorenzetti¹, J. Staniek¹, B. Bannert⁴, D. Kyburz⁴, C. Glaser¹, M.-T. Schleyer¹, R. E. Voll¹, M. Rizzi¹. ¹Rheumatology and Clinical Immunology; ²Clinic for Pediatrics and Juvenile Medicine; ³Gynecology Department, University Medical Center Freiburg, Freiburg, Germany; ⁴Department of Rheumatology, University of Basel, Basel, Switzerland

Background: Rheumatoid arthritis is an immune-mediated disease, in which immune cell activation leads to destructive inflammation of the joints. In this context the treatment with JAK inhibitor tofacitinib has been proven to be effective. In mice treatment with tofacitinib resulted in reduced specific antibody responses, failure to generate germinal centres, and partial block of B cell development in the bone marrow. Conversely *in vivo* treatment of psoriasis arthritis patients as well as of rheumatoid arthritis patients with tofacitinib results in an increase of relative

and absolute numbers of B cells in the first 4–8 weeks from beginning of treatment. It is known that mouse early B cell development is strongly dependent on IL-7 signalling, while this is not essential in humans. Nevertheless other cytokines are important in determining the fate and development of B lymphocytes. Hence, the outcome of JAK inhibition in early B cell development remains to be studied

Objectives: To assess the impact of JAK inhibition on early B cell development *in vitro*

Methods: We used a *in vitro* model in which CD34+ cells isolated from cord blood are cultivated subsequently in SCF, Flt3-L and IL-6, then SCF, Flt3-L and IL-7 and finally in cytokine-free medium.¹ The culture reproduces all stages of development from common lymphocytes progenitors to immature B cells.

Results: With the addition of tofacitinib to the *in vitro* culture we observed an increase in the absolute numbers of lymphoid precursors developing *in vitro* especially at week 5 and 6 of culture. Specifically, JAK inhibition led to an increase of pre-B and immature B cells by week 5 and 6. These data are in contrast with the early B cell development block observed in the tofacitinib treated mouse, but are in line with the rapid increase of B cells in peripheral blood after 4–8 weeks of tofacitinib treatment in patients. Analysis of induction of fate determining genes (EBF, E2A, PAX-5) showed an earlier and stronger induction of fate determining genes.

Conclusions: Our data indicate that JAK inhibition may promote early B cell development by enhancing the commitment of lymphoid precursors to the B cell compartment, contributing to a temporary increase in relative and absolute numbers of B cell in peripheral blood of treated patients. These data contribute to our understanding of human B cell development, prompt us to further analyse the quality of B cell output from the bone marrow in JAK inhibited patients, and may provide cues to understand the outcome of JAK inhibition treatment in rheumatic diseases.

REFERENCE:

- [1] Kraus H, et al. A feeder-free differentiation system identifies autonomously proliferating B cell precursors in human bone marrow. *J Immunol*. 2014 Feb 1;192(3):1044–54.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5190

THU0041 THE EFFECTS OF ABATACEPT ON HUMAN B CELL ACTIVITIES

M.-H. Chen¹, C.-T. Ssu², C.-Y. Tsai¹, C.-M. Leu³. ¹Division of Allergy, Immunology and Rheumatology, Department of Medicine, Taipei Veterans General Hospital; ²Institute of Microbiology and Immunology; ³Institute of Microbiology and Immunology, and Infection and Immunity Center, National Yang-Ming University, Taipei City, Taiwan, Province of China

Background: Abatacept is a cytotoxic T lymphocyte antigen-4 (CTLA-4) fusion protein approved for rheumatoid arthritis (RA) treatment worldwide. Abatacept mimics the natural CTLA-4 and competes with CD28 for binding the CD80/CD86 on antigen presenting cells to prevent T cell activation. However, the impacts of CTLA-4-Ig through its interaction with CD80/CD86 on B cells are not fully understood.

Objectives: The aim of this study was to test whether CTLA-4 regulates human B cell functions

Methods: We assayed the effect of abatacept on human B cells in both *in vitro* and *in vivo* conditions. Blood was taken from 20 patients with RA before and after abatacept treatment and the expression of surface proteins on B cells was detected using immunofluorescent staining. Serum level of rheumatoid factor (RF) and anti-cyclic citrullinated peptide antibody (ACPA) was measured by ELISA. Purified human B cells from healthy donors were stimulated in the presence of abatacept and cell proliferation, cytokine production, plasma cell differentiation, and antibody production were measured.

Results: In RA patients, abatacept treatment transiently reduced the level of CD80/CD86 on peripheral blood memory B cells and increased the naïve to memory B cell ratio. Also, abatacept reduced IgM-RF level in 10 out of 12 patients, which correlates with patients' disease activity, but it had no significant effects on serum levels of anti-citrullinated protein antibody or anti-tetanus toxoid antibodies during the 6 months of abatacept treatment. In the *in vitro* assays, we observed that the CD80 and CD86 induced by T-independent (TI) but not T-dependent (TD) stimulation was significantly downregulated by abatacept at both the mRNA level and protein level. Some TI-induced cytokine production by B cells from healthy donors was also reduced by abatacept. Neither TI nor TD-stimulated B cell proliferation was reduced by abatacept in ³H-thymidine incorporation assay. Abatacept had no significant effect on CD38^{high} CD27^{high} plasma cell differentiation. Finally, abatacept inhibited Daudi-B cell induced allogeneic T cell proliferation, indicating a significant blockade of T-B interaction by abatacept.

Conclusions: In RA patients, abatacept may decrease RF level by interfering the interaction of CD28 with CD80/CD86, therefore preventing B cells from T cells'

help for differentiation into plasma cells. Our results also indicate that abatacept binding CD80/CD86 may provoke a negative reverse signal to B cells to further regulate B cells activation.

Disclosure of Interest: M.-H. Chen Grant/research support from: BMS, C.-T. Ssu: None declared, C.-Y. Tsai: None declared, C.-M. Leu Grant/research support from: BMS

DOI: 10.1136/annrheumdis-2018-eular.1474

THU0042 CANCER IMMUNOTHERAPY ARRAY: A NOVEL SCREENING TOOL FOR IMMUNE SYSTEM PROFILING IN CANCER IMMUNOTHERAPY BRIDGING AUTOIMMUNITY AND CANCER

P. Schulz-Knappe, P. Budde, H.-D. Zucht, S. Konings, L. Steeg, E. Friedrich, C. Gutjahr, R. Steil, S. Bhandari, M. Tuschen. *Protagen AG, DORTMUND, Germany*

Background: Recent FDA-approved checkpoint inhibitors targeting the cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4) and programmed death 1 (PD-1)/PD-L1 pathway represent milestones in the field of cancer immunotherapy. In general, cancer immunotherapy works only in a subset of patients, but some patients experience prolonged responses. Cancer immunotherapy can cause severe immune-related adverse events (irAE) in patients, who are increasingly seen by rheumatologists. We propose that that autoantibody profiling will reveal novel B-cell associated mechanisms of therapy response and side effects. This may yield minimally-invasive biomarkers to identify patients at risk to develop irAE and monitor cancer patients over the course of their life under immunotherapy

Objectives: We have developed a novel Cancer Immunotherapy Array, which includes a combination of antigens important in autoimmune diseases, anti-tumour immunity, and oncogenes and tested the array in patient sera from a diverse set of cancer immunotherapy trials.

Methods: The Cancer Immunotherapy Array consists of a bead-based multiplex array using minimal patient serum samples incubated with antigen-coated, color-coded Luminex beads. Run in microtiter plate format, the Array permits quantification of the autoantibody reactivity in thousands of serum samples towards approximately 900 human protein antigens in each sample. Magnetic beads are employed to enable automated pipetting and washing steps.¹ We selected human protein antigens from groups A) tumor-associated antigens (TAA), B) autoimmune disease antigens, C) cytokines, and D) cancer signalling pathway proteins

Results: In total, over 2000 serum samples from diverse cancer indications plus hundreds of samples from autoimmune diseases such as RA, SLE, Sjogren's disease and healthy controls were screened with the Cancer Immunotherapy Array. As key findings we report autoantibody panels which can differentiate patients with irAEs and those without irAEs. Also, but less prominent, individual autoantibodies are associated with overall survival. Autoantibodies that target antigens involved in cancer signalling pathways are associated with irAEs. Also, patients with increased levels of a distinct autoantibody against an inflammatory cytokine do not develop irAEs across multiple tumours.

Conclusions: The Cancer Immunotherapy Array is a high throughput array suitable for the analysis of thousands of cancer patient serum samples. Its first application presents novel autoantibody signatures for therapy-related toxicities (irAEs) as well as response. These signatures have the potential to serve as useful tools that will broaden our understanding of the mechanisms of therapy response and irAE occurrence.

REFERENCE:

- [1] Budde P, et al. *Lupus*. 2016;25:812–22.

Disclosure of Interest: P. Schulz-Knappe Shareholder of: Protagen AG, P. Budde Employee of: Protagen AG, H.-D. Zucht Employee of: Protagen AG, S. Konings Employee of: Protagen AG, L. Steeg Employee of: Protagen AG, E. Friedrich Employee of: Protagen AG, C. Gutjahr Employee of: Protagen AG, R. Steil Employee of: Protagen AG, S. Bhandari Employee of: Protagen AG, M. Tuschen Employee of: Protagen AG

DOI: 10.1136/annrheumdis-2018-eular.6712

THU0043 AUTOANTIBODY PROFILING IN PROSTVAC AND IPIILIMUMAB TREATED PROSTATE CANCER PATIENTS REVEALS POTENTIAL BIOMARKERS OF IMMUNE-RELATED ADVERSE EVENTS

P. Budde¹, J.L. Marte², H.-D. Zucht¹, S. Bhandari¹, M. Tuschen¹, P. Schulz-Knappe¹, R.A. Madan², J.L. Gulley², J. Schlom². ¹PROTAGEN AG, Dortmund, Germany; ²National Institutes of Health/National Cancer Institute, Bethesda, USA

Background: Autoantibodies (AAB) targeting self-antigens can be found in two clinically and immunologically opposing diseases, autoimmune diseases and cancer. While in autoimmune diseases, the immune system is hyperactivated against self-antigens, many tumours suppress the anti-tumour immune response. The therapeutic cancer vaccines PSA-Tricom (Prostvac) is designed to generate an antigen-specific tumour response in metastatic castration-resistant prostate cancer (mCRPC), which is in phase 3 testing. To further augment the immune response, combination therapies of Prostvac with ipilimumab are currently tested in clinical studies. Ipilimumab is an antibody that blocks the immune checkpoint molecule cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4). However, treatment with ipilimumab is associated with immune-related adverse events (irAEs).¹

Objectives: Since there are no biomarkers for predicting irAEs, we aimed to investigate AAB profiles as biomarkers associated with irAE in mCRPC patients treated with prostvac and ipilimumab combination therapy.

Methods: Serum samples from 24 mCRPC patients treated with prostvac and ipilimumab therapy were tested for the presence of serum autoantibodies against 842 preselected antigens. Candidate antigens comprise immune-related and cancer signalling pathway proteins, autoimmune disease antigens, and tumor-associated antigens (TAA). Samples were collected prior to treatment (T0 samples), at 3 and 6 month. IrAEs included rash, elevated aminotransferases, neutropenia, diarrhoea, colitis and endocrine irAEs. Overall survival was also captured and correlated with AABs. Autoantibody levels were measured by Luminex FlexMap3D bead based multiplex protein arrays² and data were analysed by significance analysis of microarrays (SAM), Partial least squares regression (PLS) and Pearson's correlation.

Results: In total, 87 AABs were found that were significantly different in patients with irAEs and those without irAEs (SAM |d|>2.5; Pearson's correlation |3|>0.35). PLS analysis revealed that AABs associated with irAEs were also associated with overall survival. Gene ontology analysis of pathways, molecular function and cellular localization revealed that AABs predicting irAEs target cancer, cell cycle, cell adhesion and apoptotic pathways. We also found elevated levels of AABs in patients who do not experience irAEs. Interestingly, these 40 AABs target proteins that are involved in inflammatory, adaptive and cellular immune response pathways or are autoimmune disease antigens.

Conclusions: AABs that target antigens involved in cancer signalling pathways are associated with irAEs following prostvac plus checkpoint inhibitor combination therapy. In contrast, AABs targeting immune response pathways were found in patients who do not develop irAEs and may counteract the action of inflammatory molecules. Similarly, anti-cytokine AABs have been found in autoimmune diseases, where they appear to counteract the pathological effects of cytokines.³ Further studies in larger sample sets are needed to confirm these findings.

REFERENCES:

- [1] Madan RA, et al. *Lancet Oncol* 2012;13(5):501–8.
[2] Budde P, et al. *Lupus*. 2016;25:812–22.
[3] Cappellano G, et al. *Am J Clin Exp Immunol*. 2012;1(2):136–46.

Disclosure of Interest: P. Budde Employee of: Protagen, J. Marte: None declared, H.-D. Zucht Employee of: Protagen, S. Bhandari Employee of: Protagen, M. Tuschen Employee of: Protagen, P. Schulz-Knappe Shareholder of: Protagen, R. Madan: None declared, J. Gulley: None declared, J. Schlom: None declared

DOI: 10.1136/annrheumdis-2018-eular.2383