THU0165

OBESITY AS ONE OF THE COMMODITIES WAS THE ROBUSTEST PREDICTION FACTORS FOR POST THERAPEUTIC CLINICAL REMISSION OF RHEUMATOID ARTHRITIS WITH SHORT DISEASE DURATION – RESULTS FROM KANSAI CONSORTIUM FOR WELL-BEING OF RHEUMATIC DISEASE PATIENTS

K. Murakami¹, M. Hashimoto², K. Murata², W. Yamamoto^{2,3}, R. Hara⁴, M. Katayama⁵, A. Ohnishi⁶, K. Akashi⁶, S. Yoshida⁷, K. Nagai⁷, Y. Son⁸, H. Amuro⁹, T. Hirano⁹, K. Ebina¹⁰, K. Nishitani², M. Tanaka², H. Ito¹¹, K. Ohmura¹, T. Mimori¹. ¹Department of Rheumatology and Clinical Immunology; ²Department of Advanced Medicine for Rheumatic Diseases, Kyoto University Graduate School of Medicine, Kyoto; ³Department of Health Information Management, Kurashiki Sweet Hospital, Kurashiki; ⁴The Center for Rheumatic Diseases, Nara Medical University, Nara; ⁵Department of Rheumatology, Osaka Red Cross Hospital, Osaka; ⁶Department of Rheumatology and Clinical Immunology, Kobe University Graduate School of Medicine, Kobe; ⁷Department of Internal Medicine (IV), Osaka Medical College, Takatsuki; ⁸First Department of Internal Medicine, Kansai Medical University, Hirakata; ⁹Department of Respiratory Medicine, Allergy and Rheumatic Disease; ¹⁰Department of Orthopaedic Surgery, Osaka University Graduate School of Medicine, Osaka; ¹¹Department of Orthopaedic Surgery, Kyoto University Graduate School of Medicine, Kyoto, Japan

Background: A part of rheumatoid arthritis (RA) patients are resistant to clinical remission (CR) irrespective of therapies. In addition to known risk factors, systemic organ complications are assumed to interfere with CR.

Objectives: To extract the predictive comorbidities of clinical response of disease activities.

Methods: In Kansai consortium for well-being of rheumatic disease patients (ANSWER) cohort, which was the real world cohort of clinical database of rheumatic diseases, RA patients within 3 years of disease duration were included and followed. Using logistic regression analysis, background factors at the initial visit were extracted in order to predict CR after 1 year (1 year-non CR).

Results: 651 patients met in the inclusion criteria were under the analysis. Of those, 245 (37.6%) cases were resulted in 1 year-non CR. The average scores of DAS28-CRP at first visit and one year later was 3.51 and 2.02, respectively. Logistic regression analysis revealed that DAS28-CRP at first visit (OR 1.42/unit, 95% CI 1.24–1.63), concomitant use of methotrexate (MTX) or biologic disease modifying rheumatic drugs (bDMARDs) (OR 2.04, 95% CI 1.41–2.96) and body mass index (BMI) (OR 1.07/unit, 95% CI 1.02–1.12) were significant predictive factors of 1 year-non CR, but not in the case with gender, age, disease duration, hypertension, diabetes, dyslipidemia, lung diseases, heart diseases, digestive tissue diseases, history of malignancy nor concomitant autoimmune diseases. Using propensity score matching (1:1) stratified by gender, age, disease duration, concomitant use of MTX or bDMARDs and DAS28-CRP at baseline, the 49 pairs were matched between obese (defined as body mass index (BMI) over than 28) and non-obese patients. In these cohort, mean score of DAS28-CRP after 1 year was significantly higher in obese patients than in non-obese patients (2.63 and 2.18, respectively, see the figure 1).

Conclusions: In these early arthritis cohort, obese patients tend to remain higher disease activities even considering with gender, age and therapeutic management, although validation studies should be added to confirm these findings.

Acknowledgements: None.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1680

THU0166

THE DIASTOLIC DYSFUNCTION IN PATIENTS WITH RHEUMATOID ARTHRITIS

L. Feiskhanova. Department of hospital therapy, Kazan state medical University, Kazan, Russian Federation

Background: Rheumatoid arthritis is autoimmune rheumatic disease characterised by injury not only the joints but also other organs, including the heart. It is known that the presence of rheumatoid arthritis increases the risk of fatal cardiovascular complications by 1.5 times compared to the general population.

Objectives: to identify features of development of diastolic dysfunction in patients with rheumatoid arthritis.

Methods: we examined 180 patients with rheumatoid arthritis. The activity of disease was defined according to the scale of the DAS-28. By echocardiography in 101 patients (group 1) were identified diastolic dysfunction of the left or both ventricles and in 79 patients (group 2) it was absent. Both groups were matched for age and sex. We determined the following echocardiographic parameters: mitral E/A, tricuspid E/A, end-diastolic dimension of left ventricle. In addition to echocardiography, patients underwent the vectorcardiography with the assessment of electrophysiological parameters: the squares of loops P, QRS, T, maximum vector (MV), MV-azimuth and MV- ascent. To compare two independent groups on quantitative grounds used nonparametric methods, the rank correlation and Mann-Whitney test. Differences were considered to be valid when $p < 0.05$.

Results: when comparing the groups revealed that in the 1st group, DAS-28 was higher than in the 2nd ($p < 0.05$): 5.57^{5,17}; 6.15 and 5.32^{4,8}; 5.8 respectively. In the 1st group, the square of loop QRS and the MV- ascent directly correlated with E/A of mitral valve ($p < 0.05$), whereas in the 2nd group, we have established a direct relationship with end-diastolic dimension ($p < 0.05$).

Conclusions: the results indicate that increased activity of rheumatoid arthritis contributes to the development of diastolic dysfunction of the myocardium. In addition, the decrease in E/A observed in diastolic dysfunction, accompanied by electrophysiological remodelling and reduction in electrical activity of the myocardium of the left ventricle, diagnosed during registration of vectorcardiogram. Moreover, even in the absence of diastolic dysfunction, a tendency to its development in the presence of electrophysiological remodelling. This demonstrates the relationship between early electrophysiological, structural-geometric changes in patients with rheumatoid arthritis. Early diagnosis allows for timely start prevention of remodelling in patients with rheumatologic diseases.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1862

THU0167

FALLS IN RHEUMATOID ARTHRITIS (RA) AND ITS RELATION TO DISEASE ACTIVITY, DISABILITY AND PHYSICAL PERFORMANCE TESTS

J.Z. Gaiño¹, M.B. Bertolo², C.S. Nunes¹, C.M. Barbosa¹, Z. Sachetto², M. Davitt¹, E.D.P. Magalhães². ¹UNICAMP, Campinas, Brazil; ²Rheumatology, UNICAMP, Campinas, Brazil

Background: Rheumatoid patients(RA)are known to have an increased falls incidence with a threefold increased risk of hip fracture. Few studies have been conducted to evaluate the relation between falls, disease activity, disability and physical functioning.

Objectives: To evaluate the prevalence of falls in RA and its relation with disease activity, disability and physical performance tests.

Methods: 113 RA patients were evaluated from the outpatient clinic of the Rheumatology Division of the State University of Campinas/Unicamp. Patients were assessed for occurrence of falls in last year, fear of falling, sociodemographic and clinical data (medication, visual impairment, vertigo, physical activity, body mass index, disease duration, rheumatoid factor, lower limb swollen and tender joints, foot tactile sensitivity, disease activity-CDAI and disability-HAQ). Subjects were submitted to Berg Balance Scale-BBS, the Timed Up and Go Test-TUG and 5-Time Sit Down-to-Stand Up Test-SST5 and were divided in "Fallers" and "Non-Fallers" groups. For comparison of groups the chi-squared test, Fisher's exact test and Mann-Whitney were used. Univariate linear regression and multivariate analysis were used to analyse the relation between sociodemographic, clinical data and physical tests with the occurrence of falls. Kruskal- Wallis test was used for analysis of the association of BBS, TUG and SST5 with CDAI and HAQ. The data were analysed with a 5% level of significance.

Results: 52.21% reported the occurrence of falls in the past 12 months and 62.8%were fearful of falling. Comparing "Fallers" and "Non-Fallers", significant differences were noted for HAQ($p=0.0242$), fear of falling($p=0.0196$) and BBS ($p=0.0120$). After univariate logistic regression there was association of falls with income(OR 1.05), HAQ(OR 1.945), fear of falling(OR 2.586) and TUG (OR 1.09). In multivariate model, income was independently linked to falls(OR 1.07). BBS, TUG, SST5 were correlated with CDAI and HAQ($p < 0.05$).