(26.7%) patients were treated with biological therapy and 22 (21.0%) of them were taking tocilizumab. Patients were assessed using a combination of clinical, laboratory and radiological evaluation before and after Tocilizumab therapy. The demographic and clinical characteristics of the patients, the pre-tocilizumab and end of the follow-up acute phase values (Erythrocyte sedimentation rate (ESR), C-reactive protein (CRP)), visual analogue scales (VAS) (Pain, Fatigue, Patient Global), and concomitant therapies were recorded. Other clinical, imaging and laboratory results of the TA patients were obtained from hospital files. The comparative imaging (Computed Tomography and Magnetic Resonance Imaging) results of the patients just before the Tocilizumab therapy and during follow-up were recorded from the hospital data system.

Results: Twenty-two patients (86.4% female) were included in to the analysis. The median (minimum-maximum) age of the patients was 40²⁴⁻⁶³ years and the median disease duration (from diagnosis) was 48¹²⁻¹⁶⁸ months. Before tocilizumab therapy; cyclophosphamide (8 patients (36.4%), conventional immunosuppressives (21 patients, 95.4%) and TNFi (7 patients, 31.8%) were used in addition to corticosteroid therapy. Main indications for tocilizumab therapy was as follows; radiological progression in 9 patients (acute phase was normal in 2 of them), acute phase elevation in 8 patients (4 of them radiologically stable, 4 of them had no radiologic evaluation), physician's decision and clinical symptoms in 3 patients (acute phase normal; 2 of them radiologically stable, one had no radiologic evaluation), in two patients tocilizumab therapy was started out of our centre and data before tocilizumab was not available. Fifteen patients (median follow-up time 15³⁻⁴² months) were evaluated for response to tocilizumab treatment. There was a significant decrease in ESR, CRP, and VAS-patient-global of patients after tocilizumab therapy; median (min-max) ESR (26⁵⁻¹¹⁹ vs. 3 (2-49) mm/s, p=0.02), CRP (39.8 (2.4-149.0) vs. 7.9 (0-92.9) mg/L, p=0.017), VAS-patient-global (50 (0-90) vs. 30 (0-60), p=0.027), respectively. Total of 9 patients had follow-up imaging; 8 of them was radiologically stable and regression was seen in one patient (figure 1).

Table 1:

Patient No	Age/Sex	Age at diagnosis	Disease Duration (months)	Previous treatments	Before Tocilizumab Therapy	ESR mm/h	CRP g/L	VAS (Pain/Fatigue/Patient Global)	Imaging
1	57F	53	48	MTX	Pred. 15 mg/day	34	42.8	0/20/20	Stable
2	48M	31	72	CYC, MTX, SBZ, IFX, ADA	Pred. 10 mg/day	75	81.7	-	Stable
3	47F	37	96	CYC, SBZ, MTX, PE, ETN	MPred. 32 mg/day	24	19.6	50/70/30	Progression
4	25F	20	36	CYC, MTX, LEF, IFX, ADA	Pred. 30 mg/day	11	42.7	80/70/90	Stable
5	55F	53	12	MMF	MPred. 16 mg/day	68	78.1	-	Stable
6	40F	36	12	MTX, PE, IFX	Pred. 40 mg/day	26	28.0	-	-
7	30F	27	12	MTX, PE, IFX	Pred. 40 mg/day	119	198.0	70/90/70	Progression
8	63F	61	12	MTX	-	-	-	Tocilizumab was started out of our center	-
9	28F	23	48	MTX, PE	PE 500 mg/month	103	85.1	0/0/0	-
10	43F	35	168	CYC, MTX	Pred. 40 mg/day	35	62.7	10/20/40	-
11	43F	40	24	CYC, MTX	-	12	16.7	90/90/90	Progression
12	34F	26	72	MTX, IFX	Pred. 50 mg/day	30	3.2	0/0/20	Progression
13	35F	32	36	AZT, MTX	Pred. 10 mg/day	3	1.8	60/60/70	Stable
14	24M	13	72	MTX, IFX	Pred. 5 mg/day	-	-	Tocilizumab was started out of our center	-
15	31F	23	84	CYC, MTX	Pred. 20 mg/day	6	2.7	40/60/30	Progression
16	31F	28	34	CYC	Pred. 30 mg/day	7	2.4	40/70/60	Stable
17	50F	58	12	CYC	Pred. 5 mg/day	42	39.8	70/60/70	Stable
18	48F	46	12	MMF	MPred. 6 mg/day	5	8.8	50/80/50	Progression
19	24M	20	24	CYC, MTX	Pred. 40 mg/day	18	31.7	0/0/10	Progression
20	23F	25	12	-	MPred. 4 mg/day	14	3.9	80/60/50	Progression
21	66F	58	132	AZT	-	2	2.7	80/90/90	-
22	43F	34	60	MTX	Pred. 15 mg/day	30	-	0/70/40	Progression

CYC: Cyclophosphamide; MTX: Methotrexate; SBZ: Sulfasalazine; IFX: Infliximab; ADA: Adalimumab; PE: Prednisolone; ETN: Etanercept; AZT: Azathioprine; MPred: Methylprednisolone; IFX: Infliximab; ADA: Adalimumab; PE: Prednisolone; ETN: Etanercept; AZT: Azathioprine; MPred: Methylprednisolone.

Conclusions: The use of biological therapies in more than a quarter of TA patients suggests that the role of conventional therapies in TA is limited. Although imaging modalities and acute phase elevations are the main factors in the decision of change in treatment, presence of symptoms and physician opinion are also important. Significant response was obtained with the treatment of tocilizumab.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6756

THU0453 EFFICACY OF AB INITIO OR VERY EARLY INTRODUCTION OF IMMUNOSUPPRESSIVE THERAPY IN GIANT CELL ARTERITIS: A MULTICENTER RETROSPECTIVE OBSERVATIONAL STUDY

L. Quartuccio¹, E. Cavallaro¹, F. Angelotti², R. Capecchi², G. Vitiello³, D. Cammelli³, A. Tavoni², S. De Vita¹. ¹Department of Medical Area, Rheumatology Clinic, Udine Academic Hospital "Santa Maria della Misericordia", Udine; ²Department of Internal Medicine, Clinic of Immunology, Pisa; ³Experimental and Clinical Medicine Department, School of Allergy and Clinical Immunology, University of Florence, Florence, Italy

Background: Glucocorticoids (GC) remain the mainstay of treatment of giant cell arteritis (GCA). However, relapses occur in up to 50% of patients when GC are tapered and prolonged courses of GC are associated with serious side effects. In this setting, the outcome is frequently determined by GC-related adverse events

(AEs). Thus, several studies have been conducted on the effectiveness of a GC-sparing immunosuppressive therapy (IT), with conflicting results.

Objectives: To evaluate the effectiveness of IT in a series of GCA patients, in: 1) reducing the risk of GCA relapse; 2) lowering the exposure to GC therapy; 3) minimising the occurrence of steroid-induced AEs.

Methods: We performed a multicenter retrospective observational study including 69 patients (75.4% females; mean age $\pm$ SD 68.7 $\pm$ 7.8 y; median follow-up time 40 months [range 26-65]) diagnosed with GCA, all receiving an adjunctive IT given ab initio (71%) or within 3 months from the start of GC. All but 17 patients (with extracranial involvement) fulfilled the GCA ACR classification criteria. Risk of first relapse of GCA, GC exposure and main AEs were retrospectively analysed. Results were compared with those reported in other studies, all characterised by a lower use of IT.

Results: all patients received GC plus IT: methotrexate up to 20 mg/week in 63 patients, cyclophosphamide in 22, tocilizumab in 12. In the follow-up, a first relapse of GCA occurred in 14 (20.3%) of patients. The rates of first relapse at 12, 24, 36 and 48 weeks were 4.3%, 11.6%, 15.9% and 15.9%, respectively, in comparison with rates of 24.3%, 41.6%, 61%, 64.9%, respectively, in the GC monotherapy arm in the meta-analysis by Mahr,¹ where similar data were available. The median daily prednisone dose at beginning was 50 mg, and decreased to 10 mg at 3 months and to 5 mg at 6 months. The median time required to reduce prednisone to 7.5 mg/day and 5 mg/ was 3 and 5 months respectively (vs. 6.5 and 7.5 months in series by Proven,² where similar data were available). In our cohort, lower cumulative GC use was accompanied by a reduction of AEs, without further toxicity (tab.1).

	Present work	Proven 20032	Mahr 20071	Souza 20143	Alba 20144	Labarca 20165
Number of patients	69	120	77	45	106	286
% use of IT	100%	2%	0%	55.6%	14.2%	0%
Follow up, months	40	120	13,7	54,9	93,6	61,2
Fractility fractures	11,6%	38%	8%	NA	NA	27%
Diabetes	4,3%	9%	12%	33,4%	8,5%	NA
Hypertension	8,7%	22%	NA	66,7%	51%	1,7%
Infections	10,1% (2,9% needing hospitalisation)	31%	NA	23%	NA	17% (all needing hospitalisation)

Conclusions: The use of IT ab initio or very early in the treatment of GCA appears to be effective and safe, lowering the risk of relapses, reducing GC dose and the rate of GC-related AEs. Therefore, the present data support the early introduction of IT in the treatment of GCA.

REFERENCES:

- [1] Mahr AD, et al. Arthritis Rheum 2007;56(8):2789-97.
- [2] Proven A, et al. Arthritis Rheum 2003 49(5):703-8.
- [3] Souza AW, Okamoto KY, Abrantes F, et al. Clinics (Sao Paulo) 2013;68:317-322.
- [4] Alba MA, et al. Medicine (Baltimore) 2014;93(5):194-201.
- [5] Labarca C, et al. Rheumatology (Oxford) 2016;55(2):347-56.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3794

THU0454 INCREASED FDG UPTAKE OF THE MUSCLES IN POLYMYALGIA RHEUMATICA (PMR)

M.A. Cimmino¹, N. Vecchiola¹, M. Bauckneht², D. Camellino³, S. Morbelli², S. Paolino¹, M. Cutolo¹, G. Sambucetti². ¹Department of Internal Medicine, University of Genova; ²Division of Nuclear Medicine, Università di Genova; ³Autoimmunology Laboratory, University of Genova, Genova, Italy

Background: PMR is an inflammatory disease affecting the shoulder and pelvic girdles of elderly patients. Its causes, development mechanisms and anatomical targets of inflammatory damage are still elusive. Synovitis, vasculitis, and bursitis have all been demonstrated in PMR but data on muscles are scanty.

Objectives: Our study evaluated muscular uptake at FDG-PET/CT in patients with PMR.

Methods: Fifty patients with PMR diagnosed according to Bird et al criteria and 50 age- and sex-matched control patients without inflammatory disorders, who underwent FDG-PET/CT to rule out malignancy, were studied. Standard FDG-