joint funding of Amgen, BMS, Celgene, Generic Assays, GSK, Horizon, medac, Mundipharma, Pfizer and Roche.

DOI: 10.1136/annrheumdis-2018-eular.2732

THU0470 NON-CLONAL ELEVATION OF SERUM IMMUNOGLOBULIN FREE LIGHT CHAINS IS PREDICTIVE OF HIP FRACTURE IN BOTH WOMEN AND MEN

S. Amin, E.J. Atkinson, S.J. Achenbach, A. Dispenzieri, R.A. Kyle, S.V. Rajkumar. MAYO CLINIC, Rochester, USA

Background: Proinflammatory cytokines favour uncoupling of bone turnover and decreased bone density and strength, leading to increased fracture risk. Non-clonal elevation of serum immunoglobulin free light chains (sum of kappa and lambda chains) (Σ FLC) may be a global marker of generalised immune stimulation, and has been associated with chronic co-morbidities¹ as well as increased mortality².

Objectives: We examined whether elevated Σ FLC is associated with an increased risk for hip fractures in a population-based cohort.

Methods: We studied Olmsted County, Minnesota, USA residents, age ≥ 50 years, in whom Σ FLC was measured between March 1995 and November 2003 and research authorisation was available. Anyone with a known plasma cell disorder was excluded. Using the Rochester Epidemiology Project, a unique medical records linkage system that allows access to all (inpatient and outpatient) community medical records for Olmsted County residents, we identified all hip fractures that occurred in subjects following their Σ FLC measurement to their last available follow-up or the end of 2015. All available medical records were reviewed by trained nurse abstractors to validate hip fractures identified and to determine their antecedent cause (pathological process [e.g., malignancy], severe trauma [e.g., motor vehicle accidents] and those due to no more than moderate trauma [by convention, equivalent to a fall from standing height or less]). Charlson comorbidity index (CCI) was determined at the time of baseline Σ FLC measurement. We used a Cox proportional hazards model, stratified by sex, adjusting for age, serum creatinine and CCI, to examine whether a Σ FLC ≥ 4.72 mg/dl (levels previously associated with increased mortality in this population¹) is associated with an increased risk for hip fracture.

Results: We studied 15 814 residents [mean age (SD), 64¹⁰ yrs; 8722 women, 7092 men] of whom 796 (9.1%) women and 781 (11.0%) men had a Σ FLC ≥ 4.72 mg/dl. Women and men with an elevated Σ FLC ≥ 4.72 mg/dl had higher CCI [median (IQR) 2 (0,4) vs 0 (0,2); same results for both sexes]. We identified 687 women and 255 men with a hip fracture from any cause (628 women and 220 men had a moderate trauma hip fracture), over 1 12 171 person-years (p-y) and 89 269 p-y of follow-up, respectively. We found that both women and men with Σ FLC ≥ 4.72 mg/dl had an increased risk for hip fracture due to any cause: hazard ratio (HR) [95% CI] 1.31 [1.00, 1.73] in women, HR 1.97 [1.37, 2.83] in men; as well as for moderate trauma hip fractures: HR 1.39 [1.05, 1.84] in women, HR 2.12 [1.46, 3.09] in men.

Conclusions: We found that Σ FLC ≥ 4.72 mg/dl is associated with an increased risk for hip fracture in both women and men, independent of age and chronic co-morbidities. Elevated Σ FLC may be a marker of proinflammatory cytokines detrimental to bone health and warrants further study.

REFERENCES:

- [1] Nakano T, et al. Free Immunoglobulin Light Chain: Its Biology and Implications in Diseases. *Clinica Chimica Acta*. 2011;412(11-12):843-9.
- [2] Dispenzieri A, et al. Use of Nonclonal Serum Immunoglobulin Free Light Chains to Predict Overall Survival in the General Population. *Mayo Clinic Proceedings* 2012;87(6):517-523.

Acknowledgements: Supported by grant P01 AG04875, R01 CA107476 and R01 AG034676 from the National Institutes of Health, U.S. Public Health Service.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6041

THU0471 RISK FACTORS OF LOW BONE MINERAL DENSITY IN PATIENTS WITH SYSTEMIC SCLEROSIS

S.V. Monov¹, D. Monova², M. Stambolova³, R. Shumnalieva¹. ¹Department of Rheumatic Diseases, Medical University – Sofia; ²Department of Internal Medicine, Medical University – Sofia, Medical Institute; ³Department of Internal Medicine, Medical Institute – Ministry of Interior, Sofia, Bulgaria

Background: Systemic sclerosis (SSc) is a multisystem autoimmune disease, characterised by diffuse fibrosis, degenerative changes, and vascular

abnormalities in the skin, joints, and internal organs. The effect of systemic sclerosis on bone density is not well understood.

Objectives: The aim of this study is to evaluate the risk factors of low bone mineral density (BMD) and occurrence of fracture and fracture-related mortality in patients with SSc.

Methods: Demographics, disease manifestations of SSc, biological inflammatory parameters, functional disability, scleroderma health assessment questionnaire, immunological status, BMD (lumbar spine and femoral neck), risk factors for low BMD, fractures, and fracture-related mortality were collected in patients with SSc. BMD was measured by using a dual-energy X-ray absorptiometry in lumbar spine (L1-L4) and femoral neck. Fisher's Exact and Student's t-tests were used to evaluate differences between women with and without low BMD. Logistic regression was used for multivariate analysis.

Results: Forty eight consecutive unselected SSc women were approached. The mean age of women was 48,32 $\pm$ 15,56 years, the mean disease duration was 10,12 $\pm$ 4,78 years, Twenty-nine women (60,42%) had low BMD, of those 11 (37,93%) had osteoporosis, mean BMD in lumbar spine was $-2,86\pm 0,38$ and in femoral neck was $-2,22\pm 0,26$. Twenty three (47,92%) women with SSc were postmenopausal. In correlation analysis and in multiple regression models, there were correlations between BMD and longer duration of SSc (p<0,01), family history of osteoporosis (p<0,05), age (p<0,01), menopause (p<0,01), low body mass index (p<0,02), presence of internal organ involvement (p<0,05), malabsorption syndrome (p<0,05), joint involvement (severe joint pain and erosive arthropathy) (p<0,05), and immunological status (positivity of anti-DNA topoisomerase I antibodies) (p<0,05). 4 women (4/29) with low BMD had a fracture, compared to 2 without low BMD. Fracture-related mortality did not occur in any patients.

Conclusions: Our data suggest that women with SSc are at risk of low BMD and fracture, especially when other risk factors for osteoporosis are present. A number of clinically relevant factors (longer duration of SSc, presence of internal organ involvement, joint involvement, immunological status) are associated with low BMD.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1886

THU0472 FACTORS ASSOCIATED WITH READINESS FOR ADOPTING OSTEOPOROSIS TREATMENT CHANGE

M.I. Danila¹, E.J. Rahn¹, A.S. Mudano¹, R.C. Outman¹, P. Li¹, D.T. Redden¹, F. A. Anderson², S.L. Greenspan³, A.Z. LaCroix⁴, J.W. Nieves⁵, S.L. Silverman⁶, E. S. Siris⁷, N.B. Watts⁸, S. Ladores¹, K.M. Meneses¹, J.R. Curtis¹, K.G. Saag¹.

¹University of Alabama at Birmingham, Birmingham; ²University of Massachusetts Medical School, Worcester; ³University of Pittsburgh, Pittsburgh; ⁴Group Health Cooperative, Seattle; ⁵Helen Hayes Hospital, West Haverstraw; ⁶Cedars-Sinai Medical Center, Los Angeles; ⁷Columbia University Medical Center, New York; ⁸Mercy Health Osteoporosis and Bone Health Services, Cincinnati, USA

Background: Understanding factors associated with the readiness for adopting osteoporosis treatment change may inform the design of behavioural interventions to improve osteoporosis treatment uptake in women at high risk for fracture.

Objectives: To examine the factors associated with the readiness for adopting osteoporosis treatment change among US women with prior fractures.

Methods: US women in the Global Longitudinal Study of Osteoporosis (GLOW) with prior self-reported fractures who were not currently using osteoporosis therapy were eligible to participate in the Activating Patients at Risk for Osteoporosis (APROPOS) Study. Participants' readiness for behaviour change was assessed using a modified form of the Weinstein Precaution Adoption Process Model (PAPM). We defined pre-contemplative participants as those who self-classified in the unaware and unengaged stages of PAPM. Contemplative participants were defined by the undecided, decided not to act, and decided to act stages of PAPM. Bivariate tests and stepwise multivariable logistic regression evaluated the following factors associated with these two levels of readiness for behaviour change: sociodemographic characteristics, health literacy, self-reported history of depression and dementia, previous treatment for osteoporosis, whether participants had been told they had osteoporosis/osteopenia, and whether they had concerns about osteoporosis.

Results: A total of 2684 women were enrolled in APROPOS. Participants were 95% Caucasian, with a mean (SD) age 74.9 (8.0) years and 77% had some college education. Overall, 25% (n=544) self-classified in the contemplative stage of behaviour change. Compared to women who self-classified as pre-contemplative, contemplative women were more likely to be concerned about osteoporosis (adjusted OR [aOR]=3.2, 95% CI 2.3–4.4) and to report prior osteoporosis treatment (aOR 4.3, 95% CI 3.1–6.0). Participants who were told they had osteoporosis had a 12.4 fold odds to be in the contemplative group (95% CI 8.5–18.1), while those who were told they had osteopenia had 4.1 fold odds to be in the contemplative group (95% CI 2.9–5.9).