months decreased mean BMD by 2.6–3.5% (depending on the BMD site) compared to mean BMD decreases of 1.9–2.9% with LMWH. BMD of the spine and hip is significantly lower in patients with LMWH therapy than in subjects with UFH therapy, which is independent of age, sex, menopause, low BMI and altered body composition.

Conclusions: LMWH most likely have less effect on bone turnover when compared to UFH. LMWH for 6 months may not increase the risk of osteoporosis, but longer exposure for up to 24 months may adversely affect BMD. Clinicians should consider monitoring BMD in adults on long-term heparin therapy who are at increased risk of bone loss or fracture.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4255

THU0501 WHAT DOES TRABECULAR BONE SCORE CONTRIBUTE TO A FRACTURE LIAISON SERVICE?

S. Ojeda¹, A. Naranjo¹, A. Saavedra¹, F. Santana¹, A. Molina¹, T. Marrero², O. Suarez², C. Rodriguez-Lozano¹. ¹Rheumatology; ²Radiology, Hospital Univ. Gran Canaria Dr. Negrin, Las Palmas de Gran Canaria, Spain

Background: The utility of the trabecular bone score (TBS) is controversial.

Objectives: To analyse the clinical usefulness of FRAX-TBS in patients with fragility fracture seen in a Fracture Liaison Service (FLS) compared to FRAX without TBS.

Methods: Consecutive outpatients >50 y seen in our FLS were included, excluding patients admitted with hip fracture. The variables included were: age, sex, type of fracture, risk factors for FRAX, FRAX for major and hip fractures with and without TBS, bone densitometry (DXA) and indication of treatment to prevent new fractures.

Results: We included 251 patients, mean age 69 y, 86% women. The most frequent type of fracture was the forearm (n=122) followed by the humerus (n=64). The DXA scan results (taking the lowest value of column/hip) was osteoporosis in 41%. The average TBS was 1.307 (SD 0.103, range 0.961–1.550): forearm fracture 1.313 (SD 0.102), vertebra 1.281 (SD 0.131) and hip 1.291 (SD 0.103). 32% presented normal TBS result 53% partially degraded and 15% degraded (table 1). The results of FRAX with and without TBS were similar, with an average for major/hip fracture of 10.7/4.2 with DXA and 10.6/4.0 with TBS respectively (p>0.05). A high risk of hip fracture (FRAX ≥3) was observed in 43% of patients using FRAX-DXA and 41% using FRAX-TBS. As a high risk of major fracture (FRAX ≥10) in 40% of patients using FRAX-DXA and 41% using FRAX-TBS. In 241 cases (96%) there was concordance in level of risk for FRAX-DXA and FRAX-TBS, while the 10 discordant cases for high risk of fracture were distributed as follows: in 6 cases FRAX-DXA was high and FRAX-TBS was normal, and in 4 cases FRAX-DXA was normal and FRAX-TBS was elevated. In our FLS, a bisphosphonate was prescribed to 169 patients (67%). Regarding patients with a low FRAX risk (n=82), in one patient FRAX-TBS was in favour of treatment. Thus, in 1/251 fractures FRAX-TBS could have influenced the treatment indication compared to FRAX-DXA.

	TBS≥1.350 Normal	TBS<1.350 y≥1.200 partially degraded	TBS<1.200 Degraded
Number of patients	80	134	37
Age, mean (SD)	67 (11)	70 (10)	71 (10)
Sex, women, n (%)	65 (81)	124 (92)	33 (92)
Risk factors, n (%)			
BMI, mean	27	29	31
Previous fracture	8 (10)	34 (25)	7 (19)
Parent's hip fracture	8 (10)	21 ⁽¹⁶⁾	2 (5)
Glucocorticoids	6 (8)	5 (4)	5 (13)
Smoking	9 (11)	20 (15)	6 (16)
Alcohol	2 (3)	4 (3)	1 (3)
Rheumatoid arthritis	3 (4)	2 (1)	0
Secondary osteoporosis	13 (16)	22 (18)	10 (28)
Bone Densitometry, n (%)			
Normal	24 (30)	7 (5)	3 (8)
Osteopenia	37 (46)	65 (49)	12 (32)
Osteoporosis	19 (24)	62 (46)	22 (60)

Conclusions: In our FLS unit, TBS does not help to classify patients' risk. The values of FRAX-TBS are similar to those of FRAX-DXA, not offering advantages when classifying patients with high risk who are candidates for treatment.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2722

THU0502 ASSESSMENT OF SEVERITY OF ATHEROSCLEROTIC LESIONS OF CAROTID ARTERIES IN MEN WITH CORONARY HEART DISEASE DEPENDING ON BONE MINERAL DENSITY AND RISK OF OSTEOPOROTIC FRACTURES

T. Raskina¹, A. Voronkina², M. Letaeva¹, A. Kokov³, O. Barbarash³, E. Maluta².

¹Federal State Educational Institution of Higher Education "Kemerovo State Medical University"; ²State Autonomous Healthcare Institution Kemerovo region "Regional Clinical Hospital of Emergency Medical Care named M. Podgornitsky";

³Federal State Budgetary Scientific Institution "Scientific-Research Institute of Complex Issues of Cardiovascular Disease", Kemerovo, Russian Federation

Objectives: To assess severity of atherosclerotic lesions of carotid arteries (CA) in men with coronary heart disease (CHD) depending on bone mineral density (BMD) and risk of osteoporotic fractures according to FRAX scale.

Methods: The study involved 102 men aged 51–75 years (median age of 61⁵⁵; 65 with CHD, verified by coronary angiography method. All patients underwent dual energy X-ray absorptiometry of lumbar vertebral bodies L1-LIV and femoral neck and colour duplex scanning of extracranial arteries. Assessment of severity of carotid atherosclerosis was carried out by presence of atherosclerotic plaques (ASP) and stenosis of CA, thickness of intima-media (TIM) CA. Threshold TIM for men over age of 50 was considered 0.9 mm (recommendations of American Society of Echocardiography, 2008 Based on results of densitometry value of T-criterion (recommendations ISCD, 2007 men were divided into three groups: 33 patients with osteoporosis (OP, T-criterion ≤2.5), 48 patients with osteopenia (OPE, T-criterion of –1 to –2.5) and 21 men with normal BMD (NBMD, T-criterion ≥–1). On the basis of information on the presence of clinical risk factors of osteoporotic fractures and densitometry data at all included in the study patients FRAX calculator was used to quantify probability of major osteoporotic fractures and hip fracture in next 10 years.

Results: It was found that 86.3% of patients value TIM exceeds threshold value (0.9 mm). Thickening of intima-media complex was in men with OP in 88.0% of cases, with OPE – in 87.5%, with NBMD – in 81.0% without statistically significant differences (p>0.05). In a comparative analysis of atherosclerotic lesions CA found that men with OP significantly were more likely to have ASP in CA (75.8%) compared to patients with OPE (43.7%, p=0.010) and NBMD (38.1%, p=0.016). Stenosis of CA in all investigated men was discovered in 54 cases (53%). Comparative analysis showed that in patients with OP percentage of patients with stenosis of CA was 2 times more than men with NBMD (75.8% vs 38.1%, p=0.006) and was significantly higher than proportion of such patients in the group with OPE (44.0%, p=0.004). In all included the study patients 10 year absolute risk of major osteoporotic fracture by FRAX was equal to 9.88±7.22, risk of hip fracture – 3.97 ±6.27. There was inverse correlation of TIM with T-criterion (r=–0.21, p=0.035) and with BMD (r=–0.20, p=0.045) at level of hip and reliable direct correlation between TIM and risk of hip fracture by FRAX (r=0.21, p=0.035).

Conclusions: In men with CHD aged over 50 years low BMD (T-criterion is less than 2.5) is associated with more severe carotid atherosclerosis, and thickening of intima-media CA – with an increased risk of hip fracture.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5548

THU0503 COST ANALYSIS OF ANTI-OSTEOPOROTIC DRUGS IN REAL-WORLD CLINICAL PRACTICE

V. Orlando¹, V.M. Monetti¹, F. Guerriero¹, V. Russo¹, R. Piscitelli¹, M. Ruggeri², E. menditto¹. ¹CIRFF Center of Pharmacoeconomics, University of Naples Federico II, Naples; ²School of Economics, Catholic University of Sacred Heart, Rome, Italy

Background: Nonadherence can reduce treatment effectiveness and can negatively affect healthcare costs and thus the treatments' cost-effectiveness. Adherence in the setting of osteoporosis has been shown to be just as problematic, if not worse, than that in other chronic diseases. Economic evaluations based on modelling are commonly used to compare alternative treatment strategies in osteoporosis, to support decision-makers and to inform treatment guidelines.

Objectives: The aim of this study is to analyse healthcare costs of osteoporosis and to build a economic model cost-effectiveness of pharmacological interventions based on real world data.

Methods: The data of the analysed subjects were drawn from the administrative databases of four Local Health Authorities in the Abruzzo Region. Patients 60 years of age or older were included if at least one prescription for any