antiosteoporosis drugs had been filled in between January 1, 2006 and December 31, 2006. The patients were classified as exposed or not exposed to osteoporotic fracture. For each group, compliance with antiosteoporosis drugs was calculated. Patients were considered compliant if their Medication Possession Ratio (MPR) was $\geq 80\%$. The cost analysis was conducted taking each healthcare service into account, i.e. drug therapy, diagnostic tests and hospitalisation admissions, during the study period. A hypothetical scenario based on the real-life available evidence was constructed. The mean level of adherence to populate the hypothetical scenario of "full adherence" was set at $MPR > 80\%$. The model built by adding a step value, constrained by a normal random variable, to the real-world adherence of each subject so that the subject shifted to the hypothetical scenario of full adherence, in order to quantify the clinical outcome (number of fractures) achievable in the hypothetical scenario. Cost-effectiveness of full adherence compared to real world adherence was expressed in terms of Incremental Cost Effectiveness Ratio (ICER) and the number of fractures avoided was set as an effectiveness unit of measure.

Results: The mean annual healthcare cost per fracture avoided was € 247.44, of which medical treatments and diagnostic tests accounted for € 103.60 (41.9%) and € 143.84 (58.1%), respectively. The mean annual healthcare cost per fractured patient was € 1,044.85, of which medical treatments, diagnostic tests and hospitalizations for osteoporotic fracture accounted for € 88.73 (8.5%), € 169.48 (16.2%) and € 786.65 (75.3%), respectively.

Conclusions: Costs per fractured patients resulted to be about four times greater than those of not fractured patients. Therefore, only enhancing adherence to medication may lead to reductions in the number of patients requiring hospitalisation.

REFERENCES:

- [1] Hilgismann M, Boonen A, Rabenda V, Reginster JY. The importance of integrating medication adherence into pharmacoeconomic analyses: the example of osteoporosis 2012;159–166.
- [2] Scotti L, Arfè A, Zambon A, Merlino L, Corrao G. Cost-effectiveness of enhancing adherence with oral bisphosphonates treatment in osteoporotic women: an empirical approach based on healthcare utilisation databases. *BMJ open* 2014;4(3),e003758.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6620

THU0504 UTILISATION OF ANTI-OSTEOPOROTIC DRUGS IN REAL-WORLD DATA: A STUDY OF GENDER-RELATED DIFFERENCES

V. Orlando¹, V.M. Monetti¹, F. Guerriero¹, V. Russo¹, A. Moretti², A. Piscitelli¹, R. Piscitelli¹, G. Iolascon², E. Menditto¹. ¹CIRFF Center of Pharmacoeconomics, University of Naples Federico II; ²Department of Medical and Surgical Specialties and dentistry, University of Campania Luigi Vanvitelli, Naples, Italy

Background: Osteoporosis is mostly defined as the disease of women, because the prevalence and fracture rates are much higher in postmenopausal women than in older men. However, there has been increasing recognition that male osteoporosis also represents an important burden as a common cause of morbidity, mortality and health care expenditure. Also, men are more likely than women to have osteoporosis that is undiagnosed and undertreated.

Moreover, bone fracture are important factors of high mortality and morbidity rates in osteoporotic patients. Lack of persistence is common among subjects using oral anti-osteoporotic drugs, and leads to increased risk of fragility fracture.

Objectives: The aim of this study is to perform gender specific analysis regarding the persistence to antiosteoporosis drugs.

Methods: We conducted a retrospective cohort study using administrative data from four local health authorities in the Abruzzo Region (Central Italy), which comprise about 9 00 000 inhabitants (68% of the overall regional population). Patients 60 years of age or older were included if at least one prescription for any antiosteoporosis drugs had been filled in between January 1, 2006 and December 31, 2006. Persistence estimates over time (discontinuation rates were assessed at 365 days) were derived using Kaplan–Meier survival analysis, stratifying for gender, considering treatment discontinuation as failure event and comparing differences using Log-rank test (1 degree of freedom).

Results: The final cohort consisted of a total of 7867 patients (87.2% women). The mean patient age for both genders at the index date was 74.5 years. The crude analysis of long-term gender persistence showed a significant difference between women and men: the relative number of persistence patients after 1 year was 66.4% in men and 44.7% in women. The Kaplan Meier plots of time to persistence start to differ for men vs women approximately 60 days after treatment start (figure 1).

Conclusions: The results showed that gender influences adherence to therapy and this is an issue that could be taken in strong consideration in tailor intervention to improve adherence.

REFERENCES:

- [1] Cawthon PM. Gender differences in osteoporosis and fractures. *Clinical Orthopaedics and Related Research* 2011;469(7):1900–1905.
- [2] Bor A, Matuz M, Gyimesi N, et al. Gender inequalities in the treatment of osteoporosis. *Maturitas* 2015;80:162–169.
- [3] Christopher J, et al. Sex and gender considerations in male patients with osteoporosis. *Clinical Orthopaedics and Related Research* 2011;469(7):1906–1912.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5572

THU0505 INFLUENCE OF ORAL PREDNISOLONE ON EFFECT OF DENOSUMAB ON OSTEOPOROSIS IN PATIENTS WITH JAPANESE RHEUMATOID ARTHRITIS; 36 MONTHS OF FOLLOW-UP ~A MULTICENTER REGISTRY STUDY~

Y. Kanayama¹, Y. Hirano², N. Takahashi³, S. Asai³, N. Ishiguro³, T. Kojima³, on behalf of TBCR-BONE study group. ¹Orthopedic Surgery And Rheumatology, Toyota Kosei Hospital, Toyota; ²Rheumatology, Toyohashi Municipal Hospital, Toyohashi; ³Orthopedic Surgery, Nagoya University Graduate School of Medicine, Nagoya, Japan

Background: Denosumab (DMB) is a fully human monoclonal antibody to the receptor activator of nuclear factor-kappaB ligand (RANKL) that blocks its binding RANK, inhibiting the development and activity of osteoclasts, decreasing bone resorption, increasing bone density and reducing fracture risk. Osteoporosis(OP) is more frequent in patients with rheumatoid arthritis(RA) than in the general population due to active systemic inflammation as well as the use of glucocorticoid and immobility. We investigated the influence of oral prednisolone on effect of DMB in patients with Japanese RA from initiation to 36 months at this time.

Objectives: This prospective study investigated the efficacy of DMB for 36 months on glucocorticoid-induced OP in RA patients.

Methods: Patients with a diagnosis of RA according to the 2010 ACR/EULAR criteria who had been prescribed DMB from Tsurumi Biologics Communication Registry (TBCR)-BONE between October 2013 and October 2014 were enrolled. The final study cohort of 56 patients received continuous DMB therapy more than 36 months. The DMB dose was 60 mg at once every 6 months. In all cases native or activated vitamin D has been used. We reviewed the results for 36 months about the increase and decrease of bone mineral density(BMD) of lumbar spine (LS) and total hip (TH) by DEXA and bone turnover markers, intact n-terminal propeptide type I procollagen(PINP) and tartrate-resistant acid phosphatase form 5b (TRACP-5b).

Results: In the patients receiving oral prednisolone group(n=20, GC+) and not receiving group(n=36, GC-), the number of female was each 18 (90%) and 35 (98%) cases(p=0.288). The mean age was 69.5±7.0 and 70.5±6.6 years old (p=0.700); disease duration was 15.9±9.5 and 16.3±13.7 years (p=0.688); the body mass index was 20.6±3.5 and 19.5±3.0 (p=0.225) and the FRAX was 33.8