THU0537

ASSOCIATION BETWEEN BIG FIVE PERSONALITY TRAITS AND RESPONSE TO MULTIDISCIPLINARY PROGRAM IN PATIENTS WITH CHRONIC LOW BACK PAIN: A PROSPECTIVE STUDY

M.E. Ibrahim^{1,2}, K. Weber³, S. Genevay¹. ¹Rheumatology, University Hospitals of Geneva, Geneva, Switzerland; ²Rheumatology, Suez Canal University, Ismailia, Egypt; ³Medical Direction, University Hospitals of Geneva, Geneva, Switzerland

Background: Chronic back pain is a multifactorial condition, which cannot be thoroughly addressed without understanding the psychological factors that influence its onset, maintenance and the response to its management. Patient attitudes and personality traits often determine how they experience pain, how they adjust to it, and their response to the different management strategies.

Objectives: This study aimed to identify whether patients with specific Big Five personality traits are more likely to seek medical care, and to uncover whether there is an association between personality and the patient's response to multidisciplinary functional rehabilitation program.

Methods: 97 patients (57% male) aged 41.1 (10.1) with chronic low back pain, enrolling in an intensive 4 week multidisciplinary functional rehabilitation program, completed the NEO Personality Inventory-Revised (NEO PI-R) at baseline. Outcome was assessed at treatment admission and discharge, and at 6 months follow up, including Core Outcome Measure Index (COMI), Tampa Scale for Kinesiophobia (TSK), Oswestry Disability Index (ODI) and Hospital Anxiety and Depression Scale (HADS).

Results: Both men and women had significantly lower scores in the Openness to Experience domain and significantly higher scores in the Conscientiousness domain than the average general population. At baseline, Neuroticism was found to correlate positively with TSK and HADS. Low Openness to Experience, and low Extraversion correlated with high HADS at baseline as well. Analysis of the follow up data showed a significant reduction in the COMI, ODI, HADS and TSK at the end of the program, and at 6 months follow up. There was no correlation between the improvement in COMI, ODI or TSK scores after 6 months and any of the 5 domains of the NEO PI-R.

Conclusions: Low Openness to Experience and high Conscientiousness seem to be key factors of chronicity in patients with low back pain. Significant decrease in pain, disability as well as depressive and anxious moods showed that these patients were effectively treated by the multidisciplinary functional rehabilitation program regardless of their personality traits.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4416

THU0538

PATIENTS WITH CHRONIC LOW BACK PAIN ORIENTED TO FUNCTIONAL RESTORATION PROGRAM ARE YOUNGER, WITH HIGHER SMOKING AND LONGER SICK LEAVE: A CROSS-SECTIONAL STUDY OF 166 PATIENTS

M. Assadourian¹, F. Bailly¹, P. Letellier¹, S. Ascione¹, M. Le Ralle¹, A. Potel¹, L. Morardet¹, V. Foltz¹, A. Dupeyron², N. Ibrahim³, I. Griffoul-Espitalier⁴, B. Duplan⁵, J. Beaudreuil⁶, B. Fautrel¹, L. Gossec¹. ¹Pitié Salpêtrière Hospital, Paris; ²Nîmes Hospital, Nîmes; ³La Source Hospital, Orléans; ⁴Tours Hospital, Tours; ⁵Reine Hortense Hospital, Aix les Bains; ⁶Lariboisière Hospital, Paris, France

Background: Functional Restoration Programs (FRP) are multi-disciplinary programs that have demonstrated effectiveness to promote functional status and return to work in patients with chronic low back pain (cLBP)^{1, 2, 3} and moderate or high disability.

Objectives: To explore the characteristics of cLBP patients oriented or not towards a FRP.

Methods: This was an observational cross-sectional, multi-centric study in France in 2017. Consecutive patients with cLBP (>6 weeks) with or without radiculalgia, aged 18–66 years and visiting the rheumatologist for cLBP were included. Patient and disease characteristics were collected. Patients oriented or not towards a FRP were compared in particular for demographic characteristics, duration of sick leave over the past 12 months, smoking, ongoing aerobic physical activity (patient-reported as <or at least 30 min sessions twice a week), anxiety or

depression (Hospital Anxiety and Depression Scale ≥11), disability (Oswestry >40%) and kinesiophobia (Tampa Kinesiophobia Scale ≥40).

Results: In all, 166 patients were analysed: median age 48 years, median cLBP duration 5 years, and 100 (60%) were females. Overall, 62 (37%) were proposed for a FRP. Patients oriented towards a FRP, versus not, were more frequently male (50% vs 44%, p=0.04), younger (median 46 vs 52 years, p=0.02), and had lower pain levels (median 6 vs 7 of 0–10 numeric scale, p=0.006). They had longer sick leave (mean 18 vs 10 weeks/year, p=0.006), were more frequently smokers (33% vs 23%, p=0.02), had similar pain duration (60 month vs 72, p=0.32), functional disabilities (39% vs 49%, p=0.22), kinesiophobia (62% vs 55%, p=0.65), anxiety (52% vs 55%, p=0.60) and depression score (40% vs 35%, p=0.54), but less physical activity practice (38% vs 54%, p=0.04).

Conclusions: Patients oriented towards a FRP were younger and more often males, with higher smoking and longer sick leave. It appears that kinesiophobia, functional status and duration of low back pain didn't influence the orientation toward a FRP. More comprehensive assessments of patients with cLBP are needed; multifactorial questionnaires such as StartBack or BP2 may be useful to orient patients towards FRPs.

REFERENCES:

- [1] Poulain C, et al. Eur Spine J 2010 Jul;19(7):1153–61.
- [2] Véron O, et al. Ann Readapt Med Phys 2008;51(8):642–9.
- [3] Roche-Leboucher, et al. Spine 2011 Dec 15;36(26):2235–42.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3968

THU0539

MECHANICAL TRACTION FOR LUMBAR RADICULAR PAIN: SUPINE OR PRONE? A RANDOMISED CONTROLLED TRIAL

M. Bilgilişoy Filiz¹, Z. Kılıç², A. Uçkun³, T. Çakır⁴, S. Koldaş Doğan¹, N. F. Toraman¹. ¹Physical Medicine and Rehabilitation, Antalya Training and Research Hospital, Antalya, Turkey; ²Physical Medicine and Rehabilitation, Ankara Atatürk Training and Research Hospital, Antalya, Turkey; ³Physical Medicine and Rehabilitation, Mut State Hospital, Mersin; ⁴Physical Medicine and Rehabilitation, Private Likya Hospital, Antalya, Turkey

Background: Mechanical lumbar traction has been used to treat spinal disorders and low back pain for centuries, since Hippocrates.^(2,3) Although the supine position is generally preferred, the patient may also be positioned prone during traction treatment.^(4,5) However, to the best of our knowledge, no studies have compared the effects of lumbar traction in these two different positions for patients with chronic LBP.

Objectives: To compare the effects of mechanical lumbar traction either in the supine or prone position with conventional physical therapy (PT) in patients with chronic low back pain (LBP) and lumbosacral nerve root involvement in terms of disability, pain, and mobility.

Methods: Participants (n=125) were randomly assigned to receive 15 sessions of PT with additional mechanical lumbar traction either in the supine position (supine traction group), or in the prone position (prone traction group), or only PT without traction (PT only group). Patients were assessed at baseline and at the end of the PT sessions in terms of disability, pain, and mobility. Disability was assessed using the modified Oswestry Disability Index (ODI); pain was assessed using a visual analogue scale (VAS); and lumbar mobility was assessed using the modified lumbar Schober test (mLST).

Results: 118 patients completed the trial. All groups improved significantly in the ODI, VAS, and mLST (p<0.05) (table 1). In the between-group analysis, improvements of ODI and VAS were found significantly better in the prone traction group compared with the PT only group (adjusted p=0.031 and 0.006, respectively).

Conclusions: Addition of traction in prone position to other modalities resulted in larger immediate improvements in terms of pain and disability, and the results suggest that when using traction, prone traction might be first-choice. Further research is needed to confirm the benefits of lumbar traction in the prone position.

REFERENCES:

- [1] Marketos SG, Skidas P. Hippocrates. The father of spine surgery. Spine 1999;24:1381–7.

Abstract THU0539 – Table 1

	Supine traction group (n=39)			Prone traction group (n=39)			Only PT group (n=40)		
	preintervention	postintervention	P	preintervention	postintervention	P	preintervention	postintervention	P
ODI	50.4±22.5	33.2±15.5	.001	53.8±15.1	35.9±15.9	.001	49.5±15.1	40.9±17.6	.002
VAS	8.0±1.6	4.3±2.2	.001	8.0±1.7	3.7±2.7	.001	7.7±1.5	5.2±2.3	.001
mLST	19.4±1.9	20.7±1.8	.001	18.8±1.8	20.5±1.3	.001	19.7±1.9	20.8±1.7	.001

- [2] Krause M, Refshauge KM, Dessen M, et al. Lumbar spine traction: evaluation of effects and recommended application for treatment. *Manual therapy* 2000;5:72–81.
- [3] Harte AA, Gracey JH, Baxter GD. Current use of lumbar traction in the management of low back pain: results of a survey of physiotherapists in the UK. *Archives of physical medicine and rehabilitation* 2005;86:1164–9.
- [4] Fritz JM, Lindsay W, Matheson JW, et al. Is there a subgroup of patients with low back pain likely to benefit from mechanical traction? Results of a randomized clinical trial and subgrouping analysis. *Spine* 2007;32:E793–800.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1244

THU0540

INCREASED WORK CAPACITY IN CHRONIC LOW BACK PAIN PATIENTS AFTER A MULTIDIMENSIONAL PROGRAM ASSOCIATED WITH DECREASE IN FEAR AND APPREHENSION

M. Norberg^{1,2}, L. Belgrand². ¹Rehabilitation, Centre medical et thérapeutique La Lignière, Gland; ²Rehabilitation, Lavey Medical SA, Lavey Les Bains, Switzerland

Background: Chronic low back pain has a high burden in our society. Almost 85% of the population would be affected from low back pain. Less than 10% would be chronic but they have an important economic impact since they have the highest costs. According to Vlayen, the kinesiophobia (avoidance of movement) is the most important prognostic factor when evaluating the return to work.

Objectives: To study the importance of kinesiophobia, fear and anxiety in the chronic low back (CLB) pain patient and their relationship to workability after a multidimensional intensive treatment program.

Methods: We included 850 patients who had followed an outpatient program of functional restoration during 3 weeks. The program was composed of physical exercises, occupational therapy and psychological group discussions. They were followed over 1 year. Using different questionnaires (TSK -Tampa scale of Kinesiophobia, FABQ, Pact -subjective work capacity-, Phoda, SF 36), physical performances tests (muscular endurance: Shirado, Biering-Sörensen, Bruce; lumbar mobility, Pile lifting test) we analysed the important factors for their work capacity.

Results: There were a clear relationship between a decrease in kinesiophobia and an increase of work capacity. Globally, the work capacity increased from 41.2% to 79%. There were no long standing increases in muscular performances, but the important change appeared in the decrease in the physical part of FABQ (14 to 9/24) and the SF36 limitations physical health/emotional problems (19.4% to 51.8%/36 to 65.7%).

Conclusions: A multidimensional intensive program including approaches on fear and apprehension has an important impact on work capacity. This observation is important to take into account in creating functional restoration programs.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2887

THU0541

THE EFFECT OF PHYSICAL THERAPY ON CLINICAL AND QUALITY OF LIFE IN CHRONIC NECK PAIN PATIENTS: A RANDOMISED CONTROLLED TRIAL

N. Şahin¹, M. Şahin². ¹Physical Medicine and Rehabilitation, Balıkesir University, Balıkesir; ²Physical Medicine and Rehabilitation, Magnet Hospital, Kayseri, Turkey

Background: In the treatment of chronic neck pain (CNP), education, medical treatment, exercise and physical therapy (PT) modalities are in place. However, there are not enough studies on the efficacy of PT modalities in CNP.

Objectives: To evaluate the effectiveness of the addition of PT modalities to exercise and medical treatment in relieving pain and improving the functional status of patient with CNP.

Methods: 80 patients with CNP were included in a randomised, controlled trial. Patients were assigned in two groups randomly. Treatment group (TG) received conventional PT (hot pack (HP), ultrasound (US), Transcutaneous Electrical Nerve Stimulation (TENS)) treatment in addition. PT was applied ten sessions. HP treatment was applied in 20 min. US treatment was applied with 1.5watt/cm² dose and continuous type in 10 min. TENS treatment was applied with conventional type in 30 min. All patients were informed about correct posture and daily life activities. Both groups received home-based exercises program and analgesic medical treatment if it is necessary. Patients were evaluated before and after therapy and 3th month later by Visual Analogue Scale (VAS), cervical range of motion (ROM), Beck Depression Scale (BDS) and short form-36 (SF-36).

Results: In both groups there is significant improvement in VAS, cervical ROM, SF 36 and BDS after treatment ($p < 0.01$). In TG significant improvement was seen on 3th month follow up. But the significant improvement was not seen on 3th month in control group (CG).

There is no significant difference between groups for VAS, SF-36 parameters and BDS before and after treatment ($p > 0.05$). There was a significant improvement in VAS, SF-36 parameters and BDS in the TG compared with the CG at the end of therapy and 3 months post-treatment ($p < 0.01$).

Conclusions: Medical treatment and exercise with HP, US and TENS therapy was effective on both pain and disability during the treatment. This improvement kept on 3th month follow up. Also same improvement was seen on mood and life quality. Exercise has better effects on after treatment, but these good effects decrease on 3th month follow up. So we think physical medicine modalities should be used in CNP with disability.

REFERENCES:

- [1] Binder A. The diagnosis and treatment of nonspecific neck pain and whiplash. *Eura Medicophys* 2007;43:79–89.
- [2] Leaver AM, Refshauge KM, Maher CG, et al. Conservative interventions provide short-term relief for non-specific neck pain. *J Physiother* 2010;56:73–84.
- [3] Wang WTJ, Olson SL, Campbell AH, et al. Effectiveness of Physical Therapy for Patients with Neck Pain. A Individualized Approach Using a Clinical Decision-Making Algorithm. *Am J Phys Med Rehabil* 2003;82:203–218.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2689

THU0542

EVALUATION OF THE EFFECTIVENESS OF ULTRASOUND GUIDED EPIDURAL CORTICOSTEROID INJECTION AND PULSED ELECTROMAGNETIC FIELD STIMULATION IN CHRONIC LOW BACK PAIN

S.A. Tabra¹, M. ElSergany², M.E. Mowafy¹, H. Khalaf¹, G. Elsharnoby³.

¹Rheumatology and Rehabilitation, Faculty of Medicine, Tanta University;

²Rheumatology and Rehabilitation, Faculty of medicine Tanta university; ³clinical pathology, Faculty of Medicine, Tanta University, Tanta, Egypt

Background: Epidural injections are one of the most common nonsurgical interventions for managing chronic low back pain. They have been used to treat radicular pain from herniated discs, spinal stenosis, and axial spinal pain. Pulsed electromagnetic field stimulation therapy (PEMFs) provides a noninvasive and safe method to treat the site of injury, the source of pain, inflammation by modulating factors involved in pain signalling and the inflammatory response.

Objectives: To assess the improvement in patients with chronic low back pain treated with epidural steroid injection or Pulsed electromagnetic field stimulation. To compare the efficacy of epidural steroid injection and pulsed electromagnetic field stimulation in treatment of patients with chronic low back pain.

Methods: In this study; sixty patients with chronic discogenic low back pain (diagnosed clinically and by magnetic resonant imaging of lumbosacral region) with or without radicular pain of at least 6 months duration were selected. We excluded patients with other causes of back pain as spondylolithesis, inflammatory, infective, neoplastic, traumatic causes. Patients were randomly divided into two equal groups (30 patients each); after informed consent; group I treated by ultrasound guided caudal epidural injection of 40 mg methylprednisolone and 2 ml 2% lidocaine and 20 ml of 9% NaCl twice one week in between and group II received PEMFs daily for 4 weeks. And all patients will be instructed to follow an exercise program. All patients were assessed clinically, functionally by Oswestry Disability Index (ODI) and by measuring serum level of beta-endorphin by ELIZA before, at the end treatment and six months after the end of treatment.

Results: In both groups; there was highly significant improvement in pain after treatment ($P1 < 0.0001$) the mean value of the (VAS) was 8.13+0.63, 7.70 +1.34 respectively before treatment and 3.33+2.63, 2.30+2.32 respectively after treatment. Still further significant improvement at the end follow up ($P1 < 0.0001$) in group I. There was highly significant improvement ($p < 0.0001$) of functional status in both groups after treatment and at follow up period as compared to before treatment but there was significant decrease of functional status at follow up period as compared to after treatment in group II. There was significant improvement of serum level of beta endorphin ($p > 0.05$) in both groups after treatment and follow up period as compared to before treatment but there was insignificant difference at follow up period as compared to after treatment. Our result showed insignificant difference between two groups in clinical, functional or laboratory parameters.

Figure 1. Comparison regarding low back pain as measured by VAS before, after treatment and at 6 months follow up

Figure 2. Comparison regarding serum beta endorphin before, after treatment and at 6 months follow up

Conclusions: Epidural corticosteroid injection and Pulsed electromagnetic field stimulation are effective tools in management of chronic low back pain

REFERENCES:

- [1] Abdi S, Datta S, Trescot AM, et al. Epidural steroids in the management of chronic spinal pain: a systematic review. *Pain Physician* 2007;10(1):185–212.
- [2] Rohde C, Chiang A, Adipoju O, et al. Effects of pulsed electromagnetic fields on interleukin-1 beta and postoperative pain: a double-blind, placebo-controlled, pilot study in breast reduction patients. *Plast Reconstr Surg* 2010;125(6):1620–9.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2069

THU0543

COMPARISON OF THE EFFECTIVENESS OF LOW LEVEL LASER THERAPY AND STEROID INJECTION IN MODERATE CARPAL TUNNEL SYNDROME: A RANDOMISED CONTROLLED TRIAL

S. Badil Güloğlu¹, M. Bilgilişoy Filiz¹, K.K. Kılıç², S. Koldaş Doğan¹, I. Erdem Toslak², N.F. Toraman¹. ¹Physical Medicine and Rehabilitation; ²Radiology, Health Sciences University, Antalya Training and Research Hospital, Antalya, Turkey, Antalya, Turkey

Background: Recent studies have reported that low-level laser therapy (LLLT) reduces pain and improves function in carpal tunnel syndrome (CTS). However, to the best of our knowledge, there exist no studies that compares the effectiveness of LLLT and steroid injections in moderate CTS.

Objectives: Is to evaluate the effectiveness of LLLT and steroid injection in moderate CTS, by comparing the symptoms, the electrophysiologic and the Magnetic Resonance Imaging (MRI) findings of the patients.

Methods: 87 patients with moderate CTS were included in the study. 87 patients were randomised to the steroid or LLLT groups. LLLT was applied to the carpal tunnel area (6 joules/5 min) five times per week, for a total of 10 sessions. Steroid injections were performed once after randomization. Numbness and pain of the patients were assessed using a Visual Analogue Scale (VAS), disability was assessed using the Quick arm-shoulder-hand (Quick DASH) disability questionnaire. Rough and fine grasping strength tests, Tinel and Phalen tests and

electrophysiological test were performed. All the patients were evaluated by MRI in terms of median nerve cross-sectional area, median nerve intensity and intensity rate, and median nerve palmary spring rate. All tests were performed before the treatment, and 1. month and 6. months after the treatment, except the MRI evaluations, which were done before and after 6. months of the treatment.

Results: The study was completed with 80 patients, 133 wrists at the end of six months. There were no significant differences between the groups in terms of demographic and baseline clinic characteristics, electrophysiologic and MRI findings of the patients ($p>0.05$). While the VAS scores of numbness and pain and Quick Dash symptom scores were better in the steroid group in the first month, there were no significant differences between the groups in the sixth month. In both groups, there were similar significant improvements in the Phalen and Tinel tests, rough and fine grasping forces, median nerve motor distal latency value in the first and sixth months after the treatment in comparison to pretreatment values ($p<0.05$). Median nerve sensory distal latency and median sensory NCV showed significant improvements in the 1st and the 6th months after treatment in comparison to pretreatment only in the steroid group ($p<0.05$). In both groups, there were similar significant improvements in the median nerve intensity rate in the radioulnar plane and median nerve palmary spring rate on the MRI after the treatment. Median nerve intensity rate was improved in both groups also at the pisiform plane but this improvement was significant only in the steroid group.

Conclusions: It is possible to conclude that LLLT reduces the signs and symptoms of the patients who has moderate CTS as well as the steroid treatment. Our study offers a data regarding LLLT's efficiency in short-intermediate terms. Further studies are needed to make comments about long-term efficiency of LLLT. LLLT is a noninvasive treatment option in moderate CTS and with a similar efficacy as steroid treatment.

REFERENCE:

- [1] Evcik D, Kavuncu V, Kahir T, Subasi V, Yaman M. Laser therapy in the treatment of carpal tunnel syndrome: a randomized controlled trial. *Photomed Laser Surg* 2007;25:34–39.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4218

THU0544

STUDY OF THE EFFECTS OF OZONE VERSUS STEROID INJECTION ON PATIENTS WITH RECALCITRANT TENNIS ELBOW; A CLINICAL TRIAL

S.A. Raeissadat¹, M.H. Abdollahzadeh². ¹Physical Medicine and Rehabilitation Department of Shahid Beheshti University of Medical Sciences, Research Development Center of Shahid Modarres Hospital; ²Physical Medicine and Rehabilitation Department of Shahid Beheshti University of Medical Sciences, Physical Medicine and Rehabilitation Research Center, Tehran, Iran, Islamic Republic of Iran

Background: Tennis elbow is a prevalent musculoskeletal disorder. In recent years ozone injection has been proposed as a treatment for many musculoskeletal disorders. This study aims to compare the effect of ozone injection with steroid in patients with recalcitrant tennis elbow.

Objectives: The aim of this study is to compare the effect of ozone with the standard treatment of steroid injection in patients with tennis elbow, resistant to conservative treatments. This comparison is made according to VAS score, Pressure pain threshold (PPT) and modified Mayo clinic performance index for elbow.

Methods: In this study 64 patients with tennis elbow, which have had the symptoms for more than 3 months and were resistant to conservative treatments, were randomised to two groups. In steroid injection group 40 mg of methylprednisolone acetate and in ozone injection group 4 ml of ozone with concentration of 15mcg/ml was injected. 32 patients in steroid and 29 patients in ozone injection group finished this study. Data were gathered before injection and 2 and 6 months after that by VAS score, modified Mayo clinic performance index for elbow and PPT (measured by the means of an algometer). Variables were compared between the two groups and also the changes in each group have been measured in reference to baseline data.

Results: In both groups VAS score, Mayo clinic performance index and PPT improved significantly in 2 and 6 month follow up. Steroid injection was significantly better than ozone injection in improving 2 and 6 months pressure pain threshold and 6 months Mayo clinic performance index. Other data didn't show significant difference between the two groups.

Conclusions: Both steroid and ozone injection improved pain and function in patients with recalcitrant tennis elbow for at least 6 months. Steroid injection was superior to ozone injection in improving PPT during the 2 and 6 month follow up and function improvement in steroid group was significantly more than ozone group after 6 month.