significant, among those treated with other non-TNFs (HR 2.4; 95% CI 0.9, 6.3; $p=0.08$) compared with those treated with abatacept. Infection-related medical costs were lower for abatacept compared with other non-TNFs as shown in the table 1.

Conclusions: In the real-world setting in the United States, pts treated with abatacept versus other tDMARDs tended to have a lower risk for infection-related hospitalisation and lower infection-related medical costs.

Disclosure of Interest: D. Paul Employee of: Bristol-Myers Squibb, L. McDonald Employee of: Bristol-Myers Squibb, A. Marshall Shareholder of: Bristol-Myers Squibb, Employee of: Bristol-Myers Squibb, T. Curtice Employee of: Bristol-Myers Squibb, M. Lingohr-Smith Grant/research support from: Bristol-Myers Squibb, Employee of: Novosys Health, B. Menges Grant/research support from: Bristol-Myers Squibb, Employee of: Novosys Health, J. Lin Grant/research support from: Bristol-Myers Squibb, Employee of: Novosys Health

DOI: 10.1136/annrheumdis-2018-eular.2323

THU0646 EVALUATING THE QUALITY OF CARE FOR RHEUMATOID ARTHRITIS

C. Barber¹, D. Marshall¹, E. Szefer², D. Thompson², D. Lacaille³. ¹University of Calgary, Calgary; ²Emmes Canada; ³University of British Columbia, Vancouver, Canada

Background: The Arthritis Alliance of Canada (AAC) has developed performance measures (PMs) to evaluate RA care quality.

Objectives: To operationalize and report on 4 PMs using administrative data for British Columbia (BC), Canada: PM1) percentage of patients with new onset RA with at least one visit to a rheumatologist in the first year after diagnosis, PM2) percentage of RA patients under the care of a rheumatologist seen in follow-up at least once per year, PM3) percentage of RA patients dispensed a disease modifying anti-rheumatic drug (DMARD) during the measurement year, PM4) time to DMARD therapy in new onset RA.

Methods: All patients with RA visits between 01/01/1997 and 31/12/2009 in BC were identified using health administrative data and followed until December 2014. Cases were identified by ≥ 2 physician billing codes for RA ≥ 8 weeks but ≤ 5 years apart. For this study, only cases age >18 who were seen by a rheumatologist at some point over follow-up were included. PM1: The percentage of incident RA cases with at least one visit to a rheumatologist within one year of their first RA visit was evaluated. PM2: The proportion of prevalent RA cases having at least one visit per year was calculated for those under rheumatology care. PM3: The percentage of prevalent RA patients dispensed a DMARD (including biologic agents and small molecule inhibitors) was calculated. PM4: time from RA onset (defined as first RA visit) to DMARD therapy was reported (in the calendar year of RA incidence), using median and 90th percentile wait time, as well as the proportion meeting the benchmark of 14 days.

Results: The cohort included 18 976 incident and 29 639 prevalent RA cases. The percentage of patients seen by a rheumatologist within 1 year from RA diagnosis increased over time, while the percentage of RA patients under the care of a rheumatologist seen in yearly follow-up declined steadily overtime. Further analysis (data not shown) suggests this is due to having more people with longer follow-up in the latter years, and lost to follow-up increasing over time, rather than a true

calendar year effect. The percentage of RA patients dispensed a DMARD was suboptimal (56% $>$ 65%). Of note, patients were not necessarily seen by a rheumatologist during the measurement year. The median time to DMARD improved over time to 23 days in 2009, with roughly one third receiving a DMARD within the benchmark of 14 days.

Conclusions: The present study represents the first time the AAC's PMs have been tested in administrative data and highlights where the measures are being met and potential gaps in care which require further examination.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5023

THU0647 LONGITUDINAL STUDY OF LONG-TERM POVERTY AND PERSISTENT DEPRESSIVE SYMPTOMS IN SLE

E. Yelin¹, J. Yazdany¹, L. Trupin¹, N. McCormick², P. Katz¹. ¹Medicine/Rheumatology, University of California, San Francisco, San Francisco, USA; ²Arthritis Research Centre of Canada, University of British Columbia, Vancouver, Canada

Background: A prior study found that persons with SLE in long-term poverty have greater accumulation of disease damage over 6 years than those exiting poverty or never in poverty¹. The present study evaluates the effect of long-term poverty status on depressive symptoms over the same duration of time.

Objectives: Analyse the impact of long-term poverty on prevalent and incident persistent depression after accounting for other risk factors for depression among persons with SLE.

Methods: Data are from the UCSF Lupus Outcomes Study in which persons with SLE were recruited in 2003 throughout the U.S. and interviewed annually through 2015. In each year we characterised respondents' poverty status based on household income and family size and administered the CESD measure of depressive symptoms, defining a high level of depressive symptoms using a validated SLE-specific cutpoint (≥ 24) associated with a formal diagnosis of depression². Prevalent persistent depression was defined as having high levels of depressive symptoms for ≥ 3 years between 2009 and 2015. Incident persistent depression used the same criteria, measured only among those who had low levels of depressive symptoms between 2006 and 2009. Logistic regression was used to estimate the impact of being poor in every year from 2003–2009, permanently leaving poverty by 2009, or never being poor on prevalent and incident persistent depression, with and without adjustment for gender, age, marital status, race/ethnicity, education, disease duration, extent of accumulated damage by 2009 using the Brief Index of Lupus Damage³, smoking status, and BMI.

Results: 535 persons with SLE were interviewed in each year from 2003 to 2015 (94% female, 65% non-Hispanic whites, mean age in 2003 50 years, range 20–83, mean disease duration 17 years, range 1–51). Between 2003 and 2009, 81% were never poor, 8% exited poverty, and 11% were poor in every year. 89 of the 535 (16.6%) met the study definition of prevalent persistent depression; 23 (7.4%) of the 312 free of high levels of depressive symptoms from 2006–2009 had incident persistent depression as of 2015. Table 1, below, indicates that those who were poor in every year had significantly higher rates of prevalent and incident persistent depression than those exiting poverty or never poor.

Abstract THU0646 – Table 1. Results of Four Performance Measures for RA.

Measurement Year ¹	Prevalent RA cases	Incident RA cases	PM1: Patients seeing rheumatologist within 1 year of RA onset	PM2: % of Patients under rheum care with yearly rheumatology visits	PM3: % Patients dispensed a DMARD	PM4: Median (90th percentile) days to DMARD	PM4: % with DMARD within 14 days of onset
2004	17 472	1647	80%	57%	59%	31 (825)	27%
2005	19 097	1614	82%	56%	60%	26 (579)	30%
2006	20 612	1704	85%	54%	60%	23 (411)	31%
2007	21 764	1430	83%	53%	60%	29 (399)	28%
2008	22 721	1305	86%	50%	60%	26 (339)	29%
2009	23 386	921	92%	49%	61%	23 (188)	34%
2010	23 122	-	-	46%	60%	-	-
2011	22 781	-	-	44%	56%	-	-
2012	22 370	-	-	42%	56%	-	-
2013	21 966	-	-	41%	57%	-	-
2014	21 412	-	-	41%	57%	-	-

¹PMs reported on a prevalent cohort until 2014, and on an incident cohort reported until 2009