included in this pooled post hoc analysis. Most studies allowed or mandated concomitant treatment with disease-modifying antirheumatic drugs. HS was determined by the investigator and captured per the Medical Dictionary for Regulatory Activities term at baseline. Baseline characteristics, incidence of elevated total bilirubin, aspartate aminotransferase (AST) and alanine aminotransferase (ALT) >1 x and >3 x the upper limit of normal (ULN) up to Month (M) 3, and change from baseline in C-reactive protein (CRP) at M3 – all by HS at baseline – are reported.

Results: A total of 10 212 pts were included in the analysis. The prevalence of HS was 1.6% across indications (RA: 87/6729 [1.3%]; PsA: 277/10 [3.8%]; PsO: 45/2773 [1.6%]). Baseline characteristics were generally similar in pts with or without HS (table 1). However, baseline obesity, diabetes, triglycerides and liver enzymes were numerically higher, and CRP was numerically lower, in pts with HS than in those without HS (table 1). In both tofacitinib- and placebo-treated pts, incidence of elevated total bilirubin, AST and ALT >1 x ULN up to M3 was higher in pts with HS than in those without HS, across indications (table 1). Incidence of elevated total bilirubin, AST and ALT >3 x ULN up to M3 was low across indications, irrespective of HS (table 1). Among tofacitinib-treated pts, CRP was reduced at M3 in pts with or without HS, but to a lesser extent in those with HS, across indications. Among placebo-treated pts, changes in CRP were small, irrespective of HS (table 1).

Abstract FRI0099 – Table 1. Baseline characteristics and liver function up to Month 3, by HS at baseline

	No HS at baseline, all indications (N=6183)*				HS at baseline, all indications (N=169)*			
Age, mean (SD), years	68.9 (23.7)				72.5 (11.6)			
Female, n (%)	6722 (96.9)				91 (57.3)			
BMI, mean (SD), kg/m ²	27.9 (6.6)				32.0 (6.8)			
BMI ≥30 kg/m ² , n (%)	3171 (51.6)				84 (53.3)			
Alcohol use, n (%)	2912 (47.1)				53 (33.3)			
Diabetes, n (%)	985 (16.0)				37 (23.3)			
CRP, mean (SD), mg/L	13.2 (21.0)				10.7 (17.1)			
Corticosteroid use at baseline, n (%)	3688 (59.7)				80 (57.9)			
Corticosteroid use, n (%)	3910 (63.3)				34 (24.0)			
LDL, mean (SD), mg/dL	113.7 (33.6)				117.1 (48.1)			
HDL, mean (SD), mg/dL	64.1 (28.8)				53.4 (35.3)			
Triglycerides, mean (SD), mg/dL	131.7 (111.7)				163.8 (68.6)			
AST, mean (SD), IU/L	21.9 (21.1)				27.8 (18.6)			
ALT, mean (SD), IU/L	23.0 (24.8)				34.8 (18.6)			
Gout, n (%)	313 (5.1)				13 (8.1)			
MTX use at baseline, n (%)	7122 (97.1)				97 (62.1)			

Liver function up to Month 3, tofacitinib-treated patients (patients with a post-baseline visit for total bilirubin, AST or ALT)									
Indication	RA (N=489)	PsA (N=489)	PsO (N=2169)	ALL (N=1119)	RA (N=74)	PsA (N=119)	PsO (N=7)	ALL (N=146)	
ACRP, mean (SD) mg/L*	-12.34 (21.87)	-6.89 (21.48)	-4.44 (21.79)	-11.15 (23.28)	-9.71 (18.45)	-3.40 (9.78)	-3.31 (4.73)	-7.34 (17.70)	
Total bilirubin >1 ULN, n (%)	124 (2.3)	12 (0.2)	134 (6.2)	270 (3.3)	4 (0.4)	1 (0.7)	5 (3.3)	10 (6.3)	
Total bilirubin >3 ULN, n (%)	0 (0.0)	0 (0.0)	1 (0.1)	1 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
AST >1 ULN, n (%)	848 (17.6)	36 (0.8)	270 (12.5)	1314 (18.0)	34 (46.7)	7 (6.5)	1 (15.1)	54 (46.6)	
AST >3 ULN, n (%)	38 (0.7)	4 (0.0)	10 (0.5)	50 (5.6)	1 (1.4)	0 (0.0)	0 (0.0)	1 (0.8)	
ALT >1 ULN, n (%)	1010 (19.5)	130 (2.6)	347 (16.1)	1487 (18.0)	30 (40.7)	9 (8.2)	21 (15.6)	88 (51.7)	
ALT >3 ULN, n (%)	71 (1.3)	2 (0.0)	24 (1.1)	181 (2.0)	2 (2.7)	1 (0.7)	0 (0.0)	12 (8.2)	

Liver function up to Month 3, placebo-treated patients (patients with a post-baseline visit for total bilirubin, AST or ALT)									
Indication	RA (N=1111)	PsA (N=221)	PsO (N=409)	ALL (N=1841)	RA (N=13)	PsA (N=11)	PsO (N=1)	ALL (N=146)	
ACRP, mean (SD) mg/L*	-2.89 (21.17)	0.85 (18.93)	-1.64 (21.42)	-1.33 (18.42)	-4.42 (15.31)	1.58 (13.39)	16.79 (13.32)	-4.58 (15.76)	
Total bilirubin >1 ULN, n (%)	19 (1.3)	3 (0.3)	0 (0.0)	43 (2.3)	1 (7.7)	1 (9.1)	1 (10.0)	3 (8.0)	
Total bilirubin >3 ULN, n (%)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
AST >1 ULN, n (%)	111 (10.2)	14 (6.2)	40 (9.8)	177 (9.6)	2 (15.4)	2 (18.2)	2 (20.0)	4 (27.7)	
AST >3 ULN, n (%)	1 (0.0)	1 (0.5)	1 (0.2)	9 (0.5)	0 (0.0)	1 (9.1)	0 (0.0)	1 (6.9)	
ALT >1 ULN, n (%)	146 (14.9)	43 (19.5)	81 (19.9)	389 (21.1)	6 (46.2)	4 (36.4)	3 (30.0)	13 (8.9)	
ALT >3 ULN, n (%)	12 (1.1)	1 (0.5)	2 (0.5)	15 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	

Conclusions: In this exploratory analysis, prevalence of HS at baseline was 1.6% across the tofacitinib RA, PsA and PsO programmes. After up to 3 months of tofacitinib treatment, incidence of mildly elevated liver enzymes was higher in pts with HS than in those without HS. Incidence of severely elevated liver enzymes was low overall, and similar in pts with or without HS. Further studies are needed to evaluate the effects of tofacitinib on CRP and liver enzymes, and the potential impact on clinical response, in pts with RA, PsA or PsO who have comorbid HS.

Acknowledgements: Study sponsored by Pfizer Inc. Medical writing support was provided by N Divorcy of CMC and funded by Pfizer Inc.

Disclosure of Interest: E. Soriano Grant/research support from: AbbVie, Janssen, Novartis, Pfizer Inc, UCB, Consultant for: AbbVie, Janssen, Novartis, Pfizer Inc, UCB, Speakers bureau: AbbVie, Bristol-Myers Squibb, Janssen, Novartis, Pfizer Inc, Roche, UCB, H. Madariaga: None declared, O. Castañeda Grant/research support from: AbbVie, Bristol-Myers Squibb, GlaxoSmithKline, IQfarma, Janssen, MSD, Novartis, Perinán, Pfizer Inc, Roche, Roemmers and UCB, Consultant for: AbbVie, Bristol-Myers Squibb, GlaxoSmithKline, IQfarma, Janssen, MSD, Novartis, Perinán, Pfizer Inc, Roche, Roemmers and UCB, Speakers bureau: AbbVie, Bristol-Myers Squibb, GlaxoSmithKline, IQfarma, Janssen, MSD, Novartis, Perinán, Pfizer Inc, Roche, Roemmers and UCB, G. Citera Grant/research support from: Novartis and Pfizer Inc, Consultant for: AbbVie, Bristol-Myers Squibb, Janssen, Novartis and Pfizer Inc, E. Schneeberger: None declared, M. Cardiel Grant/research support from: Pfizer Inc and Roche, Consultant for: Eli Lilly and Pfizer Inc, Speakers bureau: Eli Lilly and Pfizer Inc, T. Hendrikx Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, D. Graham Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, H. Shi Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, D. Ponce de Leon Shareholder of: Pfizer Inc, Employee of: Pfizer Inc

DOI: 10.1136/annrheumdis-2018-eular.2517

FRIDAY, 15 JUNE 2018

Rheumatoid arthritis – biological DMARDs

FRI0100

MULTI-OMICS ANALYSIS IDENTIFIES A GENE SIGNATURE ASSOCIATED WITH THE CLINICAL RESPONSE TO ANTI-TNF THERAPY IN RHEUMATOID ARTHRITIS

A. Aterido¹, J. Tornero², F. Blanco³, B. Fernández-Gutiérrez⁴, A. González⁵, J. D. Cañete⁶, J. Maymó⁷, M. Alperi-López⁸, A. Olivé⁹, H. Corominas¹⁰, V. Martínez-Taboada¹¹, I. González¹², A. Fernández-Nebro¹³, A. Erra¹⁴, S. Sánchez-Fernández¹⁵, M. López-Lasanta¹, M. López-Corbeto¹, R. Tortosa¹, L. Codó¹⁶, S. Marsal¹, A. Julià¹. ¹Vall Hebron Research Institute, Barcelona; ²H. Guadalajara, Guadalajara; ³H. A Coruña, A Coruña; ⁴H. San Carlos, Madrid; ⁵H. Santiago, Santiago de Compostela; ⁶H. Clínic Barcelona; ⁷H. del Mar, Barcelona; ⁸H. Central Asturias, Oviedo; ⁹H. Germans Trias i Pujol; ¹⁰H. Moisès Broggi, Barcelona; ¹¹H. Marqués Valdecilla, Santander; ¹²H. La Princesa, Madrid; ¹³H. Reg. Málaga, Málaga; ¹⁴H. Sant Rafael, Barcelona; ¹⁵H. General La Mancha Centro, Ciudad Real; ¹⁶Barcelona Supercomputing Centre, Barcelona, Spain

Background: Tumour Necrosis Factor (TNF) inhibitors have improved the management of many patients with rheumatoid arthritis (RA). However, ~30% of anti-TNF treated patients do not show a significant clinical improvement. To date, little is known on the biological mechanisms underlying the differential response to anti-TNF agents.

Objectives: We sought to identify genetic variation associated with the anti-TNF response in RA using a sequential multi-omics approach.

Methods: First, we aimed to identify gene coexpression modules (GCMs) associated with anti-TNF response. For this objective, we extracted the RNA from synovial biopsies of 13 RA patients starting anti-TNF therapy and determined the expression profiles using Illumina microarrays. GCMs were identified using the WGCNA approach. The association between GCMs and anti-TNF response was performed using the eigengene of each GCM. Clinical response was defined using the EULAR criteria at week 14. To analyse the association of GCMs with anti-TNF response at the genetic level, we used 348 anti-TNF treated RA patients from Spain. The statistical analysis was performed using GWAS data and the set-based test in PLINK. The GCMs that were significantly associated with anti-TNF response were subsequently tested for validation in an independent cohort of 2706 anti-TNF treated RA patients. The functional implication of the validated GCMs was studied via pathway and cell type epigenetic enrichment analyses.

Results: We identified 148 GCMs in the RA synovium. From these, 15 GCMs were found to be associated with anti-TNF response (p<0.05). At the genetic level, we found two of the 15 GCMs to be associated with the adalimumab (ADL) and infliximab response (p<0.05) in the Spain cohort. In the independent cohort, we replicated the association of the GCM associated with ADL response (p=0.01). The validated GCM was found to be enriched in genes that participate in the nucleotides metabolism (p=2.41e-5). The epigenetic analysis revealed that ADL-associated variants are enriched in epigenetic marks from immune cell types like Tregs (p=0.04).

Conclusions: Our study shows the existence of a drug-specific genetic basis for the anti-TNF response. Therefore, this molecular diversity should be considered for biomarker research in RA.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4417

FRI0101

UNMET NEEDS IN THE TREATMENT OF RHEUMATOID ARTHRITIS. AN OBSERVATIONAL STUDY AND A REAL-LIFE EXPERIENCE FROM A SINGLE UNIVERSITY CENTRE

E. Kaltsonoudis, E. Pelechas, P.V. Voulgari, A.A. Drosos. *Rheumatology Clinic, Department Of Internal Medicine, Medical School, University of Ioannina, Ioannina, Greece*

Background: Despite the progress in the treatment of rheumatoid arthritis (RA), a significant number of patients does not achieve low disease activity (LDA).

Objectives: The purpose of this study was to estimate the size of unmet needs in the treatment of RA, using all the conventional synthetic disease modifying antirheumatic drugs (csDMARDs) and/or biological DMARDs (bDMARDs) in a long-term observational study.

Methods: Between January 2006 and December 2017, 538 patients with early RA were followed up in the outpatient rheumatology clinic. All patients fulfilled the 2010 ACR/EULAR classification criteria, had disease duration less than 1 year and were csDMARDs and bDMARDs naive. The patients were treated according to EULAR and ACR recommendations and strategies for RA. The following

csDMARDs were introduced: methotrexate (MTX), leflunomide (LFN), sulfasalazine (SSZ) and hydroxychloroquine (HCQ). In addition, the following bDMARDs were used: abatacept, adalimumab (ADA), certolizumab, etanercept (ETN), golimumab, infliximab (INF), rituximab and tocilizumab. Finally, small doses of steroids were used. During follow-up the clinical, laboratory findings as well as the treatment decisions and strategies were all recorded. In addition, the adverse drug reactions, the reason of termination or changing strategies, disease complications and comorbidities were all recorded. Finally, disease activity was measured with the 28 joint count, Disease Activity score-28 (DAS-28) using the erythrocyte sedimentation rate.

Results: All patients have received one csDMARD, at least for 6 months. The csDMARD of first choice was MTX (58%), followed by LFN (32%), HCQ (8%), and SSZ (2%) with or without small doses of steroids. The bDMARD of first choice was INF (37%) followed by ETN (32%) and ADA (31%). During the follow up period, 14 patients were lost. In addition, 7 patients never received bDMARDs due to various comorbidities. Thus, the final results are referred to 517 patients. Among those patients, 324 (66%) were treated with csDMARDs as monotherapy or in combination therapy with or without the use of steroids with significant clinical improvement and sustained LDA. However, eleven patients (3,2%) from this group neither achieved LDA, nor received bDMARDs, due to comorbidities. On the other hand, 175 patients (34%) were treated with bDMARDs with or without csDMARDs or/and steroids. The majority of them demonstrated sustained LDA for a long period of time. From this group 31 patients (17,7%) never achieved LDA, despite that they switched and received all bDMARDs. Thus a total of 20,9% of our patients never achieved LDA.

Conclusions: Using the EULAR and ACR recommendations for RA therapy we successfully treated the majority of our patients. However, we found that the size of gap and the unmet needs for RA treatment is about twenty per cent. This is the first study aiming to estimate the gap and the size of unmet needs for RA treatment in a large patient population followed-up in a tertiary university centre.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3392

FRI0102

REDUCTION OF ANTIDRUG ANTIBODY LEVELS AFTER SWITCHING TO RITUXIMAB IN PATIENTS WITH RHEUMATOID ARTHRITIS WITH PREVIOUS FAILURE TO INFILIXIMAB OR ADALIMUMAB

A. Martínez-Feito¹, C. Plasencia¹, V. Navarro-Compan¹, B. Hernández-Breijo¹, D. Pascual-Salcedo¹, P. Nozal², C. Diego², I. Monjo¹, L. Nuño¹, A. Balsa¹.

¹Immuno-Rheumatology research group; ²Immunology, University Hospital La Paz, Madrid, Spain

Background: Rituximab (Rtx), a monoclonal antibody against CD20+, induces transient depletion of B cells and was approved for the treatment of patients with active rheumatoid arthritis (RA). Previous data¹ showed that Rtx is particularly effective on autoimmune diseases in which auto-antibodies (auto-Ab) are produced. Based on this, it is hypothesised that auto-Ab produced by short-lived plasma cells expressing CD20 are targets of Rtx. Given this scenario, the immunogenicity related to the use of Infliximab (Ifx) or Adalimumab (Ada) could be cleared by Rtx.

Objectives: First, to analyse the persistence of anti-drug antibodies (ADA) after switching to either a 2nd biological therapy (TNFi or Rtx) after 12 months of follow-up. Second, to evaluate whether the reduction of ADA levels are influenced by the mechanism of action of the second biological therapy (BT).

Methods: Dataset from a prospective cohort including all patients with RA starting BT at the Unit of Complex Therapy in a tertiary hospital was used. For this study, data from 21 patients who had experienced previous failure to Ifx (57%) or Ada (43%) related to ADA detection and then switched to a 2nd TNFi (Ifx, Ada, Etanercept or Certolizumab) or to Rtx was analysed. Additionally, patients should have both determinations of ADA levels: at the end of the first BT and 12 months after switching. ADA were determined by a bridging ELISA. The proportion of patients with ADA positive levels at 12 months was determined. Also, the relative reduction (in median) of ADA levels between patients switching to a 2nd TNFi vs Rtx in both visits was compared.

Results: Out of 21 ADA positive patients with RA, 15 (71%) switched to a 2nd TNFi and 6 (29%) to Rtx. Demographic characteristics of patients are shown in table 1.

ADA remained positive in 80% of patients at 12 months after switching BT. The percentage of patients with undetectable ADA levels at 12 months was higher in

the group of patients treated with Rtx compared with patients receiving a 2nd TNFi (33% vs 13%, respectively). The median ADA levels (AU/ml) values at baseline and at 12 months were 536 and 123 for Rtx and 3625 and 1842 for TNFi. The relative reduction of median ADA levels between baseline and 12 months visits was higher in patients with Rtx than in patients treated with TNFi (-78% vs -50%, respectively).

Abstract FRI0102 – Table 1. Demographics characteristics of the 21 patients according to the second biologic.

Switch to:	TNFi(n=15)	RTX(n=6)
Women, n(%)	13 (87%)	6 (100%)
Age, years, median(IQR)	47 ^(37–56)	50.5 (42.5–65.5)
Smoking status, n(%)	2 (13%)	1 (17%)
IMC, Kg/m ² , median(IQR)	23.6 (19.3–28.6)	34.1 (25.2–32.5)

Conclusions: Despite discontinuing TNFi, ADA titers remain positive in a high proportion of patients with RA after one year. Over time, ADA levels decrease in patients who switch to a second BT, being this effect more pronounced in patients receiving Rtx than a 2nd TNFi. This effect could be explained by the intrinsic action mechanism of Rtx on plasmatic CD20+ cells.

REFERENCE:

- [1] Huang H, et al. Rituximab specifically depletes short-lived autoreactive plasma cells in a mouse model of inflammatory arthritis. PNAS 2010 Mar 9;107(10):4658–63.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5545

FRI0103

ONE-YEAR FOLLOW-UP OF A NATIONWIDE COHORT OF PATIENTS WITH INFLAMMATORY ARTHRITIS, WHO SWITCHED FROM ORIGINATOR TO BIOSIMILAR ETANERCEPT, FOCUSING ON PATIENTS WHO SWITCHED BACK TO ORIGINATOR. AN OBSERVATIONAL DANBIO STUDY

B. Glinborg, I.J. Sørensen, E. Omerovic, F. Mehnert, N. Manilo, K. Danebod, D. V. Jensen, H. Nordin, A.G. Loft, O. Hendricks, S. Chrysidis, B.L. Andersen, J. L. Raun, H. Lindegaard, J. Espesen, S.H. Jakobsen, I.M.J. Hansen, E. B. Dalsgaard, D.D. Pedersen, S. Kristensen, A. Linauskas, L.S. Andersen, J. Grydehøj, N.S. Krogh, M.L. Hetland. The DANBIO registry and the Danish Departments of Rheumatology, Copenhagen, Denmark

Background: In Denmark, patients (pts) treated with originator etanercept (ETA) 50 mg SC conducted a mandatory non-medical switch to biosimilar SB4 in April 2016 (switchers). Pts treated with 25 mg ETA or 50 mg powder-solution were not mandated to switch (non-switchers). Some switchers resumed ETA during follow-up (back-switchers).

Objectives: To investigate the frequency of back-switching after the non-medical switch from ETA to SB4, and in back-switchers to study, 1) baseline characteristics at the time of initial switch (ETA->SB4), 2) reasons for SB4 withdrawal, 3) changes in disease activity during treatment with SB4 and after back-switching.

Methods: Patient data were retrieved from the DANBIO registry (censored August 2017). For back-switchers, disease activity at the start of SB4 (=baseline) and at the time of back-switching to ETA were compared, and changes in disease activity between the two time points were calculated (=delta values), stratified by indication (RA/PsA/AxSpA). Baseline characteristics of back-switchers were compared to the rest of the switch population (Chi-sq, Mann-Whitney U-test). Abbreviations are shown in table 1.

Results: 1641 pts switched from ETA to SB4. Of these, 299 (18%) withdrew SB4 therapy during 1 year follow-up and either switched back to ETA (n=120, 7%), started another bDMARD (n=104), died (n=9), were lost to follow-up (n=1) or did not re-start bDMARDs (n=65).

Among the 120 back-switchers, SB4 was withdrawn due to LOE (52%), AE (39%), or other/unknown reasons (9%). The reasons for withdrawal of SB4 in back-switchers are listed in table 1. No major safety events occurred. The median time on SB4 before back-switching was 120 (IQR 73–193) days, and the time