has improved the possibility of controlling disease activity and slowing the progression of joint damage. However their impact on work participation is unclear.

Objectives: To assess the effect of biological and tsDMARDs versus conventional treatments in patients with CIAs on work outcomes: employment, presenteeism and absenteeism.

Methods: A systematic review of the literature using Pubmed-Medline and the Cochrane library was performed until January 2017. All randomised controlled trials (RCT) and controlled cohorts (CC) comparing work outcomes in patients with rheumatic diseases such as rheumatic arthritis (RA), ankylosing spondylarthritis (AS) and psoriatic arthritis (PsA) treated with biological or tsDMARDs versus conventional therapies were selected. Statistical analysis determined in each study effect size (ES) or odds-ratios (OR) as appropriate to assess the magnitude of treatment effect. Pooled ES and OR were computed by meta-analysis. A random effect model was used in case of heterogeneity.

Results: Thirty six RCTs and eight CCs were analysed ie 12 769 patients with conventional treatment and 19 875 patients with bDMARD or tsDMARD (4619 Infliximab, 4629 Etanercept, 3872 Adalimumab, 670 Golimumab, 2101 Certolizumab, 691 Abatacept, 444 Sirukumab, 1668 Baricitinib, 672 Tofacitinib, 365 Sarilumab, 444 Sirukumab etc); 34 studies included 30 423 patients with RA, 7 studies included 1496 patients with AS and 3 studies included 725 patients with PsA.

This meta-analysis showed in patients treated by bDMARD vs conventional treatment:

- a significant decrease of accumulated missed workdays at week 24: ES -0.34 IC95%[-0.6; -0.08] and at week 52: ES -0.04 IC95%[-0.29; 0.2],

- a significant decrease of patients losing hours due to CIAs: RR 0.63 IC95% [0.48; 0.83],

- a significant improvement in VAS productivity: ES -1.81 IC95%[-2.61; -1.01],

- For the employment loss, the positive effect of bDMARDs was nearly significant: OR 0.60 IC95% [0.33; 1.09].

Conclusions: Despite the heterogeneity of the data, this meta-analysis showed the beneficial effect of bDMARDs on both absenteeism and presenteeism in CIAs. Thus the high cost of biologic agents could be partly balanced with savings in indirect costs.

Disclosure of Interest: C. Traverson: None declared, A. tubery: None declared, C. Hua Consultant for: Pfizer, BMS, abbvie, F. Barchechath-Flaisler Consultant for: Roche Pharmaceuticals, C. Lukas Consultant for: Abbvie, BMS, Celgene, Janssen, MSD, Novartis, Pfizer, Sanofi, Schering, Roche- Chugai, UCB, B. Combe Grant/research support from: Pfizer, UCB, Consultant for: Abbvie, BMS, Janssen, Lilly, MSD, Novartis, Pfizer, Roche-Chugai, UCB, Speakers bureau: BMS, Janssen, Lilly, MSD, Pfizer, Roche-Chugai, UCB, J. Morel Consultant for: Abbvie, BMS, Celgene, Janssen, Medac, MSD, Novartis, Pfizer, Sanofi, Schering, Roche- Chugai, UCB, C. Gaujoux-Viala Grant/research support from: Pfizer, Consultant for: Abbvie, BMS, Celgene, Janssen, Medac, MSD, Nordic Pharma, Novartis, Pfizer, Sanofi, Roche- Chugai, UCB

DOI: 10.1136/annrheumdis-2018-eular.4839

FRI0107 FIVE SUCCESSFUL PREGNANCIES WITH ANTENATAL ANAKINRA EXPOSURE

C.J.F. Smith¹, C. Chambers². ¹Rheumatology, Allergy, Immunology; ²Pediatrics, University of California San Diego, La Jolla, USA

Background: Current recommendations are to discontinue the interleukin-1 (IL-1) inhibitor anakinra prior to pregnancy given lack of safety evidence.¹ A total of 39 previous exposed pregnancies have been documented in the literature, and two resulting cases of fetal renal agenesis have been described.^{2,3}

Objectives: Our aim is to add to the limited existing data on IL-1 inhibitor use in pregnancy.

Methods: Data were obtained from the Organisation of Teratology Information Specialists (OTIS) Autoimmune Disease in Pregnancy Project, a prospective cohort study of pregnancy outcomes in the U.S. and Canada. Eligible women enrolled prior to 19 weeks' gestation between 2004 and 2017. Women who consented to participate were interviewed two to three times during pregnancy with a standard questionnaire regarding medical history, exposures during pregnancy, and demographic characteristics. Outcomes were obtained by maternal interview and medical record abstraction.

Results: Five pregnancies with anakinra exposure were identified, all resulting in full-term singleton live births, mean gestational age at delivery of 38.9 weeks, with no major or long-term infant complications. Three maternal subjects used anakinra for adult-onset Still's disease (AOSD) and two for juvenile idiopathic arthritis (JIA). One mother was diagnosed with AOSD during pregnancy, and anakinra was started at 20 weeks' gestation. All other subjects had a known diagnosis prior to pregnancy, and four subjects used the medication into the third trimester. All maternal subjects used 100 mg dosing of anakinra, four with daily use, one with weekly use. For all subjects who discontinued anakinra, some amount of steroid medication was necessary for treatment of disease flare. Two subjects developed oligohydramnios, one also with pregnancy-induced hypertension. Two women

had caesarian sections, one medically-indicated and one scheduled. One infant had low birth weight, but follow-up records indicated normal adjusted weight at one year. Three women successfully breastfed their infants, at least two of whom continued anakinra while breastfeeding.

Conclusions: Anakinra was used successfully in five full term pregnancies. Two subjects developed oligohydramnios, however. With the two previous reported cases of fetal renal agenesis, a process leading to low fluid levels, and the fact that maternal hyperthermia has been previously linked to fetal renal anomalies,⁴ it is worth further exploring the potential link between anakinra use, uncontrolled maternal febrile disease, and fetal outcomes. These data add to the limited knowledge regarding antenatal anakinra use and support its use in pregnant women without access to effective alternative therapies that have a larger volume of reassuring safety data.

REFERENCES:

- [1] Götestam Skorpen C, et al. The EULAR points to consider for use of anti-rheumatic drugs before pregnancy, and during pregnancy and lactation. *Ann Rheum Dis* 2016;75:795–810.
- [2] Chang Z, et al. Anakinra use during pregnancy in patients with cryopyrin-associated periodic syndromes (CAPS). *Arthritis Rheumatol* 2014;66:3227–32.
- [3] Youngstein T, et al. International multi-centre study of pregnancy outcomes with interleukin-1 inhibitors. *Rheumatology (Oxford)*. 2017;56:2102–2108.
- [4] Abe K, et al. Maternal febrile illnesses, medication use, and the risk of congenital renal anomalies. *Birth Defects Res A Clin Mol Teratol* 2003;67:911–8.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.1849

FRI0108 GOLIMUMAB IMPROVES WORK PRODUCTIVITY AND ACTIVITY AND QUALITY OF LIFE IN PATIENTS WITH RHEUMATOID ARTHRITIS (RA), PSORIASIS ARTHRITIS (PSA) AND AXIAL SPONDYLOARTHRITIS (AXSPA): INTERIM RESULTS FROM A NON-INTERVENTIONAL STUDY IN AUSTRIA (GO ACTIVE)

C. Dejaco¹, T. Mueller², O. Zamani³, U. Kurtz⁴, S. Egger⁵, J. Resch-Passini⁶, A. Totzauer⁷, W. Eisterer⁸, B. Yazdani-Biuki⁹, T. Schwingenschloegl¹⁰, P. Peichl¹¹, A. Kraus¹², G.W. Naer¹³, V. Rickett¹³. ¹Medical University Graz; ²Private Office, Graz; ³Rheuma Zentrum Favoriten, Vienna; ⁴Rheuma Praxis Dr. Kurtz, Gleisdorf; ⁵Ordination Gesundheitsquadrat; ⁶Rheuma-Zentrum Wien-Oberlaa GmbH, Vienna; ⁷Private Office Dr. Anna Totzauer, Gmünd; ⁸Klinikum-klagenfurt am Wörthersee, Klagenfurt; ⁹Dr. Schrenk and Dr. Yazdani-Biuki OG -Gruppenpraxis, Fürstenfeld; ¹⁰Private Office Dr. Thomas Schwingenschloegl, Wiener Neudorf; ¹¹Private Office Prim. Univ. Doz. Dr. Peter Peichl, Vienna; ¹²Private Office Dr. Angelika Kraus, Melk; ¹³Merck Sharp and Dohme GmbH, Ares Tower, Donau-City-Straße 11, 1220 Vienna, AUSTRIA, Vienna, Austria

Background: Golimumab has shown clinical efficacy and tolerability within its clinical trial program. No systemic outcome data regarding patient-reported outcomes and health economic parameters reflecting the real world use of golimumab in Austria are currently available.

Methods: Go Active is a prospective, non-interventional, multi-centre study in Austria. The impact of golimumab therapy on work productivity and activity (WPAI) and quality of life (RAQoL for RA patients, AsQoL for axSpA patients, PsAQoL for PsA patients) is assessed by using patient reported outcomes. Patients (target recruitment: n=320) are followed up to 2 years. For the current interim analysis (data cut-off: 18DEC2017) changes in the primary endpoint from baseline to month 3 were analysed in 167 patients.

Results: 167 patients are included in the current analysis (74 patients with RA, 49 patients with axSpA, and 44 patients with PsA). At study entry, most patients were biological-naïve and employed.

Median age at registration was 52 years (patients with RA: 57 years, patients with axSpA: 41 years, and patients with PsA: 44 years). Almost 2/3 of patients were female (84% of patients with RA, 37% patients with axSpA, and 55% of patients with PsA). Most patients were biological-naïve at study entry (77% of all patients, 73% of patients with RA, 80% of patients with axSpA, and 82% of patients with PsA). 42% of patients were not employed (58% of patients with RA, 29% of patients with axSpA, and 30% of patients with SpA); 14% due to incapacity for work (12% of patients with RA, 21% of patients with axSpA, and 16% of patients with SpA) and 54% due to age-related pension (60% of patients with RA, 21% of patients with axSpA, and 69% of patients with SpA). Most of the patients, who worked for pay, worked full time. 159 of all patients and 66 of employed patients completed the WPAI questionnaire at baseline and after 3 months. Overall work productivity improved by -33 (-40 for patients with RA, and -31 for patients with axSpA and PsA) and activity impairment by -30 (-40 for patients with RA and