Conclusions: Biomarkers are able to improve prediction of radiographic spinal progression in axSpA, especially if used in addition to the clinical parameters, but the added value seems to be rather small.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3795

FRI0155

GENES REGULATING BONE HOMEOSTASIS ARE DIFFERENTIALLY EXPRESSED AT PERIPHERAL VERSUS AXIAL ENTHELIAL SITES IN AN ANIMAL MODEL OF SPONDYLOARTHRITIS

Z. Stavre, Y. Maeda, T. Huang, E. Gravalles. *Department of Medicine-Rheumatology, University of Massachusetts Medical School, Worcester, USA*

Background: In spondyloarthritis (SpA), mechanisms of enthesial bone formation are not well understood. Currently, no treatment can reverse this process, although TNF inhibitors may modify bone formation if initiated early. SKG mice exhibit features of SpA with both peripheral joint and axial inflammation after induction by curdlan via the Dectin-1 pathway residing upstream of TNF α , IL-23 and IL-17.

Objectives: The goal of this study was to identify genes and pathways that regulate enthesial bone formation. SKG mice were previously reported to develop axial bone formation within 12 weeks of curdlan injection, as well as enthesial bone formation around peripheral joints¹.

Methods: 9 week-old SKG or control BALB/c mice were injected IP with curdlan. Histology and microCT scanning were performed at serial time points. Laser capture microscopy was performed on formalin-fixed paraffin-embedded (FFPE) tissue sections from enthesial sites around the ankle and tail vertebrae 3 weeks after curdlan injection. RNA was extracted using the Qiagen RNeasy FFPE kit and whole transcriptome analysis was performed using Affymetrix gene array MTA 1.0.

Results: Inflammation at axial entheses developed earliest at the tail base. Contrary to previous reports, we did not observe axial bone formation, but rather saw significant erosion of vertebral bodies, throughout the spine. The absence of axial bone formation was noted as late as 24 weeks post curdlan. In contrast, exuberant enthesial bone formation occurred around ankles at predictable sites of tendon/ligament insertions into bone. We identified distinct expression of genes involved in bone resorption and formation at ankle when compared to spine enthesial sites. IPA upstream regulator analysis identified significant gene expression of upstream regulators of TGF- β , known to recruit early osteoblast precursors to bone (z-value 2.83), and WNT3A, a WNT signalling pathway agonist that promotes osteoblast differentiation and function (z-value 2.54), in the ankle, but not the spine. These findings support the histologic observation of bone formation at ankle, but not spine, enthesial sites. In addition, significant increased gene expression of upstream regulators of the aryl hydrocarbon receptor (AhR) (z-value 2.14) was found at spine, but not ankle, enthesial sites (z-value -1.78). AhR is a ligand-activated transcription factor that is a positive regulator of osteoclastogenesis via c-fos-mediated RANKL signalling.

Conclusions: Whole transcriptome analysis revealed increased expression of upstream regulators of genes in anabolic pathways, including WNT3A and TGF- β , at ankle enthesial sites where bone formation occurs, but not at spine enthesial sites where bone erosions persist without bone formation. These pathways may thus contribute to peripheral enthesial bone formation. Significantly increased gene expression of upstream regulators of AhR, a positive regulator of osteoclastogenesis, was found at spine enthesial sites. AhR is thus a candidate regulator of continued bone resorption at spine enthesial sites in this murine model of SpA.

REFERENCE:

- [1] Ruutu M, et al. β -Glucan Triggers Spondylarthritis and Crohn's Disease-like Ileitis in SKG Mice. *Arthritis Rheum* 2012 Jul;64(7):2211.

Acknowledgements: This work was supported by a grant provided by AbbVie.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.2973

FRI0156

AN ANTIMICROBIAL MECHANISM, NETOSIS, IS POTENTIALLY INVOLVED IN ANKYLOSING SPONDYLITIS PATHOGENESIS

P. Ruiz-Limon¹, M.C. Castro¹, C. Lopez-Pedraza¹, M.C. Abalos-Aguilera¹, N. Barroja¹, C. Perez-Sanchez¹, I. Arias-de la Rosa¹, R. Ortega¹, P. Font¹, J. A. Gonzalez-Reyes², J.M. Villalba², C. Lopez-Medina¹, A. Escudero¹, E. Collantes¹, Y. Jimenez-Gomez¹. ¹Rheumatology Service, IMIBIC/Reina Sofia University Hospital/Cordoba University; ²Dept. Cell Biology, Physiology and Immunology, Cordoba University; ceiA3, Cordoba, Spain

Background: Ankylosing spondylitis (AS) is a chronic inflammatory disease, of unknown etiology, that mainly affects the axial skeleton and the sacroiliac joints in the pelvis. Neutrophils are critical actors in innate immunity and their activation has been involved in the progression of AS symptoms. Recently, NETosis has been suggested to play a central role in several pathologic states, including rheumatology diseases. Nevertheless, this process has not been described yet in AS patients

Objectives: 1) To evaluate and characterise the presence of NETosis in AS patients. 2) To explore the relationship among NETosis markers and clinical characteristics of this disease

Methods: Thirty patients with AS and 32 healthy donors (HD) were included in the study. Disease activity was determined by BASDAI index and, CRP and ESR levels; in parallel, inflammatory markers were determined in plasma by ELISA kits. Spinal mobility of AS patients was measured by the BASMI index, and structural damage was calculated by the mSASSS index. *Ex vivo*, spontaneous NETosis generation in purified neutrophils from AS patients and HDs were measured by fluorescence (n=6) and scanning electron (n=3) microscopy after 6 hour of incubation. PMA, known to promote NETosis, was used as positive control. DNA extrusion was analysed by fluorescence microscopy and fluorimetry after SYTOX staining, whereas elastase percentage (NE) was analysed by fluorescence microscopy after staining of neutrophils with NE antibody. *In vivo*, myeloperoxidase (MPO) and NE protein expression were measured by flow cytometry (FACS Calibur), whereas extracellular DNA was examined in plasma using fluorimetry after SYTOX staining

Results: Compared to HDs, AS neutrophils showed spontaneous extracellular release of a meshwork of DNA nuclear and granule proteins (NETs), as demonstrated by fluorescence microscopy, fluorimetry, and scanning electron microscopy. Indeed, analysis of DNA fibres staining by SYTOX revealed that NETosis rate was above baseline levels after 6 hour of *ex vivo* AS neutrophil incubation as compared to those from HDs (p<0.05). Furthermore, higher cell free-DNA levels were observed between AS patients and HDs at 6 hour (p<0.05). Increased spontaneous NETs production in this pathology was additionally corroborated by the observation of an enhanced percentage of NE-staining cells after 6 hour of incubation (p<0.05). *In vivo*, higher intracellular NE levels (p=0.036), and plasma DNA concentrations (p=0.021) were also found in AS patients as compared to HDs. Correlation studies showed that plasma DNA levels positively correlated with inflammatory markers (i.e. CRP, ESR and TNF α), and with the spinal mobility index BASMI. In addition, a positive correlation was found between intracellular NE levels and plasma IL-1 β concentration

Conclusions: 1) NETosis is increased in AS patients. 2) Raised NETosis in AS is associated with several markers of inflammation and mobility. Thus, NETosis might act as a key mediator in the etiopathogenesis of AS. Funded by JA PI-0314-2012, SER, ISCIII (RIER RD16/0012/0015)

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4291

FRI0157

ALTERED EXPRESSION OF THE SPLICEOSOME COMPONENTS IN LEUKOCYTES SUBSETS FROM PATIENTS WITH ANKYLOSING SPONDYLITIS: ASSOCIATION TO DISEASE PATHOGENESIS AND THERAPEUTIC RESPONSE

L. Ladehesa¹, M.C. Castro¹, C. Lopez-Pedraza¹, R. Ortega¹, S. Pedraza², M. del Rio², M.C. Abalos-Aguilera¹, P. Ruiz-Limon¹, C. Perez-Sanchez¹, A. Ibañez-Costa¹, Y. Parra-Manso¹, N. Barroja¹, I. Arias-de la Rosa¹, P. Font¹, A. Escudero¹, J.P. Castaño², R.M. Luque², E. Collantes¹, Y. Jimenez-Gomez¹. ¹Rheumatology Service, IMIBIC/Reina Sofia University Hospital/Cordoba University; ²Dept. Biología Celular, Fisiología e Inmunología, IMIBIC/Universidad de Córdoba/Hospital Reina Sofia; CIBERObn y ceiA3, Cordoba, Spain

Background: Ankylosing spondylitis (AS) is a chronic inflammatory disease for which the etiopathogenesis has not yet been fully clarified. Splicing is a post-