Disclosure of Interest: K. Zobbe: None declared, D. Prieto-Alhambra Grant/research support from: Amgen, Servier, and UCB, Consultant for: UCB, Speakers bureau: Amgen, R. Cordtz: None declared, L. Mellemkjær: None declared, P. Højgaard: None declared, L. E. Kristensen Grant/research support from: UCB, Biogen, Janssen pharmaceuticals, and Novartis, Speakers bureau: Pfizer, AbbVie, Amgen, UCB, BMS, Biogen, MSD, Novartis, Eli Lilly and Company, and Janssen pharmaceuticals, L. Dreyer: None declared
DOI: 10.1136/annrheumdis-2018-eular.2302

FRI0222

ASSOCIATION BETWEEN BLOOD LEAD LEVELS AND GOUT: THE UNITED STATES NATIONAL HEALTH AND NUTRITION EXAMINATION SURVEY 2007–2014

T.T. Cheung, M.F. Tsoi, B.M.Y. Cheung, W.C.S. Lau. *Medicine, The University of Hong Kong, Hong Kong, Hong Kong*

Background: Lead is a heavy metal with no physiological role in humans. It is well known that excessive exposure to lead is associated with cardiovascular, gastrointestinal and neurological complications. However, recent studies have demonstrated that lead can increase the tubular reabsorption of urate, resulting in hyperuricaemia and the development of gout.

Although blood lead levels are dropping in the US general population, the association between low blood lead levels and gout remains inconclusive.

Objectives: To evaluate the relationship between serum blood lead levels and the development of gout in the US general population.

Methods: Adult participants with blood lead measurements and self reported gout in NHANES 2007–2014 were included in the analysis.

Results were analysed using SPSS complex sample module version 22. Logistic regression with sample weight adjustment was used to study the association between blood lead levels and gout. Odds ratio (OR) and 95% confidence interval (95% CI) were estimated. Sub-group analysis was conducted in participants with blood lead level <5 µg/dL.

Results: 18837 adult participants were included in the analysis. 18 270 participants had blood lead levels below 5 µg/dL (The current reference range for acceptable blood lead levels in healthy persons without excessive exposure to environment source of lead).

The prevalence of gout increased with blood lead levels. In the total population, the odd ratio of gout was 4.96 in subjects with blood lead levels above 2.21 µg/dL. After adjusting for body mass index, hypertension, renal function and use of diuretics, the odd ratio of gout was 2.65 which remained significant.

The subgroup analysis of subjects with blood lead levels below 5 µg/ml showed similar findings. The risk of gout is increased when blood lead levels are higher than 1.43 µg/dL.

Abstract FRI0222 – Figure 1. (A) Prevalence of gout according to quintiles of blood level. (B) Association between quintiles of blood lead level and gout at blood lead level <5µg/dL

Conclusions: Low blood lead levels are associated with gout. The risk of gout is increased when the blood lead level is higher than 1.43 µg/dL. Therefore, measures should be taken to minimise the environmental exposure to lead.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7067

FRI0223

IS PERFORMANCE OF THE 2015 ACR/EULAR GOUT CLASSIFICATION CRITERIA AFFECTED BY DISEASE DURATION AND GENDER? RESULTS FROM AN ITALIAN MULTICENTRE STUDY FOCUSED ON THE MANAGEMENT OF CRYSTAL-INDUCED ARTHRITIDES (ATTACK STUDY)

N. Ughi¹, A. Zanetti¹, P. Frallonardo², A. Hoxha³, M. Lorenzin³, A. Ariani⁴, F. Ceccarelli⁵, M.A. Cimmino⁶, C. Scirocco⁷, B. Raffener⁷, A. Bortoluzzi⁸, A. Di Matteo⁹, F. Furini⁸, M. Manara¹, F. Salaffi⁹, G. Carrara¹, C.A. Scire¹, R. Ramonda³.
¹Epidemiology Unit, Italian Society for Rheumatology, Milan; ²Università degli Studi di Padova; ³Università degli Studi di Padova, Padova; ⁴AOU di Parma, Parma; ⁵Sapienza Università di Roma, Roma; ⁶Università di Genova, Genova; ⁷Ospedale di Bolzano, Bolzano; ⁸AOU Sant'Anna, Ferrara; ⁹Università Politecnica delle Marche, Ancona, Italy

Background: Gout is a common form of arthritis, but early disease and phenotypes related to gender¹ may favour diagnosis misclassification, unless monosodium urate (MSU) crystals are searched for in synovial fluid (SF). The 2015 ACR/EULAR gout classification criteria² were validated in Rheumatology and primary settings,³ and the impact of disease features on performance were observed,⁴ yet little studied.

Objectives: To assess the performance of the 2015 ACR/EULAR gout criteria in a cohort of crystal-induced arthritides stratified by disease duration and gender in a real-life setting in Italy.

Methods: In/outpatients referring to Rheumatology Units for acute arthritis were enrolled into an ongoing multicentre cohort study designed for achieving improvement in the management of crystal-induced arthritis (ATTACK) by the Italian Society for Rheumatology. Gout was defined as MSU +SF (gold standard), irrespective of the clinical diagnosis, and the ACR/EULAR criteria were applied as full (all domains) and clinical-only set excluding imaging and SF analysis. To classify a patient as having clinical gout, the sufficient criterion (MSU +SF) was ignored and cutoff score ≥8 was used. Overall sensitivity and specificity were calculated. Disease duration (early if <2 years) was considered and gender effect explored.

Results: Of 199 adult patients (65±12 years, 75% male, 96% Caucasian) recruited in 11 hospitals, complete SF analysis and ACR/EULAR score were obtained in 105 (53%). MSU+ were 62/105 (59%), 47/62 (76%) were non-tophaceous and 25 (40%) early diseases. Calcium pyrophosphate (CPP) was the most prevalent crystal in non-MSU+. Details on performance (95% confidence intervals) are shown in table and on a receiver operating characteristic plot (figure 1).

Abstract FRI0223 – Table 1

	Sensitivity%		Specificity%	
	Full	Clinical-only	Full	Clinical-only
Overall n=105	76 (63–86)	71 (58–82)	95 (86–99)	81 (67–92)
<2 years n=49	48 (28–69)	40 (21–61)	100 (86–100)	92 (73–99)
≥2 years n=56	95 (82–99)	92 (78–98)	89 (67–99)	68 (44–87)
Male n=74	77 (64–87)	74 (60–85)	94 (71–100)	65 (38–86)
Female n=31	60 (15–95)	40 (5–85)	96 (80–100)	92 (75–99)

Overall good performance and high specificity of full and clinical-only sets were observed. In early disease criteria had high specificity, but low sensitivity, while performance was stable in longstanding gout. Specificity (clinical-only) and sensitivity were low when criteria were applied to males and females, respectively.