Results were analysed using SPSS complex sample module version 22. Multiple regression was used to analyse the difference in prevalence of gout and utilisation of urate lowering agents.

Results: 23482 adults participants were included in the analysis. There was no further increase in the prevalence of gout in the general US population. The prevalence was 3.82% and 3.84% in NHANES 2007–2010 and 2011–2016, respectively ($p=0.966$).

Although there is a decreasing trend in the serum urate levels among patients with gout, the utilisation of urate lowering agents remained low. Only 28.2% and 29.4% of patients with gout were prescribed urate lowering agents in 2007–2010 and 2011–2014, respectively.

Among patients with gout, history of chronic kidney disease and use of thiazide diuretics are the most significant negative predictors for achieving the therapeutic target of $\text{SUV} < 6 \text{ mg/dL}$ (Odds ratio of 0.23 and 0.41, respectively). Use of aspirin was not a significant predictor for treatment failure.

NHANES survey cycle	2007–2010	2009–2010	2011–2012	2013–2014	2015–2016	P-value
No. of participants included	5524	6215	5567	5735	5719	
Prevalence of gout (%)	3.9 [3.4–4.5]	3.7 [3.4–4.1]	3.8 [3.5–4.0]	4.3 [3.5–5.1]	3.9 [3.4–4.3]	0.392
Age and gender adjusted	3.9 [3.4–4.5]	3.9 [3.4–4.3]	3.8 [3.5–4.1]	4.3 [3.5–5.1]	3.9 [3.4–4.3]	

No. of patients with gout prescribed (%)	2007–2010	2009–2010	2011–2012	2013–2014	P-value
Urate-lowering therapy	27.1 [21.5–32.7]	25.8 [21.9–29.7]	26.1 [22.5–29.7]	32.5 [24.3–40.7]	0.008
Aspirin	28.7 [23.5–33.9]	27.8 [23.5–32.1]	24.1 [20.7–27.5]	29.5 [22.7–36.3]	0.006
Chronic kidney disease	0	0	0.6 [0.2–1.0]	2.6 [1.9–3.3]	0.009
Thiazide diuretic	18.0 [13.3–22.7]	12.1 [9.3–14.9]	9.7 [8.2–11.2]	14.0 [10.3–17.7]	0.019
Secondary prevention	8.78±0.29	5.65±0.17	5.50±0.24	6.43±0.11	0.009
Urate-lowering therapy success	5.25±0.22	5.22±0.17	5.59±0.27	6.63±0.17	0.001
Aspirin prescription	5.77±0.16	5.27±0.11	5.41±0.17	6.88±0.17	0.03
Thiazide diuretic prescription	7.10±0.29	5.80±0.25	7.67±0.39	7.79±0.51	0.576
Secondary prevention	0.55±0.22	0.49±0.24	0.64±0.14	0.57±0.23	0.039

	Univariate analysis	Multivariate analysis
Age	1 [reference]	1 [reference]
25–34	0.72 [0.37–1.35]	0.67 [0.35–1.29]
35–44	0.48 [0.28–0.82]	0.68 [0.44–1.05]
45–54	0.44 [0.22–0.91]	0.62 [0.40–0.95]
55–64	0.58 [0.36–0.93]	0.55 [0.35–0.86]
65–74	0.29 [0.14–0.61]	0.22 [0.11–0.43]
75–84	0.16 [0.08–0.31]	0.11 [0.06–0.19]
85–94	0.08 [0.04–0.15]	0.05 [0.03–0.09]
95–104	0.04 [0.02–0.08]	0.03 [0.02–0.05]
Gender	1 [reference]	1 [reference]
Male	1 [reference]	1 [reference]
Female	0.72 [0.37–1.35]	0.67 [0.35–1.29]
Ethnicity	1 [reference]	1 [reference]
White	1 [reference]	1 [reference]
Black	0.72 [0.37–1.35]	0.67 [0.35–1.29]
Hispanic	0.48 [0.28–0.82]	0.68 [0.44–1.05]
Other	0.44 [0.22–0.91]	0.62 [0.40–0.95]
Education	1 [reference]	1 [reference]
Less than high school	0.58 [0.36–0.93]	0.55 [0.35–0.86]
High school	0.29 [0.14–0.61]	0.22 [0.11–0.43]
Some college	0.16 [0.08–0.31]	0.11 [0.06–0.19]
College graduate	0.08 [0.04–0.15]	0.05 [0.03–0.09]
Postgraduate	0.04 [0.02–0.08]	0.03 [0.02–0.05]

Conclusions: The prevalence of gout in the US general population has not increased over the past 10 years. Although the use of urate lowering agents among patients with gout remained low, the percentage of patients with gout achieving the therapeutic target has increased over the past 10 years.

History of chronic kidney disease and use of thiazide diuretics are the most significant negative predictors for treatment success.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5636

FRI0252 TESTING FOR A CAUSAL ROLE OF MITOCHONDRIAL VARIATION IN THE DEVELOPMENT OF GOUT

T.R. Merriman¹, A. Shaikat¹, A. Gosling¹, M. Bixley¹, N. Dalbeth², L. Stamp³, E. Matisoo-Smith¹, J. Harre Hindmarsh⁴, L. Joosten⁵, T. Jansen⁶, M. Janssen⁶, A.-K. Tausche⁷, P. Riches⁸. ¹University of Otago, Dunedin; ²University of Auckland, Auckland; ³University of Otago, Christchurch; ⁴Ngati Porou Hauora Charitable Trust, Te Puia Springs, New Zealand; ⁵Radboud University Medical Centre, Nijmegen; ⁶VieCuri Medical Centre, Venlo, Netherlands; ⁷University of Dresden, Dresden, Germany; ⁸University of Edinburgh, Edinburgh, UK

Background: Mitochondria execute critical roles in diverse cellular pathways. As a danger signal mitochondria induce inflammation in response to stress through NLRP3 inflammasome activation, central to gout development. We recently reported association of reduced mtDNA copy number (CN) with prevalent gout in New Zealand Māori and Pacific (Polynesian) populations¹. However the cause-effect relationship is unknown. This could be evaluated by testing for association with gout using nuclear genetic variants that associate with mtDNA CN.

Objectives: 1) Genome wide association study (GWAS) for mtDNA CN to identify nuclear and mitochondrial loci controlling mtDNA copy number 2) test any such loci for association with gout.

Methods: The mtDNA CN GWAS comprised 1340 Eastern Polynesian (EP), 816 Western Polynesian (WP) and 4579 European samples (New Zealand, Germany, The Netherlands, Scotland) genotyped on the Illumina CoreExome v24 array. 343 mitochondrial single nucleotide polymorphisms (SNPs) were evaluated. As

previously described² the median of the absolute difference in X and Y probe intensities was used as a measure of mtDNA CN, and additional 10 000 randomly selected autosomal SNPs were used to calculate the principal components (PCs). A mtDNA CN GWAS was run on chromosomes 1–22 and the mitochondrial genome using Plink 1.9 .v2, adjusting for the first 10 PCs, age and sex followed by association analysis with gout adjusting by age, sex and the first 10 PCs generated from a separate set of 3000 autosomal SNPs.

Results: The association of reduced mtDNA CN with gout in the EP and WP groups was reproduced but there was no evidence of association of mtDNA CN with gout in Europeans. Two genome-wide significant ($p < 1 \times 10^{-7}$) variants *MUC17 rs78010183* (T-allele) and *SLC16A8 rs75640043* (T-allele) were associated with increased mitochondrial CN in EP and WP, respectively, and mitochondrial variant *rs3928306* was associated with mtDNA CN ($p = 4.49 \times 10^{-12}$) in Europeans. *MUC17 rs78010183* also associated with increased mtDNA CN in Europeans, with the T allele also increasing CN ($\beta = 0.06$, $p = 1.07 \times 10^{-4}$). The T-allele of *rs78010183* was associated with gout in Europeans (OR=9.32, $p = 5.53 \times 10^{-3}$) and the *SLC16A8 rs75640043* T-allele was associated with gout in the WP group (OR=6.85, $p = 5.50 \times 10^{-3}$). The mitochondrial variant *rs3928306* A-allele (very rare in Polynesian) was not associated with gout in Europeans (OR=1.09, $p=0.36$).

Conclusions: That genetic variants associated with mitochondrial copy number also associate with gout provides evidence for a potential causal role of mitochondrial copy number in gout. However, the nuclear genetic variants support a causal relation of increased mtDNA CN with gout, conflicting with our previous observational report of association of reduced mtDNA CN with gout¹.

REFERENCES:

- Gosling A, et al. Mitochondrial genetic variation and gout in Māori and Pacific people living in Aotearoa New Zealand. *Ann Rheum Dis* 2017 Dec 15. [Epub ahead of print].
- Asher FN, et al. Association of mitochondrial DNA levels with frailty and all-cause mortality. *J Mol Med* 2015;93:177–86.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5017

FRI0253 TRENDS FOR GOUT IN ADULTS IN AN URBAN AREA FOR A 5-YEAR PERIOD: INCIDENCE, PREVALENCE AND HOSPITALISATION RATES

V. Dostanko¹, V. Yagur¹, V. Apanasovich¹, N. Dostanko¹, A. Rekun². ¹2-d Department of Internal Medicine, BSMU; ²Rheumatology Department, 9-th Minsk City Clinical Hospital, Minsk, Belarus

Background: Gout is one of the most common arthritis nowadays which has a great influence on patient's quality of life, course and outcomes of cardiovascular and renal pathology.¹ Some recent papers demonstrate that the prevalence of gout has risen over the last decades and underline the scarcity or lack of epidemiologic data in different countries as well as the notable variations among them.^{2,3}

Objectives: We estimated incidence, prevalence and hospitalisation rates for gout in Minsk (the Republic of Belarus) for the 5 year period (2011–2015).

Methods: Minsk is a typical urban area which is considered to be representative for the urban population of the whole country. The data on the new onset gout and the first visit for gout in a corresponding year were collected from all rheumatologic services of Minsk for the 5 year period from January, 1 of 2011 to December, 31 of 2015. Only patients older than 18 years old with the diagnosis of gout according to the ICD-10 (M10) were included. The data on age and sex structure of the population of the Republic of Belarus were obtained from the annual Statistical bulletins of the National Statistical Committee of the Republic of Belarus. Hospitalisation rates were calculated on the base of statistical reports on discharges for the corresponding year.

Results: There were no significant differences in age and sex structure of the population in Belarus and Minsk for the study period. The size of the adult population, incidence and prevalence of gout, hospitalisation rates and duration of hospitalizations for gout for the corresponding years in Minsk are presented in the table 1. According to these data the gout incidence for the 5 year period changed from 25.0 (CI 95% 24.7–25.2) to 35.5 (CI 95% 35.3–35.8) per 100 000 adults, the gout prevalence for the period 2011–2015 also increased from 178.9 (CI 95% 178.3–179.5) to 290.3 (CI 95% 289.6–291.0) per 100 000 adults ($p < 0.001$ for both trends). There were 2688 hospitalizations in Minsk for gout for the study period. The mean age of hospitalised patients was 57.3 ± 10.3 years (median 58.0; range 25–87), 91.4% were men (92%; 87.8%–94.9%). It is worth noting that 45.6% of hospitalised patients were at the age 55–65% and 90.2% were under 70 years old with sharp decrease of hospitalizations after 65 years of age. Hospitalisation rates

also increased from 28.7 to 38.2 per 100 000 adults ($p<0.001$). The mean duration of hospitalisation was 12.5 ± 2.7 days (13 days; 2–26 days) and increased from 11.6 to 12.7 ($p=0.03$).

Abstract FRI0253 – Table 1

	2011	2012	2013	2014	2015
Population size in Minsk, adults	1537880	1555210	1568320	1585480	1598120
Incidence per 100 000	25.0	26.2	35.3	38.0	35.5
Prevalence per 100 000	179	228	248	255	290
Hospitalisation rate per 100 000	28.7	36.8	33.5	33.9	38.2
Hospitalisation duration, days	11.6	12.5	13.1	12.7	12.7

Conclusions: We revealed significant increase in incidence, prevalence as well as hospitalisation rates for gout in adults in Minsk (the Republic of Belarus) for the 5 year period (2011–2015) in 1.4, 1.6 and 1.3 times, correspondingly.

REFERENCES:

- [1] Kuo CF, Grainger MJ, Zhang W, Doherty M. Global epidemiology of gout: prevalence, incidence and risk factors. *Nat Rev Rheum* 2015;11(11):649–62.
- [2] Roddy E, Choi H. Epidemiology of Gout. *Rheum Dis Clin North Am*. 2014;40(2):155–175.
- [3] Dehlin M, et al. Incidence and prevalence of gout in Western Sweden. *Arthritis Res Ther* 2016;18:164.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5549

FRI0254

CLINICAL FEATURES AND RISK FACTORS FOR DEVELOPING GOUT ATTACKS DURING ANTI-TUBERCULOSIS MEDICATION: A CASE-CONTROL STUDY IN SOUTH KOREA

¹Y.-J. Ha, S.W. Chung², J. Lee³, E.H. Kang¹, Y.J. Lee¹, Y.W. Song³. ¹Division Of Rheumatology, Department Of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; ²Division Of Rheumatology, Department Of Internal Medicine, Incheon Sarang Hospital, Incheon; ³Division Of Rheumatology, Department Of Internal Medicine, Seoul National University Hospital, Seoul, Korea, Republic Of

Background: Gout is a form of inflammatory arthritis that develops in some people with hyperuricemia. Several drugs may induce hyperuricemia and gout attack. Some anti-tuberculosis (TB) medications are also known to increase uric acid level and trigger acute gout attack. However, there have been no well-described studies for gout during anti-TB therapy.

Objectives: To investigate the clinical features and risk factors of gout attacks following anti-TB treatment in South Korea.

Methods: We conducted the case-control study including 49 patients who developed gout attack following anti-TB medications and 147 age- and sex- matched TB controls without gout attack. Demographic, clinical features and laboratory findings of patients with gout attacks following anti-TB medication were analysed and compared to those of controls in order to establish risk factors associated with gout attack during anti-TB treatment.

Results: The mean age was 67.7 ± 13.2 years and 39 patients (79.6%) were male. Twenty-four patients (49.0%) had a previous history of gout. The mean time interval to develop gout attack from the initiation of anti-TB therapy was 4.13 ± 4.49 months. The attacks tended to involve lower extremity joints (43/49, 87.8%), especially the ankle and the first metatarsophalangeal joint. Factors associated with experiencing gout attack during anti-TB medication were higher body mass index (BMI, 23.7 ± 3.6 vs 21.8 ± 3.6 , $p=0.003$), higher pre-treatment uric acid level (7.2 ± 2.5 vs 6.2 ± 1.7 mg/dL, $p<0.001$), previous history of gout (49.0% vs 1.4%, $p<0.001$), absence of dyslipidemia (4.1% vs 16.3%, $p=0.008$), and decreased renal function (eGFR 63.6 ± 27.5 vs 82.6 ± 29.9 , $p=0.003$). In multivariate model, higher BMI, chronic kidney disease (CKD, eGFR <60 ml/min/1.73m²), previous history of gout, and pre-treatment hyperuricemia (uric acid ≥ 6.8 mg/dL) were

found to be independent risk factors for gout attack following anti-TB medication (table 1).

Abstract FRI0254 – Table 1. Multivariate analysis of risk factors for post-TB medication gout

	Odds ratio	95% CI	p-value
Body mass index (BMI)	1.202	1.029–1.403	0.020
Previous history of gout	71.39	12.790–398.467	<0.001
Chronic kidney disease (eGFR <60 ml/min/1.73m ²)	2.794	1.012–7.719	0.047
Pre-treatment hyperuricemia (Serum uric acid ≥ 6.8 mg/dL)	4.866	1.770–13.378	0.002

Conclusions: TB patients with obesity, history of gout, pre-treatment hyperuricemia, and CKD have higher risk for having gout attack following anti-TB medication. When starting anti-TB therapy in TB patients with these risk factors, physicians should pay attention to the development of gout attack and educate the patients.

REFERENCES:

- [1] Gerdan G, Nurullah A, Ucan ES, et al. Paradoxical increase in uric acid level with allopurinol use in pyrazinamide-induced hyperuricemia. *Singapore Med J*. 2013;54(6):e125–128.
- [2] Postlethwaite AE, Bartel AG, Kelley WN. Hyperuricemia due to ethambutol. *N Engl J Med*. 1972;286(14):761–762.
- [3] Ben Salem C, Slim R, Fathallah N, Hmouda H. Drug-induced hyperuricemia and gout. *Rheumatology (Oxford)* 2017;56(5):679–688.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6590

FRIDAY, 15 JUNE 2018

SLE, Sjögren's and APS – etiology.3, pathogenesis and animal models

FRI0255

INJECTION OF CD40 DNA VACCINE AMELIORATES THE AUTOIMMUNE PATHOLOGY OF NON-OBESE DIABETIC MICE WITH SJÖGREN'S SYNDROME

Y.-B. Zhou¹, X. Yuan¹, Q.-K. Wang¹, H. Zhang¹, G.-S. Wang¹, X.-P. Li¹, Y.-P. Wang², X.-M. Li¹. ¹Rheumatology and Immunology, The First Affiliated Hospital of University of Science and Technology of China, Hefei, Anhui, China; ²Westmead Institute for Medical Research, the University of Sydney, Westmead 2145 NSW, Australia

Background: In the patients with primary Sjögren's syndrome (pSS), overexpression of CD40 was reported. The increased CD40 expression contributes to enhance CD40-CD40L interaction and promotes the inflammatory response in various ways.

Objectives: To investigate whether CD40 DNA vaccine could inhibit the immune response and slow the disease progression of SS in non-obese diabetic (NOD) mice.

Methods: Female 8-week-old NOD mice were randomly divided into 3 groups. CD40 DNA vaccine group received pTaget CD40 DNA vaccine at a weekly dose of 50 ug for 4 weeks. Vector and NS group were administered an equivalent amount of empty vector or NS. Serum anti-CD40 antibody was measured by ELISA. Lymphocytes infiltration in the salivary glands was monitored by focus score (FS) calculation. Salivary CD40 was stained by immunohistochemistry. Splenic lymphocyte phenotypes were analysed by flow cytometry. CD40, TNF- α and IL-6 levels in the salivary glands were detected by PCR. Serum ANA was monitored by immunofluorescence.

Results: 1) CD40 DNA vaccination induced anti-CD40 antibody response. We found the positive rate on anti-CD40 antibody in vaccine group was 80% at week 6 and 100% at week 10, which was significantly increased compared with that in nonvaccinated ones ($p<0.05$ and $p<0.01$, respectively)

2) Expression of CD40 decreased in salivary glands in vaccine group