At week 10, CD40 expression on ductal epithelial and endothelial cells in NOD mice of vaccine group was significantly decreased positive staining. CD40 mRNA expression level showed a significantly reduction compared to vector group (0.51 ± 0.31 vs. 1.56 ± 0.53 , $p < 0.05$).

3) Down-regulation of lymphocytes infiltration in the salivary glands of mice in vaccine group

At week 10, infiltration of lymphocytes was inhibited in treated group while increased in control group ($F = 5.275$, $P < 0.05$). FS was significantly decreased in vaccine group as compared to NS group (2.00 ± 1.73 vs. 11.33 ± 5.51 , $P < 0.05$). Average weight of wet salivary gland and the ratio of average salivary gland weight to body weight of NOD mice in vaccine group were significantly lower than that in control groups ($p < 0.05$ and $p < 0.05$, respectively).

4) CD40 DNA vaccine reduced the expression of TNF- α and IL-6 in the salivary glands

In vaccine group, the expression level of TNF- α mRNA in salivary glands were declined significantly as compared to baseline (0.41 ± 0.25 vs. 0.98 ± 0.16 , $p < 0.05$) and IL-6 mRNA expression was down-regulated compared with control groups at week 10 ($p < 0.05$; $p < 0.01$, respectively).

5) Disturbances in spleen DC and plasma cell subpopulations

At week 6, the total numbers of CD11c⁺DC decreased as compared with two control groups ($p < 0.05$ and $p < 0.05$, respectively). CD11c⁺DC and CD19⁺CD138⁺plasma cells were significantly reduced compared to basal level ($p < 0.01$ and $p < 0.05$, respectively).

6) Level of ANA reduced in the vaccine group

At week 10, the expression of ANA in HEp-2 cells is strong positive (++++ in both control groups, but only positive (+) in vaccine group.

Conclusions: These findings indicate that CD40 DNA vaccine can downregulate the expression of proinflammatory cytokines of TNF- α and IL-6, decrease the percentage of DCs and plasma cells, ameliorate the pathologic change in NOD mice with SS, and thus inhibit the autoimmune inflammation.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3939

FRI0256 EFFECTS OF ANIFROLUMAB ON OXIDATIVE STRESS AND MACROPHAGE ACTIVATION: NOVEL BIOMARKERS AND IMPACT OF TYPE I INTERFERON BLOCKADE IN SYSTEMIC LUPUS ERYTHEMATOSUS

X. Guo¹, S. Wang¹, L. Wang², L.K. Roskos¹, W.I. White¹. ¹Medimmune; ²AstraZeneca, Gaithersburg, USA

Background: Type I interferons (IFNs) drive several aspects of systemic lupus erythematosus (SLE) etiology and pathophysiology. Anifrolumab, a fully human anti-IFN- α receptor monoclonal antibody, substantially reduced disease activity compared with placebo in a Phase IIb study of SLE patients.¹ Previous results indicated beneficial effects of anifrolumab on dysregulated lymphocyte, neutrophil, and complement systems in SLE.² However, the impact of type I IFN blockade remains to be fully elucidated.

Objectives: We characterised biomarkers for oxidative stress and macrophage activation in SLE and assessed downstream effects of anifrolumab on those two key disease pathways.

Methods: In the MUSE study,¹ 305 patients with moderate to severe SLE were randomised in a 1:1:1 ratio to receive placebo or anifrolumab (300 or 1,000 mg) every 4 weeks for 48 weeks. A four-gene expression assay was used to define type I IFN gene signature (IFNGS) test status. In our study, four protein biomarkers were measured in sera of healthy controls (HCs) and SLE patients with samples available at baseline, and 85, 169, and 365 days after administration, using a multiplex luminex immunoassay. Wilcoxon rank-sum test was used to evaluate baseline protein concentrations and posttreatment changes between different groups.

Results: Two oxidative stress biomarkers, peroxiredoxin 4 (PRDX4) and aldose reductase (AKR1B1), were found in greater serum concentrations in 195 SLE patients than in 20 HCs ($p < 0.01$). Their increased concentrations were associated with greater SLEDAI scores and abnormal anti-dsDNA concentrations ($p < 0.05$). PRDX4 concentrations were greater for IFNGS test-high ($n = 144$) vs IFNGS test-low ($n = 51$) patients ($p < 0.01$), whereas AKR1B1 was not associated with type I IFNGS-test status. Anifrolumab suppressed PRDX4 in IFNGS test-high patients compared with placebo ($p < 0.05$). In contrast, no significant changes were observed for AKR1B1 after anifrolumab treatment. Folate receptor 3 (FOLR3) and CD163 are markers of activated macrophages, the former of which has not been characterised in SLE. Our results demonstrated greater concentrations of both FOLR3 and CD163 in IFNGS test-high than in IFNGS test-low patients and HCs, along with significant associations with SLEDAI score and anti-dsDNA concentrations. Anifrolumab reduced serum concentrations of both FOLR3 and CD163 in IFNGS test-high but not IFNGS test-low patients vs. placebo ($p < 0.05$). Moreover, cellular enumeration data demonstrated anifrolumab-induced upregulation of

monocyte count in IFNGS test-high patients, which may result from reduced activation, apoptosis, and/or tissue infiltration of monocytes/macrophages.

Conclusions: Our results identified PRDX4 as an oxidative stress marker and FOLR3 as a macrophage activation marker associated with disease activities and type I IFNGS test status in SLE. Anifrolumab administration elicits reduced oxidative stress and decreased monocyte/macrophage activation, which may contribute to the clinical efficacy of anifrolumab in SLE patients.

REFERENCES:

- [1] Furie R, et al. Arthritis Rheumatol. 2017;69:376–86.
- [2] Guo X, et al. Ann. Rheum. Dis. 2016;75:118.

Disclosure of Interest: X. Guo Shareholder of: AstraZeneca, Employee of: MedImmune, S. Wang Shareholder of: AstraZeneca, Employee of: MedImmune, L. Wang Shareholder of: AstraZeneca, Employee of: AstraZeneca, L. Roskos Shareholder of: AstraZeneca, Employee of: AstraZeneca, W. White Shareholder of: AstraZeneca, Employee of: MedImmune
DOI: 10.1136/annrheumdis-2018-eular.3842

FRI0257 THE STAT4 SLE RISK ALLELE RS7574865[T] IS ASSOCIATED WITH INCREASED IL-12-INDUCED IFN-GAMMA PRODUCTION IN T CELLS FROM SLE PATIENTS

N. Hagberg¹, M. Joelsson¹, D. Leonard¹, S. Reid¹, M.-L. Eloranta¹, J. Mo², M. Nilsson², A.-C. Syvänen³, Y.T. Bryceson^{4,5}, L. Rönnblom¹. ¹Department of Medical Sciences, Rheumatology and Science for Life Laboratories, Uppsala University, Uppsala; ²Respiratory, Inflammation and Autoimmunity, IMED Biotech Unit, AstraZeneca, Gothenburg; ³Department of Medical Sciences, Molecular Medicine and Science for Life Laboratories, Uppsala University, Uppsala; ⁴Department of Medicine, Center for Hematology and Regenerative Medicine, Karolinska Institutet, Karolinska University Hospital Huddinge, Stockholm, Sweden; ⁵Department of Clinical Sciences, Broegelmann Research Laboratory, University of Bergen, Bergen, Norway

Background: Genetic variants in the transcription factor *STAT4* are associated with increased susceptibility to systemic lupus erythematosus (SLE) and a more severe disease phenotype.

Objectives: This study aimed to clarify how the SLE-associated intronic *STAT4* risk variant rs7574865[T] affects the function of immune cells in SLE.

Methods: Peripheral blood mononuclear cells (PBMCs) were isolated from 52 SLE patients in remission (SLEDAI-2K≤4). *STAT4* and *STAT1* protein levels and phosphorylation status in response to interferon (IFN)- α , IFN- γ , or interleukin (IL)-12 were determined, before and after pre-activation of cells with phytohemagglutinin (PHA) and IL-2, in CD56^{dim} NK cells, CD56^{bright} NK cells, B cells, CD4⁺ T cells, CD8⁺ T cells and monocytes by flow cytometry. The frequency of IFN- γ ⁺ cells upon IL-12 or PMA (phorbol 12-myristate-13-acetate) stimulation and the frequency of T-bet⁺ cells was determined in PHA/IL-2 pre-activated cells. Cellular responses and phenotypes were correlated with *STAT4* risk allele carriage (rs7574865[T], 21 homozygous protective, 22 heterozygous and 9 homozygous risk) using an additive linear regression model. Janus kinase inhibitors (JAKi) selective for TYK2 (TYK2i, Compound 35) or JAK2 (JAK2i, BMS-911543) were evaluated for inhibition of IL-12 or IFN- γ -induced activation of SLE PBMCs.

Results: In resting PBMCs, the *STAT4* risk allele was neither associated with protein levels of *STAT4* or *STAT1*, nor cytokine-induced phosphorylation of *STAT4* (pSTAT4) or *STAT1* (pSTAT1). However, following PHA/IL-2-activation, CD8⁺ T cells from *STAT4* risk allele carriers displayed increased levels of *STAT4* ($p = 0.04$), resulting in increased pSTAT4 in response to IL-12 ($p = 0.003$) and IFN- α ($p = 0.04$). Analysis of T cell subsets revealed that the effect was seen in CD45RA⁺CD57⁻ naïve and CD45RA⁺CD57⁻ memory CD8⁺ T cells, but not in CD45RA⁺CD57⁺ memory or CD45RA⁺CD57⁺ effector CD8⁺ T cells. A slight increase in *STAT4* protein levels and IL-12-induced pSTAT4 was also observed in CD4⁺ T cells from *STAT4* risk allele carriers ($p = 0.08$ and $p = 0.09$, respectively). *STAT4* risk allele carriers displayed an augmented IL-12-induced IFN- γ production in CD8⁺ and CD4⁺ T cells ($p = 0.03$ for both), whereas PMA-induced IL-12 production was normal ($p = 0.31$ and $p = 0.10$, respectively). T-bet expression was not correlated to the *STAT4* genotype. The TYK2i and the JAK2i efficiently blocked IL-12 and IFN- γ -induced activation of PBMCs from *STAT4* risk patients, respectively.

Conclusions: T cells from SLE patients carrying the *STAT4* risk allele rs7574865 [T] display an augmented response to IL-12 and IFN- α . This subset of patients may benefit from JAKi treatment.

Disclosure of Interest: N. Hagberg: None declared, M. Joelsson: None declared, D. Leonard: None declared, S. Reid: None declared, M.-L. Eloranta: None declared, J. Mo Employee of: AstraZeneca, M. Nilsson Employee of: AstraZeneca, A.-C. Syvänen: None declared, Y. Bryceson: None declared, L. Rönnblom: None declared