FRI0329

THE NET-EFFECT OF COMBINING RITUXIMAB WITH BELIMUMAB IN SEVERE SYSTEMIC LUPUS ERYTHEMATOSUS

O. Teng¹, T. Kraaij¹, S. Kamerling¹, E. de Rooy¹, P. van Daele², O. Bredewold¹, J. Bakker³, I. Bajema⁴, H. Scherer⁵, R. Toes⁵, T. Huizinga⁵, T. Rabelink¹, C. van Kooten¹. ¹Nephrology, Leiden University Medical Center, Leiden; ²Clinical Immunology, Erasmus Medical Center, Rotterdam; ³Clinical Chemistry; ⁴Pathology; ⁵Rheumatology, Leiden University Medical Center, Leiden, Netherlands

Background: In systemic lupus erythematosus (SLE) patients, excessive formation of neutrophil extracellular traps (NETs) is observed while their degradation is impaired. *In vitro*, immune complexes (ICx) trigger NET formation while NET-derived DNA is a postulated autoantigen for anti-nuclear autoantibodies (ANAs), found in SLE. Based on these self-perpetuating mechanisms in SLE, we hypothesised that interfering with ICx formation should regress NET formation and potentially ameliorate disease.

Objectives: Investigate the effect of Rituximab+Belimumab therapy on pathogenic autoantibodies in relation to NET formation in severe refractory SLE

Methods: A phase 2A, open-label, single arm proof-of-concept study was performed wherein 16 SLE patients with severe, refractory disease were treated with a combination of CD20-mediated B-cell depletion with rituximab and sustained inhibition of B-cell activating factor with belimumab. Besides safety, the study's endpoints were chosen to address the concept of autoantibodies in relation to excessive NET formation.

Results: We demonstrated that SLE-derived immobilised IgG, but not soluble IgG, induced excessive NET formation, confirming *ex vivo* that ICx mediate excessive NET formation in SLE. We showed that therapeutic intervention with RTX+BLM led to specific reductions in ANAs and regression of excessive NET formation. RTX+BLM appeared to be safe and achieved clinically significant responses: low lupus disease activity state was achieved in 10 patients, renal responses in 11 patients and concomitant immunosuppressive medication was tapered in 14 out of the 16 patients.

A. Effect of RTX+BLM on (auto-)antibody levels

B. Effect of RTX+BLM on excessive NET formation

Conclusions: This study provides novel insights into clinical beneficence of reducing excessive NET formation in SLE by therapeutic targeting ANA production with RTX+BLM. Altogether putting forward a new treatment concept that specifically ameliorates underlying SLE pathophysiology.

Acknowledgements: We thank professor C. Pusey (Imperial College London, UK) for kindly providing the Syk inhibitor R406 for our experiments.

Disclosure of Interest: O. Teng Grant/research support from: GlaxoSmithKline, Consultant for: Aurinia Pharmaceuticals, GlaxoSmithKline, T. Kraaij: None declared, S. Kamerling: None declared, E. de Rooy: None declared, P. van Daele: None declared, O. Bredewold: None declared, J. Bakker: None declared, I. Bajema: None declared, H. Scherer: None declared, R. Toes: None declared, T. Huizinga: None declared, T. Rabelink: None declared, C. van Kooten: None declared

DOI: 10.1136/annrheumdis-2018-eular.6415

FRI0330

RITUXIMAB THERAPY IN SLE: EARLY RETREATMENT IS ASSOCIATED WITH LOWER DISEASE ACTIVITY AND A REDUCTION IN CORTICOSTEROID USE

R.L. Batten, S. Batool, C.E. Kidd, A. Lorenzi, J. Vila, K. Walker, B. Griffiths. Rheumatology, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle, UK

Background: The recently published UK guidelines for the management of SLE recommend biologic therapy for severe or refractory disease. The British Isles Lupus Assessment Group Biologics Register (BILAG-BR) has shown rituximab (RTX) to be safe, effective and corticosteroid [CS] sparing when used to treat refractory SLE. In 2013 NHS England published an interim clinical commissioning policy statement with criteria determining when RTX can be used to treat SLE.

Objectives: We evaluated our centre's RTX retreatment strategy in patients with SLE and the consequent outcomes (disease activity and CS dose).

Methods: Records for the first 50 patients receiving RTX for refractory SLE who consented to join BILAG-BR from our centre between December 2013 and January 2016 were analysed (data cut off July 2016). Demographics, disease activity^{BILAG 2004}/SLEDAI-2K, change in CS dose, retreatment schedules and adverse events were analysed.

Results: Median(IQR) age and disease duration were 42.8 (33–53) years and 9.5 (4–15.8) years respectively. Male: female ratio was 1:25. 80% were Caucasian, 6% Asian, 4% Caribbean and 10% other. All patients met SLICC/ACR classification criteria for SLE. The median(IQR) SLEDAI-2K scores and BILAG 2004 scores reduced from 6 (4–8) to 4 (0–4) ($p<0.00001$) and 20 (10–24.5) to 9 (2–15.5) ($p<0.001$) respectively at 6 months. Complete response was achieved in 62.8% patients (defined as loss of all BILAG A and B scores to ≤ 1 B score with no new A/B scores in other organ domains). 66% patients lost all A scores at 6 months. Median(IQR) daily CS dose reduced from 10 mg(0–20) to 5 mg(0.5–9.5) at 6 months ($P<0.001$) and was 5 mg (0–6.63) at last reported visit (median (IQR) 13 (12–19.5) months). 16 patients did not fully respond to baseline treatment but 11 responded to retreatment. Serious infections (requiring hospital admission) occurred in 6 patients (12%).

30/50 patients received their 1st course of RTX at BILAG-BR baseline visit. 23 met criteria for active disease (at least 1A or 2Bs), 6 were taking an unacceptably high maintenance CS dose, and 1 was planning pregnancy. Median(IQR) CS dose in this group at baseline was 10 mg(1.5–20), and 5 mg(3–9) at 6 months. 70.8% demonstrated complete response at 6 months. 17 (57%) went on to have retreatment due to active disease, of which 11 (64.7%) had responded at 6 months post retreatment. Median(IQR) time to retreatment was 8 (6–12) months.

20/50 patients had received retreatment at predetermined intervals prior to their baseline BILAG-BR visit. Median(IQR) numbers of previous RTX courses were 3 (2–6). Median(IQR) CS dose was 8.75 mg(0–11.3) at baseline BILAG-BR visit and 5.5 mg(0–10) at 6 months. Median(IQR) time from baseline to retreatment was 6 (6–9.5) months. Median(IQR) sustained response was 18 (13.5–18) months.

Percentage change (%) in steroid dose at baseline and 6 months

Conclusions: Historically, our centre used time from 1st treatment to flare as a guide to a patient's future RTX retreatment schedule; patients were, on average, treated 2 months earlier than those treated under the current commissioning policy. Findings suggest that earlier retreatment led to sustained disease control and reduction of CS dose with no increase in adverse events. 11/16 incomplete responders responded following retreatment. Early retreatment may be associated with better outcomes for the patient and further research is needed in this area.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4310