FRI0366

PRIMARY RESPIRATORY DISEASE IN PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS: DATA FROM THE SPANISH RHEUMATOLOGY SOCIETY LUPUS REGISTRY (RELESSER) COHORT

J. Narvaez¹, H. Borrell¹, F. Sánchez-Alonso², I. Rúa-Figueroa³, F.J. Lopez-Longo⁴, M. Galindo⁵, J. Calvo-Alen⁶, A. Fernández-Nebro⁷, A. Olivé⁸, J.L. Andreu⁹, R. Blanco¹⁰, M. Andres¹¹, J.J. Alegre¹², T. Cobo-Ibañez¹³, G. Bonilla¹⁴, A. Boteanu¹⁵, E. Díez-Álvarez¹⁶, M. Freire¹⁷, M. Gantes¹⁸, P. García de la Peña¹⁹, R. García-Vicuña²⁰, J. Hernández Beirain²¹, M.L. Horcada²², J. Ibañez²³, A. Juan Mas²⁴, N. Lozano-Rivas²⁵, J.L. Marengo²⁶, I. Castelv²⁷, C. Montilla-Morales²⁸, M. Moreno²⁹, A. Pecondon-Español³⁰, E. Ruiz Lucea³¹, A. Sanchez-Atrio³², G. Santos-Soler³³, F. Toyos³⁴, E. Uriarte Isacelaya³⁵, T.R. Vazquez-Rodriguez³⁶, J.M. Nolla¹, J.M. Pego-Reigosa³⁷ on behalf of EAS-SER (Systemic Diseases Study Group of the Spanish Society of Rheumatology). ¹Rheumatology, Hospital Universitario de Bellvitge, Hospitalet de Llobregat, Barcelona; ²Unidad de Investigación, Sociedad Española de Reumatología, Madrid; ³Rheumatology, Hospital Universitario Doctor Negrin, Las Palmas de Gran Canaria; ⁴Rheumatology, Hospital Universitario Gregorio Marañón; ⁵Rheumatology, Hospital Universitario 12 de octubre, Madrid; ⁶Rheumatology, Hospital Universitario Araba, Vitoria; ⁷Rheumatology, Hospital Universitario de Málaga, Málaga; ⁸Rheumatology, Hospital Germans Trias i Pujol, Badalona, Barcelona; ⁹Rheumatology, Hospital Universitario Puerta de Hierro, Madrid; ¹⁰Rheumatology, Hospital Universitario Marqués de Valdecilla, Santander; ¹¹Rheumatology, Hospital General Universitario de Alicante, Alicante; ¹²Rheumatology, Hospital Universitario Dr Peset, Valencia; ¹³Rheumatology, Hospital Universitario Reina Sofía; ¹⁴Rheumatology, Hospital Universitario La Paz; ¹⁵Rheumatology, Hospital Universitario Ramon y Cajal, Madrid; ¹⁶Rheumatology, Complejo Asistencial Universitario de León, León; ¹⁷Rheumatology, Complejo Hospitalario Universitario de A Coruña, La Coruña; ¹⁸Rheumatology, Hospital Universitario de Canarias, Tenerife; ¹⁹Rheumatology, Hospital Madrid Norte Sanchinarro; ²⁰Rheumatology, Hospital Universitario de La Princesa, Madrid; ²¹Rheumatology, Hospital Insular de Gran Canaria, Las Palmas de Gran Canaria; ²²Rheumatology, Hospital de Navarra, Pamplona; ²³Rheumatology, Hospital Povisa, Vigo; ²⁴Rheumatology, Hospital Son Llatzer, Mallorca; ²⁵Rheumatology, Hospital Virgen de la Arrixaca, Murcia; ²⁶Rheumatology, Hospital de Valme, Sevilla; ²⁷Rheumatology, Hospital de Sant Pau, Barcelona; ²⁸Rheumatology, Hospital Clínico Universitario de Salamanca, Salamanca; ²⁹Rheumatology, Hospital Parc Taulí, Sabadell, Barcelona; ³⁰Rheumatology, Hospital Miguel Servet, Zaragoza; ³¹Rheumatology, Hospital de Basurto, Bilbao; ³²Rheumatology, Hospital Universitario Príncipe de Asturias, Alcalá de Henares, Madrid; ³³Rheumatology, Hospital Marina Baixa, Villajoyosa, Alicante; ³⁴Rheumatology, Hospital Universitario Virgen de la Macarena, Sevilla; ³⁵Rheumatology, Hospital Universitario de Donosti, San Sebastián; ³⁶Rheumatology, Hospital Universitario Lucus Augusti, Lugo; ³⁷Rheumatology, Complejo Hospitalario Universitario de Vigo, Vigo, Spain

Objectives: To investigate the primary respiratory manifestations (PRM) in SLE. **Methods:** All patients in the RELESSER cohort were retrospectively investigated for the presence of PRM. **Results:** At least one PRM was present in 11.3% (365/3215) of cases. The most common was pleurisy, occurring in 21.1% of patients, followed by ALP in 3.6%, PE in 2.9%, PPH 4%, DILD in 2%, DAH in 0.8%, and SLS in 0.8%. The variables associated with the presence of PPM are shown in the following table 1:

	Odds Ratio	95% Confidence Interval	P
Age	1,03	1,02–1,04	0,000
SLEDAI score	1,03	1,00–1,07	0,02
Raynaud phenomenon	1,41	1,09–1,84	0,01
Severe nephritis	1,56	1,18–2,06	0,002
Secondary APS	2,2	1,63–2,97	0,000
Anti-RNP positivity	1,32	1,00–1,75	0,054
Cardiac disease	2,81	1,90–4,16	0,000
Vasculitis	1,81	1,25–2,62	0,002
Hematologic abnormalities	1,31	1,00–1,71	0,048
Neuropsychiatric manifestations	1,49	1,11–2,02	0,009
Gastrointestinal involvement	2,05	1,14–3,67	0,016

After adjusting for known confounders in the multivariable model, PPM remained a risk factor for diminished survival (HR: 3.13).

Conclusions: PPM independently contributed to a decreased survival in SLE

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6436

FRI0367

ASSOCIATION OF VITALITY AND SUBSEQUENT PHYSICAL FUNCTIONING IN SYSTEMIC LUPUS ERYTHEMATOSUS: ANALYSIS OF DATA FROM THE GERMAN LULA COHORT 2002 – 2013

J. Mücke¹, G. Chehab¹, R. Fischer-Betz¹, J. Richter¹, B. Winkler-Rohlfing², M. Schneider¹, R. Brinks¹. ¹Poliklinik of Rheumatology and Hiller-Research Centre of Rheumatology, Heinrich-Heine-University Duesseldorf, Düsseldorf; ²German Lupus Self-Help Community, Wuppertal, Germany

Background: Fatigue, defined as loss of vitality, is a major burden in patients with systemic lupus erythematosus (SLE). It is well-known from cross-sectional studies that fatigue is associated with physical functioning (PF) whereas the effect of fatigue on subsequent PF has not yet been further looked on.

Methods: Data from the German *Lupus-Langzeitstudie* (LuLa) cohort was used to explore the effect of self-reported vitality on PF. The ongoing LuLa study comprises patients from the German SLE self-help group. The current analysis uses data from 2002 to 2013. We looked for associations between the PF score (outcome) and the vitality score of the SF-12 questionnaire in one to four years preceding the outcome. Statistical analysis was accomplished by a linear mixed regression model (LMM) adjusting for the potential confounders age, sex, disease duration and disease activity as surveyed by the Systemic Lupus Activity Questionnaire score (SLAQ).

Results: 1511 patients (93.3% women) provided data with a total of 7841 person-years of observation. Median age at study entry was 46 (interquartile range: 37 to 57). The table 1 shows the regression coefficients of the vitality score (and 95% confidence intervals, CI) estimated by the LMM without and with adjusting for age, sex, disease duration and SLAQ.

Exposure: Vitality scale score	Crude estimates		Adjusted estimates	
	Estimate with 95% CI	p-value	Estimate with 95% CI	p-value
1 year ahead	0.08 (0.05–0.10)	<0.001	0.07 (0.05–0.09)	<0.001
2 years ahead	0.09 (0.07–0.11)	<0.001	0.07 (0.05–0.09)	<0.001
3 years ahead	0.05 (0.03–0.08)	<0.001	0.05 (0.03–0.07)	<0.001
4 years ahead	0.03 (0.001–0.06)	0.038	0.04 (0.01–0.07)	0.002

Conclusions: We found evidence that self-reported vitality in patients with SLE is associated with PF in the subsequent years. The data indicate a "window of opportunity" for treating fatigue symptoms of up to four years for maintenance of PF.

Disclosure of Interest: J. Mücke: None declared, G. Chehab Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study, R. Fischer-Betz Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study, J. Richter Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study, B. Winkler-Rohlfing: None declared, M. Schneider Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study, R. Brinks: None declared

DOI: 10.1136/annrheumdis-2018-eular.4889

FRI0368

COMPARISON OF CLINICAL AND LABORATORY PROFILES IN 3575 SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS WITH AND WITHOUT SJÖGREN'S SYNDROME: DATA FROM A NATIONWIDE PATIENT REGISTRY (RELESSER)

J.G. Ovalles-Bonilla¹, F.J. López-Longo¹, I. Rúa-Figueroa², M. Galindo³, J. Calvo-Alen⁴, R.D. González¹, B. Serrano¹, I. Janta⁵, C. González⁵, I. Monteagudo⁵, J. C. Nieto⁵, J. Martínez⁵, C.N. Saenz⁵, J.M. Pego-Reigosa⁶, on behalf of RELESSER. ¹Rheumatology, Hospital General Universitario Gregorio Marañón, Madrid; ²Rheumatology Division, Hospital Doctor Negrin, Las Palmas GC; ³Servicio de Reumatología, Hospital 12 de Octubre, Madrid; ⁴Rheumatology, Txagorritxu Hospital, Araba, Vitoria; ⁵Rheumatology, Hospital General Universitario Gregorio Marañón, Madrid; ⁶Rheumatology Section, Hospital de Meixoeiro, Pontevedra, Vigo, Spain

Background: The clinical coexistence of Systemic Lupus Erythematosus (SLE) and Sjögren's Syndrome (SS) was recognised in 1959. The prevalence of SS among patients with SLE varies considerably among the published studies (10%–30%). There is still controversy as to whether or not SLE patients with overlapping SS have a distinct and significantly milder lupus.

Objectives: To address the clinical and serologic features of SLE and differences from SLE that occurs in overlap with SS.

Methods: A retrospective cohort of 3575 unselected SLE patients from 45 Rheumatology Units across Spain was evaluated for the presence of overlapping SS