Pulmonary fibrosis was induced by bleomycin in DPP4 knockout (KO) mice and wildtype littermates. Fibrosis of the lungs was additionally evaluated by computer tomography scans (CT). Two oral DPP4 inhibitors were tested in two concentrations in bleomycin-induced skin fibrosis and in sclerodermatous chronic graft-versus-host disease (scl-cGVHD) model. Antiinflammatory effects of DPP4 inhibition were assessed by CD45 staining of fibrotic and non-fibrotic mouse tissue upon DPP4 inhibition. Moreover, chimeric mice were generated by transplanting bone marrow from DPP4-KO mice in WT-littermates (DPP4→WT) and vice versa (WT→DPP4) and fibrosis was by intratracheal injections of bleomycin.

Results: DPP4 positive fibroblasts were increased in fibrotic skin of SSc patients and also in murine models of fibrosis. DPP4 expression is induced by TGF- β in an Erk-dependent manner. DPP4-positive fibroblasts strongly express stress fibres after TGF- β stimulation and released increased amounts of collagen. Mechanistically, inhibition of DPP4 selectively interferes with the TGF- β induced activation of ERK signalling, but does not inhibit TGF- β induced SMAD signalling, or other non-canonical TGF- β pathways involving Fra2, c-Jun, p38, Akt or STAT3. Furthermore, pharmacological inhibition of DPP4 reduced the release of collagen and the expression of myofibroblast markers. DPP4-KO mice are less sensitive to bleomycin-induced pulmonary fibrosis as shown by milder changes on CT, reduced Ashcroft scores and reduced hydroxyproline content. DPP4-KO mice also show reduced skin fibrosis upon bleomycin challenge. Moreover, treatment with DPP4 inhibitors demonstrated potent antifibrotic effects in bleomycin-induced skin fibrosis and experimental scl-cGVHD mouse model. Treatment with DPP4 inhibitors also reduced leukocyte infiltrations into the skin. The extent of pulmonary fibrosis of DPP4→WT was comparable to that of WT→WT control mice. Fibrosis was strongly ameliorated in WT→DPP4 mice and results were comparable to that of DPP4→DPP4 mice, characterising resident cells such as fibroblasts as major target cells for the antifibrotic effects of DPP4 inhibitors.

Conclusions: DPP4 characterises an activated subpopulation of fibroblasts in SSc. Moreover, inhibitors of DPP4 show a significant anti-fibrotic effect in several mouse models of SSc in well tolerated doses. These results may have direct translational implications as DPP4 inhibitors are already in clinical use for diabetes.

Acknowledgements: AS received a scientific training bursary from the European League Against Rheumatism.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7182

FRI0408 INVARIANT NATURAL KILLER T CELLS IN SYSTEMIC SCLEROSIS

A.-C. Pecher, F. Kettemann, J. Henes, S. Duerr-Stoerzer, C. Schneidawind, L. Kanz, D. Schneidawind. universitätsklinikum tübingen, Tübingen, Germany

Background: Systemic sclerosis (SSc) is a rare and heterogeneous immune-mediated disease. Its complex pathogenesis remains incompletely understood. Invariant natural killer T (iNKT) cells have been discussed in several autoimmune diseases¹. They recognise selected endogenous as well as synthetic glycolipids like α -galactosylceramide by their TCR with high affinity and when activated, they typically release high amounts of pro- and/or anti-inflammatory cytokines².

Objectives: To evaluate numbers and function of invariant natural killer T cells in patients with systemic sclerosis and analyse their correlation with disease parameters.

Methods: Human iNKT cells from 91 patients with SSc and 23 healthy controls were analysed by flow cytometric analysis. Their proliferative capacity and cytokine production was investigated following activation with CD1d ligand α -GalCer.

Results: We observed an absolute and relative decrease of iNKT cells in patients with SSc (0.036 iNKT cells/ 10^6 lymphocytes) compared with healthy controls (0.159 iNKT cells/ 10^6 lymphocytes, $p < 0.001$). We could also demonstrate that iNKT cells from patients with SSc showed a significantly decreased expansion capacity upon stimulation with α -GalCer. Interestingly, there was no difference concerning the subtype of SSc, disease severity or whether patients received immunosuppressive drugs.

Conclusions: iNKT cells are deficient and functionally impaired in patients with SSc. Therefore, adoptive transfer strategies using culture-expanded iNKT cells or administration of glycosphingolipids such as α -GalCer could be a novel approach to treat SSc patients.

REFERENCES:

- [1] Wu L, Van KL. Natural killer T cells and autoimmune disease. *Curr Mol Med* 2009;9:4–14.
- [2] Bendelac A, Savage PB, Teyton L. The biology of NKT cells. *Annu Rev Immunol*. 2007;25:297–336.

Acknowledgements: The National Institutes of Health Tetramer Core Facility kindly provided CD1d tetramer reagents. We would like to thank the Flow

Cytometry Core Facility Berg of the University Hospital Tübingen for their excellent technical support.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4531

FRI0409 MYOSITIS SPECIFIC ANTI-HISTIDYL TRNA SYNTHETASE (HISRS) AUTOANTIBODIES DISPLAY HIGH REACTIVITY AGAINST HISRS CONFORMATIONAL EPITOPES AND ASSOCIATE WITH LUNG AND JOINT INVOLVEMENT

C. Cerqueira, N. Renard, A. Notarnicola, E. Wigren, P.-J. Jakobsson, S. Gräslund, I.E. Lundberg. Department of Medicine, Rheumatology Unit, Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden

Background: Autoimmune myositis (rheumatic muscle inflammation) associated with interstitial lung disease (ILD) and arthritis is strongly correlated with the presence of anti-histidyl tRNA synthetase (HisRS) autoantibodies.¹

Objectives: The aims of this study were to investigate: 1) myositis IgG reactivity against HisRS conformational epitopes; and 2) association between clinical manifestations and anti-HisRS reactivity profiles.

Methods: Serum IgG was isolated using a protein G affinity column (from 25 anti-HisRS negative (-) and 19 anti-HisRS positive (+) myositis sera and 24 age/gender matched healthy controls, HC). Autoantibody reactivity was tested by in house ELISA developed against HisRS full-length protein and three HisRS conformational epitopes (WHEP domain – localised in the N-terminal; HisRS without WHEP (HisRS_WHEP); and ABD – anticodon-binding domain located in the C-terminal). Correlations between diagnosis, clinical manifestations and anti-HisRS IgG reactivity were evaluated.

Results: HisRS(+) myositis IgG displayed stronger reactivity against full-length HisRS and HisRS_WHEP (median 372 ng/mL and 334 ng/mL, respectively), compared to WHEP and ABD (6.38 and 6.48 ng/mL). The strongest anti-full-length HisRS reactivity (>371 ng/mL) was detected in patients presenting ILD (10/10 of patients), arthritis (6/10) and polymyositis diagnosis (PM 9/10), in comparison to patients with low anti-HisRS reactivity (<23 ng/mL, ILD – 5/6; arthritis – 3/6; PM – 5/6) or no anti-HisRS reactivity (ILD – 10/28; Arthritis – 8/28; PM – 15/28). On the contrary, myositis patients displaying no anti-HisRS reactivity were largely diagnosed with DM (11/28), skin rash (11/28) and dysphagia (6/28) when compared to patients with the highest anti-full-length HisRS reactivity (1 out of 10 patients was diagnosed with DM, skin rash or dysphagia). Similar associations were observed between the degree of anti-HisRS_WHEP, anti-WHEP or anti-ABD reactivities and manifestations of ILD, arthritis, skin rash or dysphagia, and DM or PM diagnosis. No anti-HisRS reactivity was detected in the HC group.

Abstract FRI0409 – Figure 1. Anti-full-length HisRS IgG reactivity in myositis patients diagnosed with either polymyositis (PM) or dermatomyositis (DM), and presenting manifestations of interstitial lung disease (ILD), arthritis, skin rash and dysphagia. Numbers represent the patients included in the analysis.

Conclusions: This study provides evidences for a possible underlying role of anti-HisRS autoantibodies in the pathogenesis of myositis with interstitial lung disease and joint involvement.

REFERENCE:

- [1] Hamaguchi Y, Fujimoto M, Matsushita T, et al. Common and distinct clinical features in adult patients with anti-aminoacyl-tRNA synthetase antibodies: heterogeneity within the syndrome. *PLoS One* 2013;8(4):e60442.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7048