Correlation between cbPWV/crPWV and Capillary count

Conclusions: Our findings demonstrate that arterial stiffness of the forearm has a relationship with nailfold capillary count and tends to be associated with the extent of disease in patients with SSc. These preliminary data may suggest that vascular damage may concomitantly occur in both capillaries as well as larger arteries of the forearm, which may potentially serve as a novel tool for assessing of early vascular involvement in SSc.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7213

FRI0455

INCREASING INCIDENCE OF ADULT IDIOPATHIC INFLAMMATORY MYOPATHIES: A TEN-YEAR UK EPIDEMIOLOGICAL STUDY

M.J.S. Parker^{1,2}, A. Oldroyd³, M.E. Roberts⁴, R.G. Cooper⁵, H. Chinoy³, on behalf of UKMYONET. ¹Rheumatology Department, Royal Prince Alfred Hospital, Sydney, Australia; ²Rheumatology Department, Salford Royal NHS Foundation Trust, Salford; ³Centre for Musculoskeletal Research, University of Manchester, Manchester; ⁴Greater Manchester Neurosciences Centre, Salford Royal NHS Foundation Trust, Salford; ⁵MRC-ARUK Centre for Integrated Research into Musculoskeletal Ageing, University of Liverpool, Liverpool, UK

Background: Studying the epidemiology of rare conditions such as the idiopathic inflammatory myopathies (IIM) can assist in the identification of risk factors, disease associations and temporal trends. Interrogation of differing geographically and genetically diverse populations can help to construct a more complete picture of underlying disease patterns. A number of UK centres have contributed to national and international IIM research collaborations, but to date there has been no published report detailing the incidence or prevalence of adult IIM in the UK, or to establish the relative proportion of the varying clinical subtypes. Moreover, previous international studies have focussed on specific IIM subtypes, such as inclusion body myositis (IBM) or immune-mediated necrotising myopathy (IMNM), are historic, were undertaken before recent developments in our understanding of the range of IIM subtypes, and utilised widely varying methodologies and case acquisition strategies.

The recently published combined European League Against Rheumatism and American College of Rheumatology (EULAR/ACR) classification criteria for adult and juvenile IIM represent potential progress in identifying IIM, as well as various disease subtypes¹. We present here the first epidemiological study to utilise these new criteria as part of disease verification.

Objectives: Identify and characterise all incident adult cases of IIM between Jan 1 st 2007 and Dec 31 st 2016 in the City of Salford, UK.

Methods: Adults first diagnosed with IIM within the study period were identified by: i) a Salford Royal NHS Foundation Trust (SRFT) inpatient episode IIM-specific ICD-10 coding search; ii) all new patient appointments to SRFT neuromuscular outpatient clinics; iii) all Salford residents enrolled within the UKMYONET study. All patients with 'definite' IIM by the 2017 EULAR/ACR classification criteria were included, as were 'probable' cases if expert opinion agreed. Cases were excluded if <18 years at disease onset, if they did not meet 'probable' criteria, or when 'probable' but expert opinion concluded a non-IIM diagnosis.

Results: The case ascertainment procedures identified 1156 cases which, after review and application of exclusion criteria, resulted in 32 incident cases during

the study period. 23/32 were female with a mean age of 58.1 years. The mean incidence of adult IIM was 17.6/1,000,000 person years (py), higher for females than for males (25.2 versus 10.0/1,000,000py respectively). A significant incidence increase over time was apparent (13.6 versus 21.4/1,000,000py; p=0.032). Using EULAR/ACR classification criteria, the largest IIM subtype (21/32) was polymyositis, followed by dermatomyositis (8/32), inclusion body myositis (2/32) and amyopathic dermatomyositis (1/32). Expert opinion subtype differed from EULAR/ACR Classification criteria in 19/32 cases.

Conclusions: The incidence of adult IIM in Salford is 17.6/1,000,000py, higher in females and is increasing over time. Disagreement exists between EULAR/ACR-derived and expert opinion-derived IIM subtype assignments.

REFERENCES:

- [1] Lundberg IE, Tjälmlund A, Bottai M, et al. EULAR/ACR Classification Criteria for Adult and Juvenile Idiopathic Inflammatory Myopathies and their Major Subgroups. *Ann Rheum Dis.* 2017;76:1955–64.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6079

FRI0456

PREDICTORS FOR DISEASE WORSENING DEFINED BY ORGAN FAILURE IN DIFFUSE SYSTEMIC SCLEROSIS: A EUROPEAN SCLERODERMA TRIALS AND RESEARCH (EUSTAR) ANALYSIS

M.O. Becker¹, N. Graf², R. Sauter³, Y. Allanore⁴, J. Curram⁵, C. Denton⁶, D. Khanna⁷, M. Matucci-Cerinic⁸, J. Pena⁹, J.E. Pope¹⁰, O. Distler¹. ¹Department of Rheumatology and the Centre of Experimental Rheumatology, University Hospital Zurich, Zurich, Switzerland; ²Graf Biostatistics, Winterthur, Switzerland; ³Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, Nuffield Department of Medicine, University of Oxford, Oxford, UK; ⁴Paris Descartes University, Sorbonne Paris Cité, Rheumatology A Department, Cochin Hospital, Paris, France; ⁵Bayer Plc, Reading, UK; ⁶UCL Division of Medicine, Royal Free Campus, London, UK; ⁷Division of Rheumatology, Department of Internal Medicine, University of Michigan Scleroderma Program, University of Michigan, Ann Arbor, MI, USA; ⁸Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy; ⁹Bayer US LLC, Whippany, NJ, USA; ¹⁰Department of Medicine, Division of Rheumatology, University of Western Ontario, St. Joseph's Health Care, London, ON, Canada

Background: Mortality and worsening of organ function are desirable endpoints for clinical trials in systemic sclerosis (SSc). However, these events are relatively rare, making clinical trial design challenging.

Objectives: To identify factors in a population of patients with diffuse SSc from the European Scleroderma Trials and Research (EUSTAR) group database that predict these endpoints and hence allow enrichment of those patients.

Methods: Inclusion criteria were a diagnosis of diffuse SSc and follow-up after 9–15 (12±3) months. This timeframe was chosen to reflect typical clinical trial design. Disease worsening/organ progression was fulfilled if any of the following events occurred: new renal crisis, decrease in forced vital capacity (FVC) ≥10%, new left ventricular ejection fraction (LVEF) <45% or decrease in LVEF by >10% for patients with baseline LVEF <45%, new pulmonary (arterial) hypertension, or

death. These parameters had been defined by expert nominal group technique. Methodologically, two main limitations had to be addressed:¹ the problem of missing data and² the low number of events, which prohibits the simultaneous exploration of a large set of predictors in a regression model. We addressed these issues by (a) imputing multiple predictors based on different algorithms (multiple imputation), and (b) using least absolute shrinkage and selection operator (LASSO) regression for 42 clinical parameters.

Results: Of 1451 patients who met the inclusion criteria, 706 had complete data available on all parameters for disease worsening. Of 706 patients originally evaluated, 228 (32.3%) had disease progression, most of which was either a decrease in FVC (103 patients, 14.6%) or death (92 patients, 13.0%) within the observation period (12±3 months). Of the 42 clinical parameters introduced into the model as outcome predictors, 8 remained in the final regression model, which was chosen by the Bayesian information criterion (table 1). Bootstrap with 10 000 repetitions successfully validated the model.

Abstract FRI0456 – Table 1. Predictive factors in the final LASSO regression model

Factor	p-value	Odds ratio	95% confidence interval
(Intercept)	<0.0001	0.06	0.03–0.14
Age (per 1 year)	0.001	1.02	1.01–1.04
Active digital ulcers	0.026	1.64	1.06–2.54
C-reactive protein elevation	0.002	1.80	1.23–2.63
Significant dyspnoea	0.491	1.20	0.72–2.00
Lung fibrosis	0.0004	2.21	1.43–3.41
Muscle weakness	0.015	1.64	1.10–2.45
Pericardial effusion	0.098	1.65	0.91–2.97
Proteinuria	0.064	1.75	0.97–3.16

Conclusions: The use of the predictive factors presented here could enable cohort enrichment with patients at risk for overall disease worsening in clinical trials.

Acknowledgements: This study was supported by Bayer AG (Berlin, Germany).
Disclosure of Interest: M. Becker: None declared, N. Graf: None declared, R. Sauter: None declared, Y. Allanore Grant/research support from: Actelion Pharmaceuticals US, Bayer AG, Bristol-Myers Squibb, Inventiva, Medac, Pfizer Inc, Roche Pharmaceuticals, Genentech and Biogen IDEC Inc. Sanofi-Aventis Pharmaceutical, Servier, Consultant for: Actelion Pharmaceuticals US, Bayer AG, Bristol-Myers Squibb, Inventiva, Medac, Pfizer Inc, Roche Pharmaceuticals, Genentech and Biogen IDEC Inc., Sanofi-Aventis Pharmaceutical, J. Curram Shareholder of: Bayer AG, Employee of: Bayer Plc, C. Denton Consultant for: Actelion Pharmaceuticals US, Bayer AG, GlaxoSmithKline, CSL Behring, Merck-Serono, Roche Pharmaceuticals, Genentech and Biogen IDEC Inc., Inventiva, Sanofi-Aventis Pharmaceutical, Boehringer Ingelheim, Speakers bureau: Actelion Pharmaceuticals US, Bayer AG, GlaxoSmithKline, CSL Behring, Merck-Serono, Roche Pharmaceuticals, Genentech and Biogen IDEC Inc., Inventiva, Sanofi-Aventis Pharmaceutical, Boehringer Ingelheim, D. Khanna Shareholder of: Eicos, Grant/research support from: Bayer, Bristol-Myers Squibb, Boehringer Ingelheim, Genentech/Roche, NIH, Pfizer, Sanofi-Aventis Pharmaceuticals, Consultant for: Actelion Pharmaceuticals US, Bayer, Bristol-Myers Squibb, Boehringer Ingelheim, Chemomab, Corbus, Covis, Cytori, Eicos, EMD Serono, Genentech/Roche, Pfizer, Gilead, GlaxoSmithKline, Sanofi-Aventis Pharmaceuticals, and UCB Pharma, M. Matucci-Cerinic Grant/research support from: Pfizer, Bristol-Myers Squibb, Actelion, UCB Pharma, Consultant for: Actelion, Bayer, ChemomAb, Genentech/Roche, Inventiva, Lilly, Pfizer, J. Pena Employee of: Bayer HealthCare Pharmaceuticals Inc, J. Pope Grant/research support from: Actelion, Bayer AG, Bristol-Myers Squibb, Merck, Pfizer Inc, Roche, Consultant for: Actelion, Bayer AG, Bristol-Myers Squibb, Merck, Pfizer Inc, Roche, O. Distler Grant/research support from: Actelion, Bayer, Biogen Idec, Boehringer Ingelheim, ChemomAb, EspeRare Foundation, Genentech/Roche, GlaxoSmithKline, Inventiva, Lilly, Medac, MedImmune, Mitsubishi Tanabe Pharma, Pharmacyclics, Novartis, Pfizer, Sanofi, Sinoxa, and UCB in the area of potential treatments of scleroderma and its complications. Patent mir-29 for the treatment of systemic sclerosis licensed, Consultant for: Actelion, Bayer, Biogen Idec, Boehringer Ingelheim, ChemomAb, EspeRare Foundation, Genentech/Roche, GlaxoSmithKline, Inventiva, Lilly, Medac, MedImmune, Mitsubishi Tanabe Pharma, Pharmacyclics, Novartis, Pfizer, Sanofi, Sinoxa, and UCB in the area of potential treatments of scleroderma and its complications. Patent mir-29 for the treatment of systemic sclerosis licensed

DOI: 10.1136/annrheumdis-2018-eular.5296

FRI0457

RELATIONSHIP BETWEEN DIGITAL ULCERS AND SEVERITY OF LUNG FUNCTION TEST IN SYSTEMIC SCLEROSIS OVER A FIVE-YEAR PERIOD

O.B. Ovsyannikova, O. Koneva, L. Ananieva. Department Of Vascular Pathology, Nasonova Research Institute of Rheumatology, Moscow, Russian Federation, Москва, Russian Federation

Background: Systemic sclerosis-related interstitial lung disease (SSc-ILD) is the leading cause of death in SSc. Predictors of the outcomes of ILD in SSc are under investigation.

Objectives: to assess association of the digital ulcers with dynamics of forced vital capacity (FVC) and diffusing capacity for carbon monoxide (DLco) in patients with SSc-ILD.

Methods: It was a longitudinal study involving 83 pts with SSc-ILD (mean age was 46,2±13,4; 69% have limited subset of the disease; 95% were female). The mean duration of follow up was 58,9±12,0 months. At the end of the study a number of pts with digital ulcers (DUs) was 29 (35%). Additionally 77 pts. with SSc-ILD were investigated with HRCT and were divided into 3 groups; The 1 st group (16 pts) with improvement; 2nd group (39 pts) without any changes and 3rd group (22 pts) with worsening of fibrosis.

Results: After 5 years of follow up FVC increased significantly in all pts without DUs (n=54) from 88,5%±19 to 96%±23 (p<0,05); in group 1 – from 92%±20.5 to 106%±19 (p<0,05); in group 2 from 87%±18 to 94%±23.5 (p<0,05) and only in group 3 FVC was stable (88±22 and 87±24.5) (p>0,05). The mean value of FVC in all pts with DUs didn't change (88%±14 and 86%±16, p>0,05) with tendency to decreasing in group 3 (from 83%±12,5 to 74%±13 (p>0,05).

After 5 years of follow up DLco declined significantly in all pts with or without DUs, however in the 1 st group decline of DLco wasn't significant. The decreasing of DLCO was more prominent in group 3 than in group 2. Therefore, in group 2 in patient without DU (n=24) – from 65%±16 to 60%±11 (p<0,05) and in patients with DU (n=14) DLCO changed from 61%±15 to 57%±14 (p<0,05). In 3rd group in patients without DU (n=13) DLCO decreased from 55%±15 to 48%±15 (p<0,05) and in patients with DU (n=9) – from 50%±20 to 44,5%±15 (p<0,05).

Conclusions: In patients without DUs significant increasing of FVC during 5 years long follow up was observed. The worsening of fibrosis on HRCT in pts with DUs was associated with the lowest value of FVC and DLco at the entry and at the end of the study.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4509

FRI0458

THE ONSET, CLINICAL COURSE AND OUTCOMES OF SSC- OVERLAP WITH PM/DM OR RA (SSC-PM/DM AND SSC-RA)

O. Desinova, M. Starovoytova, L. Ananieva. VA Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: Systemic sclerosis (SSc) concurrent with other connective tissue diseases (poly/dermatomyositis, rheumatoid arthritis and others) seem to be still underexplored clinical forms of SSc.

Objectives: To study specific features of the onset, clinical course and outcomes of systemic scleroderma- poly/dermatomyositis (SSc-PM/DM) and SSc-RA overlap syndromes.

Methods: Totally 115 patients were examined, 75 – with SSc-PM/DM and 40 with SSc-RA, among them 98 women and 17 men aged 17–74 years (mean age 44 ±14,5) and disease duration from 6 months to 35 years (median 8[2–8]).

Results: In 18% of overlapping SSc pts the disease manifested with isolated Raynaud's syndrome (RS) at the onset, in 61% – RS came in combination with cutaneous and/or joint and muscle pathology, and in remaining 7% and 2% overlapping SSc manifested with the isolated articular syndrome or muscular involvement (proximal weakness), respectively. During the first 3 years SSc generalisation with emerging signs of PM/DM occurred in 61% of patients, erosive arthritis manifested in 51%; and in 20% of pts arthritis was detected later. Limited skin involvement predominated, while diffuse skin lesions were present in 23% of SSc-PM/DM cases, 1/3 of them showed signs of DM. All pts with overlapping SSc syndrome received a glucocorticoids (GCs)-based combination therapy (SSc-PM/DM 30–60 mg/day, SSc-RA –10–20 mg/day), with NSAIDs and vascular drugs, 74% were administered cytotoxic agents, more often methotrexate (48%) and 7% – antifibrotic drugs. Two types of overlapping SSc evolution patterns were identified: type I is favourable (stabilisation and slow progression without significant disease activity, and preserved work capacity); type II is unfavourable (continuing activity, rapid progression, disability and deaths). Favourable overlapping