sensitivity 0.81 (95% C.I. 0.68 to 0.92); specificity 0.66 (95% C.I. 0.52 to 0.80). Out of the 44 ILD negative patients, 22 (50%) presented a CII value lower than the cut-off, and 13 of them (59%) were found to have a diffusing lung capacity for CO (DLCO)<80% of predicted. At 1 year FU, the CII was significantly correlated with total lung capacity -TLC ($r=0.45$, $p=0.004$) and DLCO ($r=0.29$, $p=0.045$). Out of the 22 patients with a CII <0.1966 but no ILD at visual evaluation, 11 (50%) developed a FVC decline at 1 year, and 8 (36.7%) a DLCO decline.

Conclusions: Here we confirm that quantitative computer-assisted CT of the lungs could be a reliable method for SSC-ILD evaluation and found that it could also be useful in predicting the evolution of lung function in the short-term FU.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.7559

FRI0464

CLINICAL MANIFESTATIONS OF PERSISTENT ANTIPHOSPHOLIPID ANTIBODIES IN SYSTEMIC SCLEROSIS PATIENTS

S. Myriam^{1,2,3}, V. Sobanski^{1,2,4}, A. Lemaire^{1,2}, J. Giovannelli⁴, E. Hachulla^{1,2,4}, B. Lopez^{4,5}, M. Lambert^{1,2,4}, P.-Y. Hatron^{1,2}, S. Dubucquoi^{4,5}, D. Launay^{1,2,4}.

¹Département de Médecine Interne et Immunologie Clinique; ²Centre national de référence maladies systémiques et auto-immunes rares (sclérodermie systémique), CHU Lille; ³Univ Lille; ⁴Lille Inflammation Research International Center (LIRIC), Univ. Lille, U995; ⁵Département d'Immunologie centre de biologie pathologie, CHU Lille, Lille, France

Background: Antiphospholipid antibodies (aPL) can be detected in systemic sclerosis (SSc) patients. While their clinical signification is well known in the primary antiphospholipid syndrome (APS), their frequency and clinical association remain debated matter of issue. In a previous study, we reported the prevalence and their clinical association of a single aPL testing a French cohort of SSc patients. Yet, the persistence of aPL during follow-up of patients has never been assessed.

Objectives: This study aimed to assess i. the persistence of aPL at least 12 weeks after the first dosage and ii. the clinical associations.

Methods: At least 12 weeks apart after the first test, 213 SSc patients were consecutively tested for lupus anticoagulant (LA), anticardiolipin (aCL) and anti-β2glycoprotein 1 (anti-β2Gp1) antibodies. aCL and anti-β2Gp1 were measured using commercial ELISA assays (Orgentec, Trappes, France), positivity was defined as ≥ 10 UGPL/mL (aCL) or ≥ 10 UA/mL (anti-β2Gp1). Clinical associations were studied using binomial logistic regressions.

Results: The mean time between two tests was 13.2 ± 6.3 months. One or more type of aPL was persistent in 7 patients (3%). Patients with persistent aPL developed more frequently venous thrombosis (VT) [4/7 (57%) versus 23/205 (11%), $p=0.006$]. Arterial or/and venous thrombosis were more common in patients with persistent aPL than other patients [4/7 (57%) versus 33/205 (16%), $p<0.001$]. Persistent aPL was also associated with miscarriages [4/6 (67%) versus 33/155 (21%), $p=0.023$].

Association between aPL positivity persistence and clinical manifestations

	aPL positivity	aPL negativity	p
Arterial or venous thrombosis	4 (57.1%)	33 (16.1%)	<0.001
Venous thrombosis	4 (57.1%)	23 (11.2%)	0.006
Arterial thrombosis	1 (14.3%)	14 (6.8%)	0.406
Miscarriage	4 (66.7%)	32 (20.6%)	0.023
Digital ulceration	0 (0%)	40 (19.8%)	0.351
PAH	1 (14.3%)	13 (6.7%)	0.402

Conclusions: The prevalence of persistent aPL in our cohort was 3%. VT and miscarriages were associated with persistent aPL.

REFERENCES:

- Parodi A, Drosera M, Barbieri L, Rebora A. Antiphospholipid antibody system in systemic sclerosis. *Rheumatology* (Oxford) 2001;40:111–12.
- Sanna G, Bertolaccini ML, Mamali A, et al. Antiphospholipid antibodies in patients with scleroderma: prevalence and clinical significance. *Ann Rheum Dis* 2005.
- Marie I, et al. Anticardiolipin, ab2GPI antibodies and lupus-like anticoagulant in SSc.
- Miryakis S, Lockshin MD, Atsumi T, et al. International consensus statement on preliminary classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6640

FRI0465

QUANTITATIVE ASSESSMENT OF INTERSTITIAL LUNG DISEASE IN IDIOPATHIC INFLAMMATORY MYOPATHIES: VISUAL SCORE AND COMPUTERISED PROCESSING ANALYSIS OF HIGH RESOLUTION COMPUTERISED TOMOGRAPHY

S. Barsotti^{1,2}, C. Romei³, E. Cioffi², E. Perrone³, C. Giovannetti², M. Mosca², F. Falaschi³, R. Neri². ¹Department of medical biotechnologies, University of Siena, Siena; ²Rheumatology Unit, University of Pisa; ³Radiology Unit, Pisa University Hospital, Pisa, Italy

Background: High resolution computed tomography (HRCT) is an essential technique for the characterisation of interstitial lung disease (ILD) in patients with idiopathic inflammatory myopathies (IIM). Although several visual scores are available for a semiquantitative evaluation of ILD, an automatic quantitative analysis of lung involvement may represent a valuable improvement to reproducibly determine the extent of the disease.

Objectives: To compare the semiquantitative visual score to a volumetric texture and local volumetric histogram feature-based analysis software (CALLIPER) to quantify the lung involvement in IIM patients.

Methods: 21 consecutive IIM patients (Bohan and Peter criteria) who underwent a HRCT between November 2016 and November 2017 were prospectively enrolled. Twelve were affected by polymyositis and 9 by dermatomyositis. We collected smoking habits data and myositis specific autoantibodies positivity, respiratory symptoms according to MRC dyspnea scale and two patients reported outcome questionnaires: Leicester cough questionnaire (LCQ) and St. George's Respiratory questionnaire (SGRQ). Patients underwent a non-contrast, supine, volumetric CT (≤ 2 mm slice thickness) that was evaluated by an expert radiologist according to Warrick's score (WS) and also quantified by CALLIPER (Mayo Biomedical Imaging Resource – USA). CALLIPER analysis of the extent of interstitial lung abnormalities (ground glass, lung reticulation, honeycombing) was performed and a total score, expressed as percentage of lung involved (ILD%), was obtained. Analysis of the vascular involvement were performed as percent of pulmonary vessel volume (VESS%).

Results: Between the 21 patients, five (23.8%) were smokers. Four patients were positive for anti-synthetase autoantibodies -ASS (3 Jo-1, 1 PL-7), 1 SRP, 1 Mi2. Dyspnea was present in 19 patients (90%): 9 grade 1, 7 grade 2, 3 grade 3 according to MRC scale. There were strict correlations between WS and ILD ($r=0.802$ $p<0.001$) and VESS% ($r=0.746$ $p<0.001$). LCQ score was negative correlated to WS ($r=-0.728$ $p=0.007$) but the correlations were stricter with ILD score ($r=-0.868$ $p<0.001$) and VESS% ($r=-0.801$ $p<0.002$). SGRQ did not correlate both to WS and CALLIPER parameters. Patients with ASS have higher ILD and VESS% compared to seronegative patients (respectively $p=0.003$ and $p=0.006$).

Conclusions: CALLIPER represents an innovative technique to quantify the lung involvement in IIM patients: the main advantage of CALLIPER is the reproducibility that avoids the inter and intra-reader bias. Although more data are needed in larger cohorts, the use of CALLIPER may open new routes for the evaluation of ILD in IIM patients, both in medical practice and in randomised controlled trials.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4265

FRI0466

ARTICULAR AFFECTATION IN SYSTEMIC SCLEROSIS: CORRELATION OF ULTRASOUND FINDINGS WITH CLINICAL, BIOLOGICAL AND RADIOGRAPHICAL FINDINGS

C. De la Puente, J. Quiñones, S. Zagarra. *University Hospital Ramón y Cajal, Madrid, Spain*

Objectives: To evaluate the prevalence of subclinical synovitis in systemic sclerosis (SS), as well as the correlation of sonographic findings in hands with radiographs and with clinical-analytical parameters. In addition, the correlation with scales of functionality will be determined.

Methods: Cross-sectional observational-analytical study that included 40 patients with ES and 23 patients with rheumatoid arthritis (RA) as a control group during the period 10/2015–03/2016. Clinical, analytical, immunological, ultrasound and the radiological characteristics of both groups and their respective statistical correlations were compiled.

Results: The mean age of the group with SS was 53.3 ± 14.1 years; 87.5% were women and 75% of the cases were cutaneously limited. The average time of evolution of the disease was 6.8 years from the diagnosis. Regarding the clinical joint involvement in the carpus and hands, frank arthritis was present in 16 cases (40%), with an average of DAN: 3.77 and TAN: 1.57; with an average of EVA global pain: 30/100 and elevated CRP in 42.5% of cases. The prevalence of subclinical synovitis with ultrasound expression of synovial effusion $\pm$ synovial hypertrophy in the SS group was at carpal level (77%), FCM (60%), PFI 32.5%) and DFI (15%). The statistical linear correlation between findings of carpal erosions at the ultrasound level versus the radiological data was positive only in the involvement at the joint level FCM and FID. In the group with SS, the HAQ questionnaire