coefficient these changes were not statistically significant, suggesting potential confounders (table 2). While women had worse baseline and 2-year WOMAC pain and function scores (all $p \leq 0.04$), this difference was not significantly influenced by neighborhood LEP (all pinteraction NS).

Methodology: Categorical variables are summarized as frequency (percent). Continuous variables are summarized as mean $\pm$ standard deviation

Methodology: Univariate and multivariable linear mixed-effects models were analyzed, with census tracts treated as random effects. In addition to neighborhood Limited English Proficiency; variables included in the model were: age, BMI, sex, ≥ 1 comorbidity; neighborhood percent poverty ($<10\%$, $10\% - <20\%$, $20\% - <30\%$, $30\% - <40\%$, $\geq 40\%$ [reference group]), neighborhood Gini coefficient was also included.

Table 1. Total Joint Replacement Patient characteristics

Characteristic	Total Hip Replacement (N=4009)	Total Knee Replacement (N=3898)
Patient demographics		
Age at surgery (years), mean $\pm$ SD	65.6 $\pm$ 10.6	67.8 $\pm$ 9.5
Sex: female, n (%)	2180 (53.9%)	2346 (60.2%)
BMI (kg/m ²)	27.9 $\pm$ 5.4	30.0 $\pm$ 5.9
Race, n (%)		
White	3794 (94.6%)	3568 (91.5%)
Black	131 (3.3%)	174 (4.5%)
Asian	25 (0.6%)	74 (1.9%)
Other	43 (1.1%)	56 (1.4%)
Unknown	16 (0.4%)	26 (0.7%)
Hispanic ethnicity, n (%)	102 (2.5%)	129 (3.3%)
Patient status		
ASA class		
I-II	3307 (82.5%)	3124 (80.1%)
III-IV	700 (17.5%)	773 (19.9%)
One or more comorbidities	895 (22.3%)	1057 (27.1%)
Census tract characteristics		
Poverty, n (%)		
$<10\%$	3325 (82.9%)	2787 (71.5%)
$10\% - <20\%$	507 (12.7%)	511 (13.1%)
$20\% - <30\%$	115 (2.9%)	129 (3.3%)
$30\% - <40\%$	49 (1.2%)	56 (1.4%)
$\geq 40\%$	13 (0.3%)	27 (0.7%)
Percent poverty, mean $\pm$ SD	6.2 $\pm$ 6.3	6.8 $\pm$ 7.2
Limited English Proficiency (LEP), n (%)		
$<10\%$	2981 (74.4%)	2787 (71.5%)
$10\% - <20\%$	614 (15.3%)	650 (16.7%)
$20\% - <30\%$	209 (5.2%)	251 (6.4%)
$30\% - <40\%$	120 (3.0%)	113 (2.9%)
$\geq 40\%$	85 (2.1%)	97 (2.5%)
Percent LEP, mean $\pm$ SD	8.6 $\pm$ 9.6	9.2 $\pm$ 10.0
Gini coefficient, mean $\pm$ SD	0.45 $\pm$ 0.07	0.45 $\pm$ 0.07
Patient-reported outcome		
Baseline survey results, mean $\pm$ SD		
WOMAC pain	53.3 $\pm$ 17.8	54.4 $\pm$ 17.3
WOMAC function	49.5 $\pm$ 18.0	53.7 $\pm$ 17.6
2-year survey results		
WOMAC pain, mean $\pm$ SD	53.7 $\pm$ 11.3	57.9 $\pm$ 15.6
WOMAC function, mean $\pm$ SD	51.1 $\pm$ 13.1	55.6 $\pm$ 18.1

Methodology: Categorical variables are summarized as frequency (percent). Continuous variables are summarized as mean $\pm$ standard deviation

Table 2. Impact of every 10% change in neighborhood limited English proficiency on WOMAC pain and function

Time-point	WOMAC pain Estimate $\pm$ SE	p-value	WOMAC function Estimate $\pm$ SE	p-value
Total Hip Replacement				
Baseline (Univariate)	-1.4 $\pm$ 0.3	<0.001	-1.7 $\pm$ 0.3	<0.001
2-year (Univariate)	-0.9 $\pm$ 0.2	<0.001	-1.2 $\pm$ 0.2	<0.001
Baseline (Multivariable)	-0.6 $\pm$ 0.3	0.10	-0.8 $\pm$ 0.3	0.07
2-year (Multivariable)	-0.4 $\pm$ 0.2	0.054	-0.6 $\pm$ 0.3	0.01
Total Knee Replacement				
Baseline (Univariate)	-1.7 $\pm$ 0.3	<0.001	-1.8 $\pm$ 0.3	<0.001
2-year (Univariate)	-1.0 $\pm$ 0.3	<0.001	-1.1 $\pm$ 0.3	<0.001
Baseline (Multivariable)	-0.8 $\pm$ 0.3	0.02	-0.5 $\pm$ 0.3	0.13
2-year (Multivariable)	-0.2 $\pm$ 0.3	0.43	-0.2 $\pm$ 0.3	0.54

Methodology: Univariate and multivariable linear mixed-effects models were analyzed, with census tracts treated as random effects. In addition to neighborhood Limited English Proficiency; variables included in the model were: age, BMI, sex, ≥ 1 comorbidity, neighborhood percent poverty ($<10\%$, $10\% - <20\%$, $20\% - <30\%$, $30\% - <40\%$, $\geq 40\%$ [reference group]), neighborhood Gini coefficient was also included.

Conclusions: Patients coming from neighborhoods with lower english proficiency have worse pain and function scores in unadjusted models. However when adjusted for potential confounders, the difference is not significant. Community factors which contribute to healthcare disparities are multidimensional, prospective studies collecting individual LEP data would be warranted to study this aspect in detail.

REFERENCES:

- John-Baptiste A, et al. The effect of English language proficiency on length of stay and in-hospital mortality. J Gen Intern Med 2004.
- Karliner LS, et al. Identification of limited English proficient patients in clinical care. J Gen Intern Med 2008.

Disclosure of Interest: B. Mehta: None declared, J. Szymonifka: None declared, S. Dey: None declared, I. Navarro-Millan: None declared, S. Grassia: None declared, L. Mandl Grant/research support from: Boehringer-Ingelheim, A. Bass Grant/research support from: Abbott, Pfizer, L. Russell: None declared, M. Parks Grant/research support from: Zimmer, Consultant for: Zimmer, M. Figgie Shareholder of: Mekanika, Y.-Y. Lee: None declared, J. Nguyen: None declared, S. Goodman: None declared

DOI: 10.1136/annrheumdis-2018-eular.2330

FRI0612

HOW EFFECTIVE ARE INTERVENTIONS TARGETING PATIENT ACTIVATION IN PEOPLE WITH LONG TERM PHYSICAL CONDITIONS? A SYSTEMATIC REVIEW

B. Jones^{1,2}, S. Hewlett^{1,2}, D. Harcourt³, A. Hunt², E. Dures^{1,2}. ¹Centre for Health and Clinical Research, University of the West of England (UWE), ²Academic Rheumatology, University Hospitals Bristol NHS Foundation Trust, ³Centre for Appearance Research, University of the West of England (UWE), Bristol, United Kingdom

Background: Patient activation is an increasingly referenced concept in the self management of long term physical conditions. It refers to someone's ability to take an active role in self-managing their health. There has been a substantial increase in the number of studies trailing interventions targeting patient activation in a variety of formats. The findings from this review will contribute to maximising the impact of current rheumatology interventions, and to increase the understanding of patient activation to develop a framework to describe its core components within a rheumatology context.

Objectives: This study aimed to identify the effectiveness of interventions targeting patient activation in people with long-term conditions, including rheumatoid arthritis.

Methods: Studies that were randomised control trials of interventions targeting patient activation in adults with long term physical conditions were included in the review. PsycINFO, Medline, AMED, CINAHL, and ProQuest were searched during January 2017, as well as a variety of grey literature locations, snowballing and hand-searching to identify potential studies for inclusion. The Cochrane Collaboration's Risk of Bias Tool was used to determine the methodological quality of included studies, and any differences were resolved by the two reviewers. Authors were contacted if any additional information was required to extract data or to clarify risk of bias.

Results: 17 papers were included in the review. There was a great deal of heterogeneity in the types of interventions available targeting patient activation in a variety of populations, and a meta-analysis was not possible as a result. Interventions were delivered via face to face, telephone, internet and resource-based approaches. The studies also included a mix of group and individually delivered interventions. Outcome measures used also varied, including both direct and proxy measures for patient activation.

The findings suggest that in studies with interventions that are more structured, there appear to be a trend towards differences in patient activation scores between groups in favour of the intervention. It is possible that these increases in patient activation may occur with an increase in health literacy, as the literature suggests an association between the two constructs. This is discussed in terms of a proposed model for patient activation for rheumatology.

Conclusions: There is some initial evidence for the benefit of increasing structure in interventions targeting patient activation. There is no evidence for a medium of delivery that is most effective in people with long term conditions. Studies would benefit from reporting information about engagement and adherence to interventions to understand the mechanisms behind engaged participants becoming more active in managing their health. There is a need for increased research into patient activation within Rheumatology.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3660

FRI0613

USEFULNESS OF SMARTPHONES APPLICATIONS IN THE FOLLOW-UP OF PATIENTS WITH INFLAMMATORY RHEUMATOLOGICAL DISEASES: ARE THEY REALLY BENEFICIAL IN OBJECTIVE TERMS?

C. A. Guillen-Astete¹, B. Laso-Jimeno². ¹Rheumatology Department, Moncloa University Hospital, ²Ramon y Cajal University Hospital, Madrid, Spain

Background: The use of computer programs developed for their use through smartphones has been incorporated into the health field. In recent years there have been countless new applications aimed at easing the work of the clinician through access to commonly used tools, image atlases, real-time information or to facilitate the monitoring of patients. In this last group, there are different applications oriented to multiple purposes.

Objectives: The aim of this study is to assess the performance of two personally developed applications for the follow-up of patients with gout and rheumatoid arthritis (ReumApp-Gota and ReumApp-AR).

Methods: We surveyed 65 patients who were users of ReumaApp-Gota and 35 patients who were users of ReumaApp-AR. All were direct users with at least 6 months of experience with the application. Both applications were developed by the author of the work on a free distribution platform and whose extension and