FRI0676

RESPONSIVENESS OF PATIENT REPORTED OUTCOMES MEASUREMENT INFORMATION SYSTEM (PROMIS®) COMPUTERIZED ADAPTIVE TESTS (CATS) IN SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

S. Kasturi¹, J. Szymonifka², J. Berman², K. Kirou², A. Levine², L. Sammaritano², L. A. Mandi². ¹Medicine/Rheumatology, Tufts Medical Center, Boston, ²Medicine/Rheumatology, Hospital for Special Surgery, New York, United States

Background: The accurate measurement of patient reported outcomes is a priority for patient-centered care in SLE, a chronic systemic disease with significant impact on quality of life. PROMIS CATs are precise measures of physical, mental, and social health with construct validity in SLE. The longitudinal responsiveness (sensitivity to change) of PROMIS CATs in SLE patients is unknown.

Objectives: To evaluate the responsiveness of PROMIS CATs in SLE outpatients using patient and physician-derived anchors.

Methods: Adult SLE patients were recruited from an SLE Center of Excellence. Subjects completed 14 selected PROMIS CATs at two visits a minimum of one month apart. SLE disease activity was measured with a patient global assessment of change, a physician global assessment and the physician-derived SELENA-SLEDAI. Responsiveness of PROMIS scores was evaluated using known-groups validity. Effect sizes were compared across groups of patients who differed in their patient global assessment of change, physician global assessment, and SELENA-SLEDAI using Wilcoxon rank-sum tests.

Results: A diverse cohort of 228 SLE patients, including 45 (19.8%) patients flaring by SELENA-SLEDAI, completed baseline surveys. Follow up surveys were completed by 190 (83%). There was poor agreement between patient and physician global assessments (weighted kappa statistic [95% CI]=0.16 [0.04–0.28]. Using the patient-based anchor, Anger, Pain Interference, and Physical Function CATs showed low to moderate responsiveness (table 1). Using the physician global assessment, only Anxiety CAT showed low to moderate responsiveness (effect size -0.27, -0.17, and 0.06 [p=0.03] with ≥ 0.5 point decrease, <0.5 point change, and ≥ 0.5 point increase respectively), while with the SELENA-SLEDAI as anchor, only Applied Cognition-Abilities CAT showed responsiveness (0.34, -0.01, 0.0 [p<0.01] with ≥ 3 point decrease, <3 point change, and ≥ 3 point increase respectively).

PROMIS CAT	Patient Global Assessment of Change			p-value
	Better (n=74)	Same (n=79)	Worse (n=33)	
Ability to Participate in Social Roles	0.14	0.00	0.00	0.19
Anger	-0.37	-0.08	0.00	0.03
Anxiety	-0.38	-0.04	-0.02	0.13
Applied Cognition-Abilities	0.15	0.00	-0.03	0.30
Applied Cognition-General	-0.08	0.00	-0.06	0.47
Concerns				
Depression	-0.15	0.00	0.36	0.15
Fatigue	-0.16	0.00	-0.15	0.51
Mobility	0.14	0.00	0.00	0.26
Pain Behavior	-0.14	0.00	0.10	0.07
Pain Interference	0.00	0.00	0.25	0.02
Physical Function	0.15	0.01	-0.14	0.02
Satisfaction with Social Roles	0.18	0.00	0.00	0.21
& Activities				
Sleep Disturbance	0.00	-0.14	-0.04	0.50
Sleep-Related Impairment	-0.07	0.00	-0.02	0.95

Conclusions: PROMIS CATs showed modest responsiveness to patient-reported, but generally not physician-derived changes in lupus health status in domains of anger, pain interference, and physical function. These data suggest that certain PROMIS CATs are precise and sensitive tools which may be used to measure and monitor important aspects of the patient experience of lupus not captured by physician-derived metrics. Further studies are needed to evaluate the responsiveness of PROMIS CATs in populations with greater SLE disease activity and more regular follow up.

Acknowledgements: Funding was provided by the Rheumatology Research Foundation Scientist Development Award.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.3152

FRI0677

PREVALENCE AND SEROLOGICAL PROFILE OF ANTI-DFS70 POSITIVE SUBJECTS: DATA FROM A ROUTINE ANA COHORT

T. Carbone^{1,2}, V. Pafundi², G. Tramontano¹, M. Gilio¹, C. Esposito¹, M. C. Padula¹, A. A. Padula¹, S. D'Angelo¹. ¹Rheumatology Institute of Lucania – IReL, ²Immunopathology Laboratory, San Carlo Hospital, Potenza, Italy

Background: Anti-Dense Fine Speckled 70 (DFS70) antibodies are a common finding in clinical laboratory referrals. High prevalence of DFS70 autoantibodies in

healthy population and usual negative association with Antinuclear Antibody (ANA)-associated autoimmune rheumatic diseases were reported.

Objectives: The aim of this study was to evaluate the prevalence of anti-DFS70 antibodies in a routine diagnostic laboratory setting and their association with other circulating serum autoantibodies.

Methods: Consecutive sera submitted for ANA screening were analyzed for anti-DFS70 antibodies by indirect immunofluorescence (IIF) (n=3175, 1030 men and 2145 women) then confirmed by Immunoblotting. IIF DFS70 positive adult patients were recruited previous written consent and tested for the following autoantibodies: anti-ENA, anti-dsDNA, -anti-TPO, anti-TG, anti-tTg, aCL, anti-PCA, AMA, ASMA, anti-LKM, anti-MPO, anti-PR3 and ASCAs.

Results: The prevalence of anti-DFS70 antibodies was 1.7% (n= 55) in the whole population and 4.6% in the ANA-positive samples. Comparison between DFS70 IIF and Immunoblotting showed an excellent correlation between the two methods (R=0.99). Analysis of anti-DFS70 antibodies titers distribution revealed that 63% of the total cohort showed high titers ($\geq 1:640$). Gender difference (female: male, 4:1) was observed in anti-DFS70 positive group and in anti-DFS70 negative/ANA positive group. The prevalence of anti-DFS70 positive female (2.1%, 45/2145) was statistically significant higher than males (1.0%, 10/1030) (p<0.05). The comparison among referring sources evidenced a prevalence of anti-DFS70 positive subjects from Endocrinology Department (9.1% versus 2.6% from Haematology, 2.1% from outpatients, 1.6% from Neurology, 1.2% from Internal Medicine, 1% from Cardiology, 0.6% from Rheumatology). Of note, our data evidenced isolated reactivity of anti-DFS70 autoantibodies in males group, while 51% of females showed concomitant disease-marker autoantibodies.

Conclusions: We found a prevalence of anti-DFS70 antibodies in adult sera from routine ANA cohort of 1.7%. The serological profile of DFS70-positive females required further investigations in order to define the presence of serum autoantibodies. Anti-DFS70 reactivity in male population may represent an exclusive biomarker predicting the absence of other autoantibodies.

Acknowledgements: The authors would like to express their especial appreciation and thanks to Prof. Ignazio Olivieri who died on July 28th, 2017. He was an example of strength and tenacity with a contagious enthusiasm for a rigorous scientific research.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.6476

FRI0678

VALIDITY OF THREE 0–10 VISUAL ANALOG SCALES (VAS) FOR QUANTITATIVE PHYSICIAN ASSESSMENT OF INFLAMMATION, DAMAGE, AND DISTRESS TO SUPPLEMENT A PHYSICIAN GLOBAL ASSESSMENT 0–10 VAS

T. Pincus¹, I. Castrejon¹, J. A. Block¹. ¹Rheumatology, Rush University Medical Center, Chicago, United States

Background: Rheumatologists generally view their primary goal as control of inflammation in order to prevent long-term damage, and quantitative assessment involves measures of inflammatory activity (lab tests, joint counts, and indices). Although structural damage and patient distress (fibromyalgia, depression, etc.) are widely recognized, these problems generally are described narratively, and not assessed quantitatively. Recent advances in control of inflammation, as well as increased degenerative diseases in an aging population and recognition of a high prevalence of fibromyalgia, may have shifted rheumatologists' patient mix more prominently toward damage and distress vs inflammation.

Objectives: To analyze physician global assessment (DOGGL) on a 0–10 visual analog scales (VAS), and 3 additional 0–10 VAS for inflammation, damage, and distress, as well as estimates of the proportion of each to explain DOGGL.

Methods: Rheumatologists at one academic site complete a 0–10 DOGGL VAS, 3 further 0–10 VAS to assess inflammation (reversible disease) (DOCINF), joint and other organ damage (irreversible disease) (DOCDAM), and patient distress (fibromyalgia, depression), etc. (DOCSTR), in routine care. The proportion of DOGGL attributed to inflammation, damage, and distress (total=100%) also is estimated. Mean values were analyzed in a cross-sectional study of 570 patients, and compared in subgroups with rheumatoid arthritis (RA), osteoarthritis (OA), or fibromyalgia (FM), using t tests and analysis of variance (ANOVA).

Results: Mean DOGGL VAS was 4.4/10 in all patients, 4.4 in 131 with OA, 4.6 in 98 with RA, and 5.2 in 89 with FM (table 1). Highest mean scores were seen for DOCINF in RA, DOCDAM in OA, and DOCSTR in FM (p<0.001), indicating face validity. Nonetheless, mean DOCDAM was higher than DOCINF in all groups, including RA, and mean estimates of the proportion of DOGGL attributed to damage was greater than to inflammation in all groups (table 1). Scores for DOCSTR were higher than for DOCINF in all groups other than RA.

Table 1 Mean VAS Scores and % of clinical management decision attributed to inflammation, damage, and distress in patients with rheumatic diseases

Mean VAS Scores	ALL N=570	RA N=98	OA N=1311	FM N=89	P*
VAS DOCGL	4.4 (1.6)	4.6 (1.8)	4.4 (1.5)	5.2 (1.6)	0.04
VAS DOCINF	1.8 (2.0)	2.8 (2.4)	0.7 (1.1)	0.8 (1.3)	<0.001
VAS DOCDAM	3.1 (2.2)	3.8 (2.3)	4.4 (1.8)	1.7 (1.9)	<0.001
VAS DOCSTR	2.1 (2.9)	1.2 (2.2)	1.5 (2.5)	6.0 (2.5)	<0.001
p (DOCINF vs DOCDAM)	0.001	0.01	<0.001	<0.001	
p (DOCINF vs DOCSTR)	0.11	<0.001†	<0.001	<0.001	
Mean % of clinical management decision attributed to...					
% inflammation	29 (31)	39 (29)	12 (19)	6 (11)	<0.001
% damage	48 (35)	52 (30)	73 (31)	18 (23)	<0.001
% distress	22 (34)	9 (20)	15 (27)	76 (27)	<0.001

*ANOVA – RA vs OA vs FM † -only comparison in which DOCINF higher than DOCSTR

Conclusions: VAS scores and estimates of the proportions of DOCGL attributed to damage were higher than for inflammation in all groups, including RA patients as a group. Control of inflammation remains a primary concern for rheumatologists, but has improved considerably in recent years, and damage and distress may have become more prominent in routine care. Systematic quantitation of inflammation, damage, and distress, in addition to DOCGL, appears feasible and of value in routine clinical care.

Disclosure of Interest: T. Pincus Shareholder of: Dr. Pincus holds a copyright and trademark for MDHAQ and RAPID3 for which he receives royalties and license fees. All revenue is used to support further development of quantitative questionnaire measures for patients and doctors in clinical rheumatology care., I. Castrejon: None declared, J. Block: None declared

DOI: 10.1136/annrheumdis-2018-eular.5588

FRI0679 HIGH SENSITIVITY CARDIAC TROPONIN T IN PSORIATIC ARTHRITIS PATIENTS: A CROSS-SECTIONAL CONTROLLED STUDY

V. Eurer¹, S. Shenhar-Tsarfaty², S. Berliner², U. Arad¹, D. Paran¹, O. Rogowski², I. Shapira², H. Matz³, O. Elkayam¹. ¹Rheumatology, ²Internal Medicine, ³Dermatology, Tel Aviv Medical Center, Tel Aviv, Israel

Background: High sensitivity cardiac troponin (hs-cTnT) is a biomarker for cardiovascular disease (CVD) in the general population. (1) Psoriatic arthritis (PsA) confers an increased risk for CVD (2) but there are no biological markers to stratify CVD risk in PsA patients.

Objectives: To determine the level of serum hs-cTnT in a PsA cohort and controls without inflammatory disease, and further characterize the PsA cohort with detectable hs-cTnT.

Methods: Serum hs-cTnT level was measured with a sandwich immunoassay method in a consecutive PsA cohort (n=96). The control cohort was based on apparently healthy individuals recruited during routine annual health examination, Tel-Aviv Medical Center Inflammatory Survey (TAMCIS) (n=6,052). hs-cTnT was measured in carefully matched controls (n=88), manually selected from the TAMCIS, based on gender, age, BMI, hypertension, and hyperlipidemia prevalence. Hs-cTnT ≥5 ng/L was used as a cutoff for the detectable level. Multiple regression analysis was used to determine the factors associated with hs-cTnT.

Results: The characteristics of the PsA cohort are presented in table 1. Remarkably, in the majority of patients, both skin and joint disease were well controlled. PsA and TAMCIS cohorts had a similar range of age (51.5 vs 48 yr) but different gender representation: 47% vs 72.5% of males (p<0.001). PsA exhibited a higher prevalence of traditional CVD risk factors compared to the TAMCIS cohort: BMI 28 vs 26.5 (p=0.002), current smokers 20.8% vs 10.1% (p=0.002), hypertension 25% vs 15% (p=0.007), dyslipidemia prevalence 34% vs 27% (p=0.101), diabetes 19.8% vs 4.6% (p<0.0001). Due to these differences, a matched control group was used for comparison of troponin. Detectable hs-cTnT was present in 29.5% of the PsA patients compared to 19.3% in the controls. (p=0.114) Factors associated with detectable hs-cTnT in PsA were consistent with traditional factors and included male gender (p=0.007), age (p=0.005), hypertension (p<0.001), and DM (p<0.001). No correlation between detectable hs-cTnT level and psoriasis/PsA duration, disease severity, treatment with DMARDs or biologics was found.

Table 1 Characteristics of the PsA cohort (n=96)

Gender, % males	47%
Age, years	51.5±13
Psoriasis duration	17±13
Psoriasis activity: remission/mild/moderate to severe	34%/56%/10%
PsA duration	9.4±8.1
PsA activity: remission/low/moderate/severe	37%/33%/22%/8%
Concomitant sDMARDs treatment (%)	49%
Concomitant biologic treatment (%)	65%
Biologic treatment exposure (past or present) (%)	71.8%

Conclusions: This is the first study to report a detectable hs-cTnT level in up to 30% in patients with well controlled PsA, asymptomatic for CVD, warranting a special attention to monitoring CVD risk factors and manifestations in this group. Traditional CVD risk factors but not measures of disease activity were associated with detectable hs-cTnT. The latter may be explained by a potential positive impact of anti-rheumatic therapies on the cardiovascular profile. Further prospective studies addressing the predictive role of hs-cTnT for CVD events in PsA are needed.

REFERENCES:

1. Saunders JT, et al. Circulation 2011;123.
2. Ogdie A, et al. Ann Rheum Dis 2015;74.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.4002

FRI0680 TETHERIN, A NOVEL TYPE I INTERFERON BIOMARKER ON BLOOD LEUCOCYTES CAN CAPTURE INTERFERON STATUS AND CORRELATES WITH USTEKINUMAB (STELARA) THERAPY RESPONSE IN PSORIATIC DISEASE.

Y. M. El-Sherbiny¹, L. Savage¹, A. Alase¹, E. Vital¹, M. Wittmann¹, P. Emery¹, D. McGonagle¹. ¹Leeds Institute of Rheumatic and Musculoskeletal Medicine, University Of Leeds, Leeds, United Kingdom

Background: Recently Ustekinumab, an IL12/23 p40 monoclonal antibody that is licensed for Psoriatic Arthritis and Psoriasis, showed promising results in phase II trials in SLE- a prototype IFN mediated disease. We previously confirmed a novel IFN gene scoring system associated with the different SLE symptoms (manuscript in review). Additionally, we also recently validated in two independent cohorts the value of BST2/ tetherin as IFN-I biomarker assay correlates with clinical activity and predicts clinical flare in SLE. Given that psoriasis has several SNPs in the IFN pathway; Thus mechanistic studies of the effect of Ustekinumab on the IFN pathway can be explored in this disease setting.

Objectives: This work tested the hypothesis that a novel interferon type I (IFN-I) status markers in the blood and skin of Ustekinumab treated psoriasis patients might correlate with therapy responses and provide insights into how p40 blockers may affect IFN pathways in a relevant human disease model.

Methods: Skin biopsies and peripheral blood at baseline (before therapy, 24weeks, 54 weeks) from 23 Ustekinumab patients with psoriasis who had ultrasound imaging confirmed subclinical enthesopathy) were recruited. Cellular immunophenotyping was performed using multi-parameter flow cytometry to detect tetherin on (Monocyte, B cells, T cells and neutrophils). All data was compared to age-matched samples from healthy controls (HC). Skin biopsies were digested and RNA extracted, qPCR of common ISGs genes were analysed in lesional and peri-lesional at corresponding time points.

Results: Tetherin showed a higher level of expression on blood subsets of psoriasis compared to HC at baseline on Monocytes, T cells, NK cells and B cells (p=0.003, =0.005, =0.035, =0.002) compared to the baseline. No significant changes observed between baseline and 2nd visit 24 weeks. Interestingly, a substantial reduction in tetherin expression at (52 weeks) in psoriasis was observed in all subsets in Monocytes, T cells, NK cells and B cells (all p<0.0001) correlating with patient response to therapy. IFN signature by TaqMan revealed higher expression in skin biopsies distinctive ISGs compared to HC, e.g. IFI27, STAT1, IFI16 and IRF7 all corrected post-Ustekinumab therapy to normal levels. Paired analysis revealed a stronger IFN signature in lesional biopsies vs non-lesional biopsies, e.g. IFI27 (p=0.0312, Wilcoxon matched pair-rank test).