CCP positive patients. Much less is known about prognostic markers in seronegative RA. Several studies report worse drug survival and worse patient reported outcomes in women with RA. This affects outcomes such as the 28-joint disease activity score (DAS28) and the health assessment questionnaire (HAQ). How these differences relate to autoantibody status is unknown.

Objectives: To investigate if the relation between sex and clinical outcomes varies by autoantibody status in patients with early RA.

Methods: An inception cohort of patients with early RA (n=233; symptoms duration ≤12 months), recruited in 1995–2005, was studied. All the patients fulfilled the 1987 American College of Rheumatology criteria for RA. The patients were managed according to usual care, with no pre-specified protocol for pharmacotherapy or rehabilitation. In a structured follow-up program, all patients were examined by the same rheumatologist. In the present study we divided the patient population in three groups according to autoantibodies status: RF and anti-CCP seropositive (double positive), RF or anti-CCP seropositive, RF and anti-CCP seronegative (double negative). We examined the relation between sex and different outcomes at 12 months (EULAR good response, clinical remission (DAS28 <2.6), HAQ≤0.5 and low pain score (VAS pain 0–100 of <20) by means of logistic regression.

Results: Complete data on autoantibody status at baseline was available for 201 patients (mean age at inclusion 61 years, 72% female, 60% RF positive and 58% anti-CCP positive). Twenty-eight% of the patients were double negative, 27% were single positive and 45% were double positive. Mean baseline DAS28 was 4.53. All patients were treated with a conventional synthetic DMARD (48% with methotrexate). Oral glucocorticoids were prescribed in 38% of patients. At the 1 year follow up, 19% had a EULAR good response, 21% were in remission, 40% had low pain and 53% low HAQ. Male patients in the double negative group were more likely to reach remission (odds ratio (OR) 6.40; 95% confidence interval (CI) 1.6–26.2) and EULAR good response (OR 4.67; 95% CI 1.2–18.3) compared to females. There were no such associations among the double positive patients (Table). Results were similar in analyses adjusted for DAS28 at baseline (Table). There was a similar pattern among double negative patients for low pain at 1 year (OR for male vs female patients 2.25; 95% CI 0.58–8.67 – adjusted for baseline pain), but no association between male sex and low HAQ at 1 year in double negative patients (OR 0.99; 95% CI 0.23–4.22 – adjusted for baseline HAQ) or the other subgroups.

Abstract SAT0093 – Table 1

Impact of sex on the chance of EULAR good response and clinical remission at the 1-year follow-up in patients with early RA. Logistic regression analysis.

	Sex	EULAR good response OR (95% CI)		Clinical remission* OR (95% CI)	
		Crude	Adjusted for baseline DAS28	Crude	Adjusted for baseline DAS28
All	Female (n=144)	(reference)	(reference)	(reference)	(reference)
	Male (n=57)	1.9 (0.9-3.7)	1.9 (0.9-3.7)	2.0 (1.0-3.9)	2.1 (1.0-4.1)
RF and anti- CCP negative	Female (n=42)	(reference)	(reference)	(reference)	(reference)
	Male (n=14)	4.7 (1.2-18.3)	6.0 (1.3-26.4)	6.4 (1.6-26.2)	6.3 (1.5-25.9)
RF or anti- CCP positive	Female (n=41)	(reference)	(reference)	(reference)	(reference)
	Male (n=13)	3.2 (0.6-17.2)	3.4 (0.6-20.6)	3.8 (0.8-17.7)	2.1 (0.3-12.3)
RF and anti- CCP positive	Female (n=61)	(reference)	(reference)	(reference)	(reference)
	Male (n=30)	1.3 (0.4-4.5)	1.2 (0.3-4.2)	0.9 (0.3-2.9)	1.1 (0.3-3.7)

*DAS 28 <2.6

Conclusions: In the subgroup of patients with seronegative early RA, male patients are more likely than female patients to reach DAS28 remission and EULAR Good Response after treatment with conventional synthetic DMARDs.

Disclosure of Interest: G. Cagnotto Paid instructor for: Novartis, E. Rydell: None declared, L. Jacobsson Consultant for: Pfizer, Abbvie, Novartis, Eli-Lilly, C. Turesson Grant/research support from: Abbvie, Bristol Myers-Squibb, Roche, Consultant for: MSD, Bristol Myers-Squibb, Roche, Paid instructor for: Abbvie, Bristol-Myers Squibb, Janssen, MSD, Pfizer, Roche and UCB

DOI: 10.1136/annrheumdis-2018-eular.1551

SAT0087 ASSOCIATION OF BIOMETRICS WITH DISEASE CHARACTERISTICS AND SYNOVIAL PHENOTYPE IN INFLAMMATORY ARTHRITIS

C. Low, R. Conway, F. Young, K. Murray, H. Convery, M. Biniecka, U. Fearon, D. Veale. Centre for Arthritis and Rheumatic Disease, St. Vincent's University Hospital, Dublin Academic Medical Centre, Elm Park, Dublin 4, Ireland, Dublin, Ireland

Background: The potential importance of altered body composition in the development of and disease course in inflammatory arthritis is increasingly being recognised. Body composition in different types of inflammatory arthritis and its influence on synovial pathology remains to be fully characterised.

Objectives: To evaluate body composition in seropositive and seronegative rheumatoid arthritis (RA) and psoriatic arthritis (PsA) patients and assess associations with disease characteristics and baseline synovial arthroscopic findings.

Methods: We performed a prospective observational study of consecutive inflammatory arthritis patients seen in outpatient clinics. Demographic and clinical characteristics were collected on all patients. Synovial biopsy was performed by needle arthroscopy, and macroscopic and histologic features recorded. The degree of synovitis and vascularity were recorded on a 0–100 mm visual analogue scale, and chondropathy on a semi-quantitative scale from 0–3. Mann-Whitney U test was used to compare groups. Spearman's Rank Correlation Coefficient was used to assess for associations between biometrics and demographic and clinical markers. GraphPad Prism Version 7 and IBM SPSS Statistics Version 24 were used for data analysis.

Results: We included 58 patients, 32 with seropositive RA, 10 with seronegative RA, and 16 with PsA. 37 (64%) were female. Mean (SD) age was 52.8 (13.9) years. Mean (SD) BMI was 29.7 (6.3) kg/m², waist circumference was 94.4 (20.3) cm, and hip circumference 104.3 (21) cm. Full demographic and clinical details are shown in table 1. Seronegative RA patients had significantly increased BMI (p=0.033) and waist circumference (p=0.017), but not hip circumference (p=0.248) compared to seropositive RA patients. PsA patients had significantly increased BMI (p<0.001), waist circumference (p=0.001), and hip circumference (p<0.001) compared to seropositive but not seronegative RA patients. There was a significant correlation between waist circumference and both synovitis (r=0.31, p=0.018) and vascularity (r=0.34, p=0.010) at arthroscopy. BMI and hip circumference did not correlate with arthroscopic findings.

	Seropositive RA	Seronegative RA	PsA
Female, n (%)	18 (56.3)	7 (70)	12 (75)
Age, mean (SD), years	55.9 (14.6)	46.7 (15.8)	50.6 (9.6)
Rheumatoid Factor+, n (%)	31 (96.9)	0 (0)	3 (18.8)
ACPA+, n (%)	24 (75)	0 (0)	2 (12.5)
BMI, mean (SD), kg/m ²	26.9 (4.8)	31.9 (7.1)	34.0 (5.8)
Waist circumference, mean (SD), cm	86.4 (20.4)	101.3 (16.3)	105.9 (15.4)
Hip circumference, mean (SD), cm	98.4 (23.9)	105.8 (16.4)	115.4 (11.1)
CRP, mean (SD), mg/L	12.8 (17.8)	18.6 (27.1)	7.5 (6.0)
ESR, mean (SD), mm/hr	28.2 (24.5)	18.9 (19.7)	17.5 (8.8)
Swollen Joint Count, mean (SD)	4.1 (5.0)	2.4 (3.1)	3.2 (6.1)
Tender Joint Count, mean (SD)	6.2 (6.8)	7.5 (7.5)	4.4 (7.1)
VAS general health, mean (SD), mm	49.1 (31.0)	51.3 (21.3)	49.1 (25.2)
DAS28-CRP, mean (SD)	3.91 (1.54)	3.98 (0.94)	3.6 (1.4)
Synovitis, mean (SD), mm	65.3 (20.3)	60.0 (14.1)	75.0 (12.6)
Vascularity, mean (SD), mm	62.5 (19.8)	58.0 (17.5)	75.0 (13.2)
Chondropathy, mean (SD)	1.6 (0.7)	1.4 (0.6)	1.5 (0.7)

Conclusions: Different types of inflammatory arthritis have distinct body composition profiles. Waist circumference, but not other biometrics, correlates with baseline synovial inflammation and vascularity.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2018-eular.5796

SAT0088 PREDICTIVE FACTORS FOR INTERSTITIAL LUNG DISEASE PROGRESSION IN PATIENTS WITH RHEUMATOID ARTHRITIS: A ROLE FOR BIOLOGICAL INFLAMMATION AND DISEASE MODIFIED ANTI RHEUMATIC DRUGS (DMARDs)

C. Lucas¹, S. Jouneau², M. Lederlin³, S. Guillot⁴, C. Salliot⁵, L. Joffres⁶, E. Solau-Gervais⁶, M. Herbet⁷, A. Sarau⁷, J.-B. Cren⁸, B. Bouvard⁸, J.-D. Albert¹, G. Coiffier¹, P. Guggenbuhl¹, A. Perdriger¹. ¹Rheumatology department, University Hospital Center; ²Pneumology department, University Hospital Center; ³Radiology department; ⁴Pneumology department, University Hospital Center, Rennes; ⁵Rheumatology department, Hospital Center, Orléans; ⁶Rheumatology department, University Hospital Center, Poitiers; ⁷Rheumatology department, University Hospital Center, Brest; ⁸Rheumatology department, University Hospital Center, Angers, France

Background: Interstitial Lung Disease (ILD) related to Rheumatoid Arthritis (RA) is frequent. ILD is associated with an increased mortality in RA patients. Predictive factors for ILD progression are not well studied.

Objectives: In RA patients with an ILD according to CT scan criteria, identify clinical and biological predictive factors for ILD progression.

Methods: It is a retrospective multicentric study. RA patients with ILD confirmed by a first thoracic CT scan (tCT at T0) were included if ILD progression could be studied with a second CT scan (Tx) done at least 6 months after T0. RA patients were classified in two groups after double tCT evaluation on a blind review of data: those with ILD progression (pILD) and those with a stable ILD (sILD). Predictive factors for tCT ILD progression were studied by comparing these two groups.