Immune checkpoint inhibitor rechallenge in patients with immune-related myositis

Therapeutic management of many cancers has been revolutionised by the development of immune checkpoint inhibitors (ICI) targeting antiprogrammed death 1 (PD-1)/ligand 1 (PDL1) and anticytotoxic T-lymphocyte antigen 4 leading to durable responses.¹ ICIs however can induce several immune-related adverse events (irAE) including musculoskeletal irAEs.² Among them, ICI-related myositis can be severe and sometimes life threatening.^{3,4} The current management includes permanent discontinuation of ICIs and steroid treatment. To date, very little is known about the risk of irAE recurrence in case of ICI rechallenge,^{5,6} especially in myositis for which no case of rechallenge has yet been reported. Through two cases, we report the safety of resuming anti-PD-1/PDL1 in patients who experienced severe ICI-related myositis.

An 87-year-old patient with metastatic Merkel cell carcinoma (MCC) received avelumab as first-line treatment. After three infusions, he developed slight head dropped syndrome with increased creatine kinase (CK) level up to 3.5 times the upper

limit normal (ULN) range. Electromyography showed myogenic syndrome (left trapezius and right sternocleidomastoid muscles) and ¹⁸F-fluorodeoxyglucose-positron emission tomography revealed significant hypermetabolism of axial muscles (table 1). Myositis-specific autoantibodies were negative. Myocarditis was ruled out. At that time the radiologic evaluation showed a partial tumour response. Avelumab was stopped and the patient received prednisone (tapering from 0.5 mg/kg) during 6 weeks, which allowed myositis remission, but MCC recurred 7 months later. Avelumab was resumed and prednisone was preventively given during 3 months, starting at 20 mg/day. With a 9-month follow-up, no irAE, including myositis, occurred and MCC returned in partial response.

A 61-year-old patient with metastatic melanoma developed ptosis, diplopia, dysphagia and muscle weakness 3 weeks after first infusion of ipilimumab combined with nivolumab as first-line treatment. CK levels raised up to 40 ULN. Electromyography showed myogenic pattern of the trapezius, without decrement. Muscular biopsy with focal necrosis/regeneration lesions, HLA-1 and C5b9 positive sarcoplasmic staining of the suffering myofibres and T cell infiltrates confirmed the myositis.

Table 1 Patient characteristics

		Patient 1	Patient 2
Cancer history	Stage IV cancer ICI	Merkel cell carcinoma Avelumab 10 mg/kg/2 weeks, 3 infusions	Melanoma Ipilimumab 3 mg/kg+nivolumab 1 mg/kg, 1 infusion
ICI-related myositis	Onset of symptoms	Week 6	Week 3
	Clinical symptoms	Dropped head syndrome Neck pain Fatigue	Myalgia, muscle weakness Ptosis, diplopia, dysphagia Fatigue
	Maximum	×3.5 ULN	×40 ULN
	Creatine kinase		
	Electromyography	Myogenic syndrome: Left trapezius and right sternocleidomastoid muscle. No decrement.	Myogenic syndrome: Right trapezius. No decrement.
	FDG-PET	Significant hypermetabolism of trapezius, erector muscles of the spine, pillar muscle of the diaphragm	Slight diffuse muscular hypermetabolism
	Muscular biopsy	Contraindication (anticoagulant treatment for atrial fibrillation)	Positive with focal necrosis/regeneration lesions, HLA-I and C5b9 positive staining, presence of T cell and macrophage inflammatory cells
	Cardiac examination ECG	Permanent atrial fibrillation (>5 years)	Normal
	High-sensitive troponin-T (<14 ng/L)	475	298
	Echocardiography	Normal	Normal
	Cardiac MRI	Contraindication (pacemaker)	—
	Cardiac FDG PET	No myocarditis	No myocarditis
	Treatment of irAE	Oral prednisone 0.5 mg/kg/day, tapered and withdrawn within 6 weeks	Methylprednisolone 1 g/day×3 days, followed by oral prednisone 1 mg/kg, tapered and withdrawn within 9 weeks
Rechallenge	Response to initial treatment with ICI	Partial response	Progressive disease
	Time from first ICI treatment to ICI rechallenge	7 months	8 months
	ICI	Avelumab 10 mg/kg/2 weeks, 20 infusions	Pembrolizumab 2 mg/kg/3 weeks, 2 infusions
	Associated treatment	Prednisone 20 mg/day, tapered and withdrawn within 3 months	—
	irAE	None	None
	ICI-myositis	No recurrence of myositis with no symptoms, normal CK level, normal electromyography and absence of muscle hypermetabolism on TEP	No recurrence of myositis without any symptoms, normal CK level
	Tumour response	Ongoing partial response after 9 months	Death due to progressive disease at week 6

CK, creatine kinase; FDG-PET, fluorodeoxyglucose-positron emission tomography; HLA, human leucocyte antigen; ICI, immune checkpoint inhibitor; irAE, immune-related adverse event; ULN, upper limit normal.

Correspondence

Neither myositis-specific nor myasthenia gravis autoantibodies were detected. ICIs were stopped and three pulses of methylprednisolone followed by tapering doses of prednisone were given leading to complete remission within 8 weeks. Because of the lack of efficacy of the single infusion of ICI combination followed by three infusions of dacarbazine, pembrolizumab was introduced 8 months after the myositis episode. The patient presented no irAEs or myositis (table 1) but died due to melanoma progression.

Despite the risk of recurrent irAEs, rechallenging ICIs after discontinuation due to previous irAEs remains critical, when considering their potential benefits in terms of survival.⁵ These two cases suggest that resuming ICIs can be safe in patients displaying persistent remission of ICI-related myositis.

Julie Delyon,^{1,2,3} Florence Brunet-Possenti,^{3,4} Sarah Leonard-Louis,^{5,6} Dimitri Arangalage,^{3,7,8} Mathilde Baudet,⁹ Barouyr Baroudjian,¹ Celeste Lebbe,^{1,2,3} Baptiste Hervier,^{6,10,11} on behalf of the PATIO Group

¹Department of Dermatology, APHP Saint Louis Hospital, Paris, France

²INSERM U976, Paris, France

³Paris Diderot, Sorbonne Paris Cité University, Paris, France

⁴Department of Dermatology, APHP Bichat Hospital; Paris VII Diderot, Sorbonne Paris Cité University, Paris, France

⁵Department of Neurology and Neuropathology, APHP Pitié-Salpêtrière Hospital, Paris, France

⁶UPMC, Médecine Sorbonne Université, Paris, France

⁷Cardiology Department, APHP Bichat Hospital, Paris, France

⁸INSERM U1148, Paris, France

⁹Department of Cardiology, APHP Saint Louis Hospital, Paris, France

¹⁰Department of Internal Medicine, APHP Pitié-Salpêtrière Hospital, Paris, France

¹¹INSERM U1135-CIMI, Paris, France

Correspondence to Dr Julie Delyon, Dermatology, APHP Saint Louis Hospital, Paris 75475, France; julie.delyon@aphp.fr

Contributors JD and FBP analysed the data and drafted the manuscript for intellectual content. SLL, DA, MB and BB had a major role in the acquisition of data. CL and BH interpreted the data and revised the manuscript for intellectual content. All authors approved the submitted version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests BH, DA, JD, FBP, MB and SLL had nothing to declare. BB received payments for lectures/boards and travel accommodations from BMS and MSD. CL received research grants or honoraria from Roche, BMS, MSD, GSK, Novartis and Amgen.

Patient consent Obtained.

Ethics approval Local ethics committee.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Delyon J, Brunet-Possenti F, Leonard-Louis S, et al. *Ann Rheum Dis* 2019;**78**:e129.

Received 24 August 2018

Accepted 30 August 2018

Published Online First 21 September 2018

Ann Rheum Dis 2019;**78**:e129. doi:10.1136/annrheumdis-2018-2-14336

REFERENCES

- 1 Wolchok JD, Chiarion-Sileni V, Gonzalez R, et al. Overall survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 2017;377:1345–56.
- 2 Kostine M, Rouxel L, Barnette T, et al. Rheumatic disorders associated with immune checkpoint inhibitors in patients with cancer—clinical aspects and relationship with tumour response: a single-centre prospective cohort study. *Ann Rheum Dis* 2018;77:393–8.
- 3 Touat M, Maisonneuve T, Knauss S, et al. Immune checkpoint inhibitor-related myositis and myocarditis in patients with cancer. *Neurology* 2018;91:e985–94.
- 4 Eggermont AMM, Blank CU, Mandala M, et al. Adjuvant pembrolizumab versus placebo in resected stage III melanoma. *N Engl J Med* 2018;378:1789–801.
- 5 Pollack MH, Betof A, Dearden H, et al. Safety of resuming anti-PD-1 in patients with immune-related adverse events (irAEs) during combined anti-CTLA-4 and anti-PD1 in metastatic melanoma. *Ann Oncol* 2018;29:250–5.
- 6 Menzies AM, Johnson DB, Ramanujam S, et al. Anti-PD-1 therapy in patients with advanced melanoma and preexisting autoimmune disorders or major toxicity with ipilimumab. *Ann Oncol* 2017;28:368–76.