

Response to: 'M-CSF and GM-CSF monocyte-derived macrophages in systemic sclerosis: the two sides of the same coin?' by Lescoat *et al*

In the quest for identification of genes for complex disorders, the functional annotation of disease-risk variants in the relevant cell type remains a challenge. Lescoat and colleagues¹ pose the question 'what is a scleroderma macrophage?' and in view of current literature (including the study by Moreno-Moral *et al*²), argued how the heterogeneous range of activated macrophages can vary between target organs of interest and possibly across different fibrotic diseases.

While we share Lescoat's point of view on the underlying complexity of functional cells in disease, it remains open to debate whether GM-CSF or M-CSF-mediated differentiation conditions of peripheral blood monocytes are sufficient to fully capture the characteristics of these cells in their disease context. As stated by Lescoat *et al*,¹ whether and to which extent the in vitro defined M1/M2 macrophage subtypes reflect the context-specific activation of macrophages observed in different organs (eg, skin, lung) in scleroderma remains to be demonstrated. In this respect, the proposed spectrum model of human macrophage activation³ argues in favour of a scleroderma-specific macrophage activation profile which does not fall into the binary M1/M2 categorisation.¹

The experimental challenges encountered in finding the 'best' cell culture conditions for modelling disease could be eased by investigating the genetic control of gene expression (ie, expression QTL or eQTL). Theoretically, these studies can test, regardless of culturing conditions, whether certain DNA variants affect the expression levels of core susceptibility genes expressed by myeloid cells, in general. Recent eQTL studies in humans showed context-specific variants (ie, on stimulation) in monocytes and macrophages but there is also evidence of some transcripts which are regulated by DNA sequence variation (eg, SNPs) independently of the stimulus⁴ or the differentiation state of the cell (monocyte vs macrophage). For instance, Fairfax *et al* identified *GSDMA* as an eQTL in activated monocytes⁴ and Moreno-Moral *et al.* (figure 2 in reference 2) showed the same gene being under local genetic control in M-CSF-derived macrophages isolated from patients with scleroderma. Notably, *GSDMA* associates with asthma susceptibility⁵ and scleroderma.⁶ This suggests genotype-dependent *GSDMA* expression in a wider monocyte/macrophage compartment, which is likely to be a risk component in these diseases. It is therefore plausible that the genetic control of *GSDMA* mRNA levels will be conserved in both M-CSF and GM-CSF-derived scleroderma macrophages but further experiments will be required to reach a definite conclusion.

Interestingly, human monocytes share eQTLs with CD4+ T cells⁷ and neutrophils,⁸ indicating that common DNA variants regulate mRNA levels of target genes in multiple immune cell types, suggesting the possibility of disease risk variants impacting on the wider immune response. As highlighted by Lescoat *et al*,¹ whether the effect of particular disease susceptibility variants on a gene's mRNA level is also conserved at the protein level remains to be examined, and we reckon that additional layers of regulation (eg, post-transcriptional, translational) are likely to be at work. Still, there are examples of robust conservation

between expression and protein QTLs associated with disease.⁹ Ultimately, the eQTL approach, while informative in linking *bona fide* disease risk variants with target cells (as shown by Moreno-Moral *et al*²), should be considered in the broader target tissue context rather than in specific cell culture conditions,¹⁰ and recent advances in ATAC-seq, single cell RNA-seq will allow the refinement of causal variants for systemic sclerosis in a setting closer to their in vivo context.

Jacques Behmoaras,¹ Enrico Petretto^{2,3}

¹Department of Medicine, Imperial College London, Hammersmith Hospital, London, UK

²Faculty of Medicine, Medical Research Council (MRC) London Institute of Medical Sciences, Imperial College London, London, UK

³Centre for Computational Biology, Duke-NUS Medical School, Singapore, Singapore

Correspondence to Dr Jacques Behmoaras, Department of Medicine, Imperial College London, Hammersmith Hospital, London W12 0NN, UK; jacques.behmoaras@imperial.ac.uk and Professor Enrico Petretto; enrico.petretto@duke-nus.edu.sg

Handling editor Josef S Smolen

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2019. All rights reserved. No commercial use is permitted unless otherwise expressly granted.



To cite Behmoaras J, Petretto E. *Ann Rheum Dis* 2019;**78**:e20.

Accepted 2 March 2018

Published Online First 9 March 2018



► <http://dx.doi.org/10.1136/annrheumdis-2018-213121>

Ann Rheum Dis 2019;**78**:e20. doi:10.1136/annrheumdis-2018-213121

REFERENCES

- Lescoat A, Jégo P, Lecœur V. M-CSF and GM-CSF monocyte-derived macrophages in systemic sclerosis: the two sides of the same coin? *Ann Rheum Dis* 2019;**78**:e20.
- Moreno-Moral A, Bagnati M, Koturan S, *et al.* Changes in macrophage transcriptome associate with systemic sclerosis and mediate *GSDMA* contribution to disease risk. *Ann Rheum Dis* 2018;**77**:596–601.
- Xue J, Schmidt SV, Sander J, *et al.* Transcriptome-based network analysis reveals a spectrum model of human macrophage activation. *Immunity* 2014;**40**:274–88.
- Fairfax BP, Humburg P, Makino S, *et al.* Innate immune activity conditions the effect of regulatory variants upon monocyte gene expression. *Science* 2014;**343**:1246949.
- Marinho S, Custovic A, Marsden P, *et al.* 17q12-21 Variants are associated with asthma and interact with active smoking in an adult population from the United Kingdom. *Annals of Allergy, Asthma & Immunology* 2012;**108**:402–11.
- Terao C, Kawaguchi T, Dieude P, *et al.* Transethnic meta-analysis identifies *GSDMA* and *PRDM1* as susceptibility genes to systemic sclerosis. *Ann Rheum Dis* 2017;**76**:1150–8.
- Raj T, Rothamel K, Mostafavi S, *et al.* Polarization of the effects of autoimmune and neurodegenerative risk alleles in leukocytes. *Science* 2014;**344**:519–23.
- Naranbhai V, Fairfax BP, Makino S, *et al.* Genomic modulators of gene expression in human neutrophils. *Nat Commun* 2015;**6**:7545.
- Ryan KJ, White CC, Patel K, *et al.* A human microglia-like cellular model for assessing the effects of neurodegenerative disease gene variants. *Sci Transl Med* 2017;**9**:eaai7635.
- Ongen H, Brown AA, Delaneau O, *et al.* Estimating the causal tissues for complex traits and diseases. *Nat Genet* 2017;**49**:1676–83.