Response to: 'Association between use of non-steroidal anti-inflammatory drugs and risk of myocardial infarction in patients with spondyloarthritis and osteoarthritis'

We wish to thank Zhou and colleagues for their letter.¹ As outlined in the published paper,² the myocardial infarction (MI) case definition was based on the diagnosis being recorded as a MI Read code in The Health Improvement Network (THIN), a validated means of identifying MI in pharmacoepidemiological studies.³ Because THIN is based on general practitioner's coded records, the additional information Zhou *et al* suggested¹ is not systematically available. This was already acknowledged in the discussion section as a limitation, and we additionally described our internal MI validation study in the discussion section, in which 89% of MI cases had an administrative code supporting the occurrence of an MI; these codes included a hospitalisation, having had an ECG or angiogram performed and a referral to a cardiologist. Similarly, the definition of osteoarthritis is also outlined in the Methods section.

We recognise the importance of considering age as an effect modifier related to non-steroidal anti-inflammatory drug (NSAID) use and attempted to partly address this through the sensitivity analysis restricted to subjects aged 55–70.

The most frequent NSAID prescriptions are described in online supplementary table 1 from our paper.² The topic of NSAID dosage and frequency certainly deserves further study.

Finally, we are not aware of data demonstrating a relation of chondroitin to risk of MI. Systemic glucocorticoids are not recommended for the treatment of knee or hip osteoarthritis nor for spondyloarthritis according to the most recent American College of Rheumatology and European League Against Rheumatism guidelines; therefore, we expect low prevalence of use among subjects in this study. Further, these medications would be considered to be intermediates in the causal pathway, and therefore adjustment for any of those types of medications would induce bias.

Maureen Dubreuil,¹ Qiong Louie-Gao,¹ Christine Peloquin,¹ Hyon K Choi,² Yuqing Zhang,¹ Tuhina Neogi¹

¹Department of Medicine, Clin Epi Research and Training Unit, Boston University School of Medicine, Boston, Massachusetts, USA

²Department of Rheumatology, Harvard Medical School, Boston, Massachusetts, USA

Correspondence to Dr Maureen Dubreuil, Department of Medicine, Clin Epi Research and Training Unit, Boston University School of Medicine, Boston, MA 02118, USA; mdubreuil@bu.edu

Handling editor Josef S Smolen

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2019. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

To cite Dubreuil M, Louie-Gao Q, Peloquin C, *et al.* *Ann Rheum Dis* 2019;**78**:e79.

Received 23 May 2018

Revised 24 May 2018

Accepted 26 May 2018

Published Online First 16 June 2018

► <http://dx.doi.org/10.1136/annrheumdis-2018-213752>

Ann Rheum Dis 2019;**78**:e79. doi:10.1136/annrheumdis-2018-213771

REFERENCES

- 1 Zhou J, Xiao Y, Li C, *et al.* Association between use of non-steroidal anti-inflammatory drugs and risk of myocardial infarction in patients with spondyloarthritis and osteoarthritis. *Ann Rheum Dis* 2019;**78**:e78.
- 2 Dubreuil M, Louie-Gao Q, Peloquin CE, *et al.* Risk of myocardial infarction with use of selected non-steroidal anti-inflammatory drugs in patients with spondyloarthritis and osteoarthritis. *Ann Rheum Dis* 2018;**77**:1137–42.
- 3 Lewis JD, Schinnar R, Bilker WB, *et al.* Validation studies of the health improvement network (THIN) database for pharmacoepidemiology research. *Pharmacoepidemiol Drug Saf* 2007;**16**:393–401.