not only professional bone-resorbing cells but also directly involved in controlling inflammatory responses. Depending on the pathophysiological context, OC can polarize the immune response towards tolerance or inflammation. Aiming at identifying key regulators of inflammatory OC functions, we have defined a miRNA-based signature, which includes miR-342-3p. MiRNAs are key regulators of gene expression that control cellular processes, including osteoclastogenesis, and few miRNAs have been described in the differentiation of myeloid precursors into mature OC.

Objectives To study the role of miR-342-3p in inflammatory OC.

Methods OC were derived from the murine monocyte RAW264.7 cells. The expression levels of miR-342-3p and OC-specific genes were monitored by qRT-PCR. RAW264.7 cells were transfected with either miR-342-3p mimics, neutralizing molecules or control miRNAs. Cell survival and proliferation were assessed at 48 hour after RANKL incubation using quantification of the caspase3-7 activity, ATP production and BrDU incorporation. Motility of OC precursors was monitored using time-laps during the course of OC differentiation. The K/BxN serum-transfer arthritis (STA) model was performed in 8 weeks old C57BL/6 males and bone marrow was flushed. Primary OC were generated from either total bone marrow or sorted CD11b⁺ and CD11c⁺ cell subsets of healthy and arthritic mice.

Results The expression of miR-342-3p was transiently up-regulated in the early phase of OC generation and was down-regulated after 24–48 hour in OC precursors. While pre-miR-342-3p promoted the motility of RAW264.7 cells, anti-miR-342-3p inhibited all motility parameters recorded ($p < 0.0001$, ANOVA test). Anti-miR-342-3p reduced the proliferation ($p < 0.01$) and cell survival of OC precursors through a pro-apoptotic effect, as assessed by increased caspase3-7 activity ($p < 0.01$). Overall, miR-342-3p neutralization in OC precursors reduced OC numbers ($p < 0.001$) compared to the pre-miR-342-3p condition. In primary cells, miR-342-3p was up-regulated in bone marrow-derived mature OC from arthritic mice compared to healthy controls ($p = 0.03$; STA $n = 5/\text{group}$).

Conclusions Our data suggest that miR-342-3p promotes the early phase of osteoclastogenesis by enhancing the cell survival and motility of OC precursors. The up-regulation of miR-342-3p in OC isolated from arthritic mice may reflect the increased osteoclastogenic potential of inflammatory precursors in arthritis.

Disclosure of Interest None declared.

P067

TOFACITINIB IMPAIRS MONOCYTE-DERIVED DENDRITIC CELL DIFFERENTIATION IN RHEUMATOID ARTHRITIS AND PSORIATIC ARTHRITIS

¹V Marzaioli*, ¹M Canavan, ¹A Floudas, ¹S Wade, ²C Low, ²DJ Veale, ¹U Fearon. ¹TBSI, Molecular Rheumatology, Trinity College Dublin; ²Centre for Arthritis and Rheumatic Diseases, St Vincent's University Hospital, UCD, Dublin, Ireland

10.1136/annrheumdis-2018-EWRR2019.56

Career situation of first and presenting author Post-doctoral fellow.

Introduction Tofacitinib is an oral Janus kinase inhibitor, recently approved for the treatment of rheumatoid arthritis (RA) and psoriatic arthritis (PsA). Although its mechanism of action has been explored in circulating cells, including neutrophils and lymphocyte, its effect on dendritic cells development

and function remains still to be elucidated. Monocyte-derived dendritic cells (Mo-DC) are a subset of inflammatory DC derived from circulating monocytes with a key role in inflammation and infection.

Objectives The aim of this project is to evaluate the effect of Tofacitinib on inflammatory Mo-DC differentiation from RA and PsA patients, an important step in innate immunity.

Methods Monocytes were isolated from blood of healthy donor (HC), RA and PsA patients by magnetic separation and differentiated in the presence of GM-CSF/IL-4 cocktail for 7 days. To evaluate the effect of Tofacitinib on Mo-DC differentiation, monocyte were pre-treated with 1 μM Tofacitinib (or DMSO as control). CD209 (immature DC marker) was evaluated by flow cytometry in the CD11c⁺ population. Non-specific macropinocytosis (using Lucifer Yellow) and receptor-mediated endocytosis (using DQTM Ovalbumin) were investigated by flow cytometry. The effect of Tofacitinib on NADPH oxidases (NOX) 5 and 2 expression, known players in Mo-DC differentiation, was evaluated by Western blot analysis. Finally, the frequency of CD209⁺ cells and their chemokine receptor expression (CXCR3/4/5 and CCR6/7) were evaluated by flow cytometry in peripheral blood (PBMC), synovial fluid (SFMC) mononuclear cells and synovial tissue cell suspensions from RA and PsA patients.

Results Pre-treatment of Mo-DC with Tofacitinib inhibited Mo-DC differentiation in RA and PsA patients, as evident by reduced CD209 marker expression. The decreased ability of monocytes to differentiate into DC in the presence of Tofacitinib was translated into a functional impairment of endocytic ability, in particular in PsA patients, as observed by the decreased uptake of both DQTM Ovalbumin and Lucifer Yellow. In addition, Tofacitinib decreased NOX5 and increased NOX2 protein expression in Mo-DC of PsA and RA Mo-DC, altering the NOX2/NOX5 balance. Finally, we identified CD209⁺ cells in the periphery of RA and PsA patients, which were enriched in SFMC and synovial tissue cell suspension, and presented with an increased expression of CCR7 and CXCR3/5.

Conclusions Together, these observations suggest a novel mechanism of action of Tofacitinib in RA and PsA, by inhibiting Mo-DC development, which may alter migration of DC to the joint and subsequent activation of the immune response.

Disclosure of Interest V. Marzaioli Grant/research support from: Pfizer, M. Canavan: None declared, A. Floudas: None declared, S. Wade: None declared, C. Low: None declared, D. Veale: None declared, U. Fearon Grant/research support from: Pfizer.

P068

RHEUMATOID ARTHRITIS PERIPHERAL CD14⁺ MONOCYTES ARE HYPER-INFLAMMATORY, HYPER-GLYCOLYTIC AND RETAIN A MEMORY BIAS TOWARD M1 MACROPHAGES

¹T McGarry*, ¹M Hanlon, ¹U Fearon, ²DJ Veale. ¹Clinical Medicine, Trinity Biomedical Sciences Institute; ²Centre for Arthritis and Rheumatic Diseases, St. Vincent's University Hospital, Dublin, Ireland

10.1136/annrheumdis-2018-EWRR2019.57

Career situation of first and presenting author Post-doctoral fellow.

Introduction Myeloid cells with a monocyte/macrophages phenotype are present in large numbers in the rheumatoid arthritis (RA) joint, significantly contributing to disease.

Objectives This study aimed to assess whether peripheral monocytes in RA are pre-programmed to become M1 pro-inflammatory macrophages.

Methods Blood was collected from healthy donors, at-risk individuals (Those with arthralgia, ACPA+/RF+, normal CRP and no evidence of synovitis) and established RA patients. CD14+ monocytes were isolated from peripheral blood mononuclear cells using a CD14 magnetic bead separation kit. Cells were stimulated with LPS (100 ng/ml) for 3–24 hours and to assess the effects of STAT3 inhibition, cells were pre-treated with STAT3i (10 μ M) for 30mins. A Human Cytokine and Chemokine PCR array was carried out and those genes most differentially expression were further validated in a larger cohort of patients using RT-PCR. The metabolic profile of cells was analysed using Seahorse XFE Technology, which concomitantly analysis glycolysis and mitochondrial respiration in real-time. Gene and protein expression of key inflammatory and glycolytic markers was also carried out by RT-PCR, western blotting and ELISA.

Results CD14+ RA monocytes are hyper-inflammatory upon stimulation, with significantly higher expression of IL-1 β , TNF α , IL-6, IL-27, CXCL10 and CXCL11 compared to healthy controls, which is indicative of a M1-like pro-inflammatory phenotype. These hyper-inflammatory cells are highly glycolytic, with increased expression of HIF1 α , HK and PFKFB3, key glycolytic enzymes. Both baseline glycolysis and baseline oxidative phosphorylation are increased in RA CD14+ monocytes, paralleled by increased ATP synthesis and maximal respiratory capacity, suggesting a hyper-energetic phenotype. This hyper-inflammatory, hyper-glycolytic phenotype is mediated by STAT3, as selective STAT3 inhibition can significantly decrease M1-like cytokines and PFKFB3 and HK2 expression. In addition, STAT3 inhibition significantly decreases both oxidative phosphorylation and glycolysis pathways. Finally, this pro-inflammatory phenotype is evident in CD14+ monocytes from arthralgia ACPA+/RF+ people at risk of developing disease, demonstrating that these processes may precede clinical manifestations in RA.

Conclusions This study demonstrates the unique inflammatory and metabolic phenotype of RA monocytes, suggesting that peripheral CD14+ monocytes may be pre-programmed to become M1-like pro-inflammatory macrophages. In addition, the observation of this phenotype in at-risk individuals indicates that these features may precede clinical manifestations of RA and therefore could be useful as a biomarker for early diagnosis.

Disclosure of Interest None declared.

P069

CONSTITUTIVE OVEREXPRESSION OF INTERLEUKIN 38 HAS A NEGATIVE IMPACT ON HUMAN NHK KERATINOCYTE FITNESS

L Mermoud*, D Talabot-Ayer, N Brembilla, J Julia Borowczyk-Michalowska, G Palmer. Department of Pathology-Immunology, University of Geneva School of Medicine, Geneva, Switzerland

10.1136/annrheumdis-2018-EWRR2019.58

Career situation of first and presenting author Student for a master or a PhD.

Introduction Interleukin (IL)-38 is a member of the IL-1 cytokine family. Based on its sequence homology with the IL-1 and IL-36 receptor antagonists, IL-38 was proposed to act as an anti-inflammatory molecule. However, few data are

available to date about its biological role and mechanism of action. Highest constitutive IL-38 expression is detected in the skin, where the cytokine is produced mainly by differentiated keratinocytes.

Objectives The aim of this study is to investigate the function of IL-38 in human keratinocytes, using the immortalized normal human keratinocyte (NHK) cell line stably transfected with vectors allowing for constitutive or inducible IL-38 overexpression.

Methods NHK cell proliferation and mortality were assessed respectively by cell counting and by measuring LDH activity in the supernatant of high density cell culture. Differentiation of NHK cells was initiated by switching to a culture medium containing high Ca⁺⁺ and serum. mRNA levels of differentiating NHK cells were determined by RTqPCR.

Results Constitutive IL-38 overexpression reduced the proliferation of NHK cells and lead to increased mortality. Differentiating NHK cells constitutively overexpressing IL-38 had increased mRNA levels for the early differentiation marker keratin 10 as compared to empty vector transfected control cells. On the contrary, expression of the late differentiation marker involucrin was reduced in NHK cells constitutively overexpressing IL-38. To further explore potential effects of IL-38 on keratinocyte differentiation, while circumventing the impact of its constitutive overexpression on cell fitness, we set up a doxycyclin-inducible Tet-On system to overexpress IL-38 in NHK cells.

Conclusions We are currently optimizing 2D and 3D culture conditions for *in vitro* differentiation of NHK cells to mimic more closely the situation in human skin. We will then study the impact of IL-38 on NHK cell differentiation using cells transfected with the inducible IL-38 expression system. In conclusion, our results suggest that constitutive overexpression of IL-38 decreases proliferation and viability of NHK cells. We will thus use an inducible expression system to further investigate the effects of IL-38 on NHK cells.

Disclosure of Interest None declared.

P070/O09

MIR-221-3P IS UPREGULATED IN RHEUMATOID ARTHRITIS AND AFFECTS JAK3/STAT3 SIGNALING IN M2-MACROPHAGES

^{1,2}L Quero*, ^{1,2}E Hanser, ^{1,2}AN Tiaden, ^{1,2}D Kyburz. ¹DBM, University of Basel; ²Experimental Rheumatology, University Hospital, Basel, Switzerland

10.1136/annrheumdis-2018-EWRR2019.59

Career situation of first and presenting author Post-doctoral fellow.

Introduction Recurrent activation of synovial macrophages is considered as intrinsic factor that precedes rheumatoid arthritis (RA)-related autonomous cytokine cascades. microRNAs (miRs) and Toll-like receptors (TLRs) play a pivotal role in regulating the inflammatory response in innate immune cells and are at dysregulated levels in RA patients.

Objectives Here we explored TLR-regulated miRs that tune the inflammatory response of M2-type macrophages.

Methods Based on a miR-screen in TLR-stimulated human M2-macrophages we selected miR-221-3p to analyze its expression profile in clinical samples from RA, other inflammatory arthritis (OIA), osteoarthritis (OA) and healthy donors (HD) by qPCR. We challenged M1- and M2-macrophages with elevated miR-221-3p and/or miR-155-5p expression