indicate HCQ may be used as a strategy against intestinal mucositis during 5-FU chemotherapy.

REFERENCES

1. Woo S-R, Fuertes MB, Corrales L, *et al.* STING-Dependent Cytosolic DNA Sensing Mediates Innate Immune Recognition of Immunogenic Tumors. *Immunity* 2014;**41**:830–42.
2. Hu B, Jin C, Li H-B, *et al.* The DNA Sensing Aim2 Inflammasome Controls Radiation Induced Cell death and Tissue Injury. *Science (New York, NY)*. 2016;**354**:765–8.
3. Miura K, Kodama Y, Inokuchi S, *et al.* Toll-Like Receptor 9 Promotes Steatohepatitis by Induction of Interleukin-1 β in Mice. *Gastroenterology*. 2010;**139**:323–34. e7.

Acknowledgements The authors confirm that there are no conflicts of interest.

Disclosure of Interest None declared.

P146 GOLIMUMAB IN RHEUMATOID AND PSORIATIC ARTHRITIS. THE RELATIONSHIP BETWEEN MINIMUM SERUM CONCENTRATION, ANTI-GOLIMUMAB ANTIBODIES AND ACTIVITY OF DISEASE

¹J Uceda*, ²MJ Fobelo, ³R Hernandez. ¹Rheumatology Unit, ²Pharmacy Unit, University Valme Hospital; ³Rheumatology Unit, University Valme Hospital, Seville, Spain

10.1136/annrheumdis-2018-EWRR2019.129

Career situation of first and presenting author Young investigator.

Introduction Currently there is very little information on optimal levels of golimumab in patients with polyarthritis.

Objectives The main objective was to analyze the minimal serum concentration of golimumab (GOL), anti-golimumab antibodies (AcGOL) and its relation to the activity of the disease measured by DAS 28 index, in rheumatoid arthritis (RA) and psoriatic arthritis (PA).

Methods We conducted a retrospective observational study with RA or PA patients on golimumab treatment between 2011 and 2016.

We analyzed the age, sex, diagnosis, dose, dose interval, duration of treatment, concomitant disease modifying drugs (DMARDs), minimum concentration of GOL (GOL Cmin), AcGOL and DAS28 levels. Sampling was performed on the day corresponding to the dose of golimumab, prior to administration, in order to obtain the minimum levels of the drug. The analytical technique used for the determination was the Promonitor® sandwich ELISA.

Results 10 RA, 5 AP patients were selected. Median age 53 years. 67% women. All with golimumab 50 mg with an administration interval of 28 to 41 days. Duration of treatment 16.73 months (median). 80% underwent combination therapy with DMARDs.

Only one patient (RA) presented levels of AcGOL, with undetectable Cmin, DAS28 of 4.05.

Golimumab levels in patients without AcGOL and DAS28 <3.2: 1 Cmin <0.25 mcg/mL, 5 Cmin 0.26–0.5 mcg/mL, 1 Cmin 0.5–1.4 mcg/mL and 1 2mcg/mL Cmin. In patients with DAS28 >3.2 were: 2 Cmin <0.25 mcg/mL, 2 Cmin 0.26–0.5 mcg/mL, 2 Cmin 0.5–1.4 mcg/mL and 1 Cmin of 2.2 mcg/mL.

Conclusions Very variable GOL Cmin values (<0.25–2.2 mcg/mL) were obtained. 80% with DAS28 <3.2 had a GOL Cmin: 0.26–0.5 mcg/mL. Because of the limited information currently available on the optimal range of golimumab levels, further

studies are needed to evaluate the relationship between serum drug levels and patient disease activity data.

Disclosure of Interest None declared.

P147 TREATMENT WITH USTEKINUMAB IN PSORIATIC ARTHRITIS IN CONDITIONS OF REAL CLINICAL PRACTICE

¹J Uceda*, ²R Hernandez, ³JL Marengo. ¹Rheumatology Unit, University Valme Hospital; ²Rheumatology Unit, University Valme Hospital; ³Rheumatology Unit, Valme University Hospital, Seville, Spain

10.1136/annrheumdis-2018-EWRR2019.130

Career situation of first and presenting author Assistant.

Introduction Psoriatic arthritis (PsA) belongs to the family of spondyloarthritis, with an incidence of 6 cases per 1 00 000 inhabitants. Ustekinumab is an inhibitor of IL 12/13 approved for patients with PsA who do not respond to disease modifying anti-rheumatic drugs (DMARDs). This drug has been shown to be effective and safe in different clinical trials.

Objectives The primary objective was to assess the effectiveness and safety of Ustekinumab in a cohort of patients with PsA, in follow-up in the Rheumatology Unit of the Valme University Hospital, under conditions of real clinical practice. We have also studied the survival of the drug as well as treatment interruptions and their causes.

Methods We conducted an observational, descriptive, longitudinal and retrospective study of a cohort of 16 patients diagnosed with PsA, according to CASPAR criteria, who receive or have received treatment with Ustekinumab from January 1, 2015 to February 1, 2018.

The effectiveness of the drug was assessed by changing the DAS 28 at 3 and 6 months. We recorded the sociodemographic characteristics of the patients and the characteristics of the disease. Also we recorded previous treatment with anti TNF alpha, and concomitant treatment with DMARDs. We evaluated the suspension of the drug and its reasons. Adverse reactions were recorded too.

Results 16 patients were studied, 56.3% women. The average age at diagnosis was 41.38 years $\pm$ 8.5 SD. 62.5% presented polyarticular form, 12.5% oligoarticular, 6.3% axial and 18.8% axial and peripheral arthritis. 87.5% had rheumatoid factor (RF) and antipeptide-citrullinated antibodies (ACPA) negative. 87.55% had received prior treatment with at least one anti-TNF alpha. 81% of the sample received Ustekinumab in combination with a DMARDs, being methotrexate the most used.

The mean of DAS 28 at the start of treatment with Ustekinumab was 4.51 $\pm$ 1.07 SD. The mean of DAS28 was 3.11 $\pm$ 1.39 SD and 2.93 $\pm$ 1.58 SD after 3 months and 6 months of treatment.

The average survival of ustekinumab in the period analyzed (January 1, 2015 to January 1, 2018) was 29.1 months $\pm$ 3.7 SD.

5 patients of our sample discontinued the treatment: 4 due to ineffectiveness and 1 due to an adverse effect that consisted of a bronchospasm crisis, resolving itself after drug suspension and not requiring hospital admission. There has been no cases of tuberculosis, neoplasia, demyelinating disease or death.

Conclusions Ustekinumab is an effective drug for psoriatic arthritis since three months of treatment. In our study Ustekinumab presented a high survival and a good safety profile in real clinical practice.

Disclosure of Interest None declared.