Quantitative assessment: A significantly thicker cartilage was found in males [0.71 ± 0.10 mm (mean $\pm$ SD)] than in females [0.68 ± 0.12 mm, (mean $\pm$ SD)] ($p < 0.01$). No significant difference was found between left and right side for each digit ($p > 0.05$).

CT value of the II MH was significantly greater than the one of the other fingers ($p < 0.01$). No difference was found between the CT values of III, IV and V MH ($p > 0.05$).

There was a significant association between the CT value and gender ($r = 0.39$; $p < 0.01$), age ($r = -0.33$; $p < 0.01$), height ($r = 0.28$; $p = 0.01$) and grade of the semiquantitative scoring system ($r = -0.19$, $p < 0.01$). No correlation was found between the CT value and weight ($p = 0.20$).

Conclusion: This study confirms the presence and provides data on the prevalence of US abnormalities at MH level in healthy individuals. Moreover, normal values for US CT of MH were reported.

REFERENCES:

- [1] Mandl P, Supp G, Baksa G, et al. Relationship between radiographic joint space narrowing, sonographic cartilage thickness and anatomy in rheumatoid arthritis and control joints. *Ann Rheum Dis.* 2015;74:2022-7.
- [2] Torp-Pedersen S, Bartels EM, Wihjelm J, Bliddal H. Articular cartilage thickness measured with US is not as easy as it appears: a systematic review of measurement techniques and image interpretation. *Ultraschall Med.* 2011;32:54-61.
- [3] Filippucci E, da Luz KR, Di Geso L, et al. Interobserver reliability of ultrasonography in the assessment of cartilage damage in rheumatoid arthritis. *Ann Rheum Dis.* 2010;69:1845-8.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5807

FRI0635

ULTRASOUND IN THE ASSESSMENT OF TENOSYNOVITIS IN JUVENILE IDIOPATHIC ARTHRITIS: SYSTEMATIC LITERATURE REVIEW

PAZ COLLADO¹, Silvia Magni-Manzoni², Martina Steiner³, Tracy Ting⁴, Patricia Vega Fernandez⁵, Clara Malattia⁶, Ana Rodriguez⁷, George Bruyn⁸, Helen Keen⁹, Lene Terslev¹⁰, The paediatric subgroup of the OMERACT Ultrasound Task Force.

¹Hospital Universitario Severo Ochoa, Madrid, Spain; ²IRCCS Ospedale Pediatrico Bambino Gesù, Roma, Italy; ³Hospital Universitario Infanta Sofia, Madrid, Spain;

⁴Cincinnati Children's Hospital Medical Center, Cincinnati, OH, United States of America; ⁵Emory University, Children's Health Care of Atlanta, Atlanta, GA, United States of America; ⁶Istituto Giannina Gaslini, Genova, Italy; ⁷Hospital Universitario Ramon y Cajal, Madrid, Spain; ⁸MC Groep, Lelystad, Netherlands; ⁹School of Medicine and Pharmacology Fiona Stanley Hospital Unit, Perth, Australia; ¹⁰Center for Rheumatology and Spine Diseases, Copenhagen University Hospital Glostrup, Copenhagen, Denmark

Background: Tendon involvement is common in patients with juvenile idiopathic arthritis (JIA). Differentiating tenosynovitis and underlying arthritis may be challenging on the basis of clinical examination (CE) alone. In daily clinical practice, ultrasound (US) imaging with Doppler information can be more versatile as it allows a dynamic assessment of joint inflammation and can be done quickly at the bedside. US can reliably evaluate the location of inflammation and provide direct visualization when local steroid injection is considered as treatment for children with JIA.

Objectives: To present the published data concerning the US assessment of tenosynovitis and the US metric properties investigated in JIA.

Methods: A systematic literature review (SLR) was conducted comprising PubMed, Embase and the Cochrane Library (before June 2018). Search terms: "arthritis, juvenile" AND, "tenosynovitis OR tendinopathy" AND "ultrasonography" OR "Doppler" AND "diagnosis OR therapy management". Selection criteria: Original articles published in English reporting US, tenosynovitis and tendinopathy in the paediatric population (up to 18 years old) with JIA. Data were extracted from manuscripts meeting the selection criteria, with particular focus on the definition used, scoring system applied and the US metric properties investigated. The quality of the studies was assessed using the same methodology developed for the SLR on paediatric synovitis [1].

Results: After selection, 11 articles were analysed. The tibialis posterior tendon group and peroneal tendons were the most commonly evaluated. Most authors applied the OMERACT definition proposed to rheumatoid arthritis (RA) [2] for evaluating paediatric tenosynovitis. Binary scoring

system on both grey scale and Doppler was applied in most of studies. Doppler function was commonly included, with only five specifying the Doppler parameters that were used. Construct validity was reported in roughly 50% of articles, including the CE as the main comparator. However, some of those articles have shown that clinical evaluation underestimates tendon involvement. US reliability and responsiveness for tenosynovitis in JIA was rarely included.

Conclusion: Although US seems a promising image modality to assess tenosynovitis in JIA, this SLR emphasizes the definitions of US elementary lesions of RA are frequently used in children. Moreover, there is insufficient evidence of studies exploring the construct validity of the US-determined tenosynovitis in JIA. Further validation studies are needed for implementation in clinical practice and trials.

REFERENCES:

- [1] Collado P. *Arthritis Care Res* 2012; 64:1011
- [2] Wakefield R. *J Rheumatol.* 2005;32:2485

Disclosure of Interests: PAZ COLLADO: None declared, Silvia Magni-Manzoni Consultant for: Abbvie, Speakers bureau: Abbvie, MARTINA STEINER: None declared, Tracy Ting : None declared, Patricia Vega Fernandez: None declared, Clara Malattia: None declared, Ana Rodriguez: None declared, George Bruyn : None declared, Helen Keen: None declared, Lene Terslev Speakers bureau: Speakers fee from : Roche, Novartis, Pfizer, MSD, BMS, Celgene

DOI: 10.1136/annrheumdis-2019-eular.3493

FRI0636

SUSCEPTIBILITY WEIGHTED SEQUENCES IN MAGNETIC RESONANCE IMAGING CAN CREATE COMPUTED TOMOGRAPHY-LIKE IMAGES AND IMPROVE THE ACCURACY OF STRUCTURAL LESION DETECTION OF THE SACROILIAC JOINT IN AXIAL SPONDYLOARTHRITIS

Torsten Diekhoff¹, Kay-Geert Hermann¹, Fabian Proff², Mikhail Protopopov², Denis Poddubnyy^{2,3}, Marcus R. Makowski², Arthritis Imaging Research Group - Berlin. ¹Charité – Universitätsmedizin Berlin, Department of Radiology, CCM, Berlin, Germany; ²Charité – Universitätsmedizin Berlin, Department of Rheumatology, CCM, Berlin, Germany; ³German Rheumatism Research Centre, Berlin, Germany

Background: In patients with axial spondyloarthritis (axSpA) the accurate depiction of structural lesions in magnetic resonance imaging (MRI) is impeded because MRI – while gaining signal from the bone marrow – allows only an indirect depiction of the cortical bone and sclerotic areas. Computed tomography (CT) is considered to be the gold standard as it displays the calcium-containing cortical and trabecular bone substance. However, the exhibited radiation exposure in CT limits its application in the clinical routine. Susceptibility weighted imaging (SWI) is an advanced MRI technique, that is able to visualise calcium due to its magnetic properties and allows a reconstruction of CT-like images without radiation exposure. (1)

Objectives: To investigate the diagnostic accuracy of CT-like images generated from SWI for the detection of structural lesions of the SI-joint.

Methods: Twentytwo patients with suspicion of or known axSpA were included. All patients underwent a 1.5-Tesla-MRI including T1 and SWI sequences and a low-dose CT of the sacroiliac joints. CT images were reconstructed in 4 mm oblique coronal reconstructions matching the orientation and slice thickness of the MRI. MRI and CT images were scored for erosions, sclerosis and joint space alterations applying the 24 regions method, a modification of the Berlin score. (2) Using CT as standard of reference, sensitivity (SE) and specificity (SP) values were calculated on patients' level. A score of 2 or higher in any quadrant was considered positive. The sumscores of T1 and SWI were compared to CT using Pearson's test.

Results: The mean erosion sumscore was 5.5 ± 5.5 , 6.3 ± 8.2 and 5.9 ± 9.2 for T1, SWI and CT respectively. The mean sclerosis sumscore was 0.78 ± 1.5 , 5.5 ± 6.5 and 6.9 ± 6.9 . Five patients had joint space alterations (pseudowidening or ankylosis) in CT. Both, T1 and SWI showed an SE of 100% and a SP of 84.5% for erosions, however, the erosion sumscore correlated better for SWI (Pearson's r : 0.93) than for T1 (r : 0.64). With respect to sclerosis T1 (SE 30%, SP 100%, r 0.42) was inferior to SWI (SE 90%, SP 92%, r 0.81). It was also less