conditions that are associated with a high risk of cardiovascular (CV) complications further beyond the increased prevalence of classical risk factors in this population. Consequently, the early identification of patients with MS-PSO and PsA who may benefit from primary CV prevention is essential. However, medical providers do not routinely screen or counsel their patients for CV risk factors or comorbidities. In addition, paucity of robust clinical data complicates the implementation of management protocols for primary CV prevention in MS-PSO and PsA.

Objectives: To generate expert-based recommendations on the management of CV comorbidities in patients with MS-PSO and PsA through the application of the Delphi technique.

Methods: The development of Delphi-based recommendations included four steps: (1) systematic literature review to explore current information and clinical recommendations on management of CV comorbidities in patients with PsO and PsA; (2) two subsequent meetings with project coordinators ($n = 3$) and scientific advisors ($n = 7$) from different medical specialties (rheumatology, dermatology, endocrinology, internal medicine, pneumology, rehabilitation, psychiatry and primary care) to examine the results of the review and decide which statements should be included in the final version of the questionnaire; (3) a two-round Delphi questionnaire aimed at specialists in rheumatology and dermatology to assess the degree of consensus on the different points raised; (4) finally, a face-to-face meeting with the project coordinators and scientific advisors to discuss and ratify the results of the Delphi process.

Results: As a result of the literature review and the assessment and approval of scientific advisors, the final version of the Delphi survey contains 52 statements grouped into 35 questions. The questionnaire is divided into 3 topic sections: (1) identification, diagnosis and referral of MS-PSO and/or PsA patients with CV comorbidities ($n = 12$ statements); (2) management and treatment of CV comorbidities associated with MS-PSO and/or PsA ($n = 36$); (3) influence of MS-PSO and/or PsA treatments on CV comorbidities and vice versa ($n = 4$).

Conclusion: This Delphi-based study may fill an important gap in the evidence by providing relevant information on the management of CV comorbidities in patients with MS-PSO and/or PsA.

Disclosure of Interests: : RAQUEL ALMODOVAR: None declared, Jose Luis López Estebaranz Consultant for: I have received fees as an advisory board member for Pfizer, Novartis, Leo Pharma, Lilly, Celgene, Abbvie and Janssen cilad, Speakers bureau: I have received fees as a speaker for Pfizer, Novartis, Leo Pharma, Lilly, Celgene, Abbvie and Janssen cilad, Eva De Higes Martínez : None declared, María Jesús López Navas : None declared, Carlos Guijarro Herraiz Consultant for: I have received payments as an advisor for Amgen, Esteve, Ferrer, MSD, Pfizer, Rubió and Sanofi., Speakers bureau: I have received payments as a speaker for Amgen, Esteve, Ferrer, MSD, Pfizer, Rubió and Sanofi., Juan José Gorgojo Martínez : None declared, David Palacios : None declared, José Carlos Peláez Álvarez Consultant for: I have received payments for consultancies from Shire Pharmaceuticals Ibérica, Pfizer, Merck Sharp Dohme España and Laboratorios Farmaceuticos Rovi., Speakers bureau: I have received lecture fees from Shire Pharmaceuticals Ibérica, Pfizer, Merck Sharp Dohme España and Laboratorios Farmaceuticos Rovi., Diana Ruiz Genao Consultant for: I have received payments for consultancies from Janssen-Cilag, Pfizer, Abbvie, Celgene, Leo Pharma, Novartis, Speakers bureau: I have received lecture fees from Janssen-Cilag, Pfizer, Abbvie, Celgene, Leo Pharma, Novartis., Lucía Pérez Carbonell : None declared, Susana Gómez Employee of: I am a current employee of Pfizer., Erica Falkelbach Employee of: I am a current employee of Pfizer., Francisco Rebollo Employee of: I am a current employee of Pfizer., Pedro Zarco Montejo : None declared
Doi: 10.1136/annrheumdis-2019-eular.6988

FRI0700-HPR LESSONS LEARNED FROM PILOT IMPLEMENTATION OF PHYSICAL ACTIVITY RECOMMENDATIONS IN AXSPA EXERCISE GROUP THERAPY – LESS MAY BE MORE

Anne-Kathrin Rausch Osthoff^{1,2}, Thea Vliet Vlieland², René Braem³, Béatrice Walker², Karin Niedermann Schneider¹. ¹Zurich University of Applied Sciences, Institute for Physiotherapy, Health, 8401 Winterthur, Switzerland; ²Leiden University Medical Center, Department of Orthopaedics, Rehabilitation and Physical Therapy, Leiden, Netherlands; ³Ankylosing Spondylitis Association of Switzerland, Zurich, Switzerland

Background: The Ankylosing Spondylitis Association of Switzerland (SVMB) offers 68 exercise groups (EG) for people with axial Spondyloarthritis (axSpA). Participants exercise weekly with focus on strength and flexibility supervised by a physiotherapist (PT). Given the EULAR physical

activity (PA) recommendations [1] and recent research [2, 3], the SVMB aims to further develop the concept of their EG by introducing: a) exercises in all dimensions, cardiorespiratory fitness, muscle strength, flexibility and neuromotor performance, according to the general PA recommendations [4]; b) individual exercise-coaching (three sessions in first six months) by the group PT, specifically trained in communication and action planning skills to support individual exercising, and c) bi-annual fitness-assessments for all exercise dimensions.

Objectives: To evaluate the new EG concept among the participating people with axSpA and PTs across four pilot groups.

Methods: After six months, a survey on the perception and satisfaction towards the new EG concept was sent to 30 people with axSpA, who rated the questions on a 0-10 numeric rating scale (0=not satisfied, 10=very satisfied). Semi-structured telephone-interviews were conducted with the PTs to explore their experiences with the new GE concept.

Results:

People with axSpA: The survey's response rate was 83% (25/30); of these, 15 (62%) participants were satisfied with the concept (mean 6.7 $\pm$ 2.23) and 20 (80%) with the exercise-coaching (mean 7.7 $\pm$ 2.92). 22 (88%) individuals defined a training goal and 15 (60%) achieved it. The exercise diary was used by 16 (64%) participants, five reported app-based. Reasons for not using the diary were the daily effort and problems with correctly allocating activities to exercise dimensions. A minority ($n=7$, 28%) reported to use a technical device for cardiovascular exercising. 17 (68%) were satisfied with the fitness-assessment (mean 7.5 $\pm$ 1.78). PTs perceived the exercise-coaching as very useful for supporting individualised exercising. However, they emphasised that the frequency of the sessions should be flexibly based on the participant's individual training goal and motivation. The PTs perceived controversial reactions to the fitness-assessments: increased but also decreased motivation towards exercising.

Conclusion: Exercise-coaching and fitness-assessments were generally appreciated; however, the frequency was perceived as too high. The GE concept will be adapted accordingly to improve feasibility for national implementation: the exercise-coachings will be reduced to two in the first six months, the first one compulsory for all participants, then bi-annually, according to individual needs and training goals. The fitness-assessments will take place annually instead of bi-annually. There is a need for easy-to-use exercise diaries, covering all exercise dimensions.

REFERENCES:

- [1] Rausch AK, et al. Ann Rheum Dis 2018; 9(77):1251-1260.
- [2] Sveaas SH, et al. Br J Sports Med 2017;51:1065-1072
- [3] Rausch AK, et al. RMD open, accepted
- [4] Garber CE, et al. Med Sci Sports Exerc 2011;43:1334-59.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7188

FRI0701-HPR EFFICACY OF INTERMITTENT AEROBIC EXERCISE ON SLEEP EFFICIENCY IN PATIENTS WITH RHEUMATOID ARTHRITIS. A RANDOMIZED CONTROLLED TRIAL

Katrine Løppenthin¹, Julie Midtgaard^{2,3}, Bente Appel Esbensen^{1,4}, Mikkel Østergaard^{1,4}, Jesper Frank Christensen⁵, Tanja Thomsen¹, Anders Tolver⁶, Poul Jennum^{4,7}. ¹Rigshospitalet, Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Centre of Head and Orthopaedics, Glostrup, Denmark; ²Rigshospitalet, University Hospitals Centre for Health Care Research (UCSF), Copenhagen University Hospital, Copenhagen, Denmark; ³University of Copenhagen, Department of Public Health, Faculty of Health and Medical Sciences, Copenhagen, Denmark; ⁴University of Copenhagen, Department of Clinical Medicine, Copenhagen, Denmark; ⁵Rigshospitalet, Copenhagen, Centre for Inflammation and Metabolism/Centre for Physical Activity Research, Copenhagen, Denmark; ⁶University of Copenhagen, Department of Mathematical Sciences Faculty of Science, Copenhagen, Denmark; ⁷Rigshospitalet, Danish Center for Sleep Medicine, Department of Clinical Neurophysiology, Glostrup, Denmark

Background: Insomnia is seen in >60% of patients with rheumatoid arthritis (RA). While cross-sectional studies have shown a positive association between regular physical activity and good sleep, there is a lack of knowledge on the effect of physical activity on insomnia in patients with RA.

Objectives: To examine the efficacy of intermittent aerobic exercise on sleep efficiency as well as the effect on fatigue and depressive symptoms.