responders (15%) to various biologics (anti-TNFs (68%), Rituximab (16%), Tocilizumab (13%), Abatacept (3%)). PBMCs were left unstimulated or stimulated (anti-CD3/CD28 beads or 10 ng/mL LPS) and stained with three flow cytometry panels including specific surface, intracellular and intranuclear markers to deeply characterise the function of several subsets of monocytes, NK cells, T cells and B cells (e.g. disease-associated cell populations such as PD-1^{hi}CXCR5⁺ peripheral T helpers (Rao DA *Nature* 2017), CD27⁺HLA-DR⁺ effector CD4⁺ T cells (Fonseka CY *Sci Transl Med* 2018) and cytotoxic PD-1⁺CXCR5⁺ CD8⁺ T cell subsets, Tbet⁺CD11c⁺ autoimmune-associated B cells, CD14⁺IL1β⁺ pro-inflammatory monocytes (Zhang F *BioRxiv* 2018)). Hypothesis-free clustering algorithms were also used for data analysis to identify unreported cell populations. Statistical association testing was performed using mixed effects ordinal (EULAR response) or linear (DAS28 or its components) regression models.

Results: Preliminary interim analysis identified associations between pan-biologic non-response and spontaneous expression of pro-inflammatory cytokines (Th1, Th17 lineage) by CD4⁺ T cells ($p < 0.003$). Good response to all biologic drugs was independently associated with the number of CD56^{dim}CD16⁺ NK cells ($p < 0.001$). Good response to anti-TNFs was associated with TNFα, IFNγ and perforin producing CD4⁺ T cells after *in vitro* stimulation with anti-CD3/CD28 ($p < 0.02$).

Conclusion: Ongoing work involves the replication of these results. The identification of immune cell types associated with response to a particular class of biologic drugs (e.g. anti-TNFs, but not other classes) might inform the choice of first line biologic drugs based on individual patients' immune profiles (stratification to treatment response categories). Cells or cytokines generally associated with non-response to all biologics could represent new therapeutic targets, at least in some patients groups.

Acknowledgement: AMP RA/SLE

Disclosure of Interests: Gemma Radley: None declared, Ben Mulhearn: None declared, Laura Donlin: None declared, Jennifer H. Anolik: None declared, Michael Brenner Grant/research support from: Roche: sponsored research agreement on stromal cells (but has nothing to do with checkpoint related disease), Consultant for: GSK: consultant. (I am part of their immunology network, a group of about 8 immunologists who advise them regularly and broadly in the areas of inflammation and infection)., Soumya Raychaudhuri: None declared, Kimme Hyrich Grant/research support from: Grants to institution: BMS, Pfizer, UCB, Ann Morgan: None declared, Gerry Wilson: None declared, John Isaacs Grant/research support from: Pfizer, Grant/research support from: Pfizer, Consultant for: Abbvie, Pfizer, Roche, Galvani, Merck, Gilead, Eli Lilly, Amgen, Janssen, Celltrion, NAPP, Consultant for: Abbvie, Pfizer, Roche, Galvani, Merck, Gilead, Eli Lilly, Amgen, Janssen, Celltrion, NAPP, Speakers bureau: Abbvie, Pfizer, Eli Lilly, Speakers bureau: Abbvie, Pfizer, Eli Lilly, Tracy Hussell: None declared, Anne Barton: None declared, Sebastien Viatte: None declared

DOI: 10.1136/annrheumdis-2019-eular.2371

SAT0063

ASSOCIATION OF SMOKING WITH TRIPLE CONCORDANT SEROPOSITIVITY IN RA PATIENTS, AND WITH RHEUMATOID FACTOR IN THE REMAINING PATIENTS

Cristina Regueiro¹, Luis Rodriguez Rodriguez², Raquel López-Mejías³, Laura Nuño⁴, Ana Triguero-Martínez⁵, Eva Perez-Pampín¹, Alfonso Corrales³, Alejandro Villalva⁴, Yolanda Lopez-Golan¹, Lydia Abasolo², Sara Remuzgo Martínez³, Ana Ortiz², Eva Herranz², Ana Martínez-Feito⁴, Carmen Conde¹, Antonio Mera Varela¹, Alejandro Balsa⁴, Isidoro González-Álvarez⁵, Miguel A. González-Gay³, Benjamin Fernandez², Antonio Gonzalez¹.

¹Instituto de Investigación Sanitaria – Hospital Clínico Universitario de Santiago, Santiago de Compostela, Spain; ²Instituto de Investigación Sanitaria San Carlos, Madrid, Spain; ³Hospital de Investigación Sanitaria Valdecilla, Santander, Spain; ⁴Instituto de Investigación del Hospital Universitario La Paz, Madrid, Spain; ⁵Instituto de Investigación del Hospital de La Princesa, Madrid, Spain

Background: The contribution of cigarette smoking to the risk of rheumatoid arthritis (RA) is larger for seropositive than for seronegative patients. Current pathogenic models explain this difference via the production of anti-citrullinated protein antibodies (ACPA) induced by protein citrullination in the lungs as a consequence of smoking. However, there are inconsistent results with this model including the van Wesemael *et al.* report, which found an association of smoking with the concurrent presence of ACPA, rheumatoid factor (RF) and anti-carbamylated protein antibodies (ACarPA), rather than with ACPA.

Objectives: We aimed to explore the relationship between smoking and the presence of FR, ACPA and ACaPA and their combinations in patients with RA.

Methods: Six cohorts with 2253 RA patients were obtained to replicate the van Wesemael *et al.* results. After replication, they were combined

with the three cohorts from that study comprising 2238 patients to a total of 4491 patients. The results from the different cohorts were combined by fixed-effects meta-analysis. Most analyses involved comparisons between two levels of smoking exposure, never and ever smokers, considering the RF/ACPA/ACaP patients as the reference.

Results: Analysis of the replication set confirmed the predominant association of smoking with the concurrent presence of the three RA autoantibodies (OR = 1.99, $P = 2.5 \times 10^{-6}$), whereas the patients with one or two antibodies were not different from the patients without antibodies (OR = 1.22, $p = 0.4$ and OR = 1.22, $p = 0.3$, respectively). The combined meta-analysis with all the cohorts clarified the dominant association of smoking with the triple concordant seropositivity (OR = 2.00, $p = 4.4 \times 10^{-16}$). In contrast, there was no association with the patient bearing one autoantibody (OR = 1.12, $p = 0.4$), and an in-between association with the concurrent presence of two antibodies (OR = 1.26, $p = 0.009$). Moreover, the association with the presence of three antibodies was significantly larger than with two antibodies (OR = 1.54, $p = 1.4 \times 10^{-6}$), whereas the association with the presence of two antibodies was not larger than with one autoantibody (OR = 1.11, $p = 0.3$). In the patients remaining after exclusion of the triple seropositive, the smokers were exclusively associated with the RF⁺ (OR = 1.28, $p = 0.03$ and OR = 1.30, $p = 0.004$ in the double and single positive patients, respectively). This association was independent of the reference, either the triple seronegative (OR = 1.29, $p = 0.001$) or all the RF⁺ patients (OR = 1.32, $p = 0.0002$).

Conclusion: Smoking increases RA susceptibility by promoting pathways leading to the concurrent presence of the three RA autoantibodies and, in its defect, to the production of RF. These actions are not covered by current pathogenic models and suggest that smoking accelerates epitope spreading.

REFERENCE

[1] van Wesemael, et al. Arthritis Res Ther. 2016;18:285.

Acknowledgement: Supported by grants PI17/01606 and RD16/0012/0014 of the Instituto de Salud Carlos III (Spain) that are partially financed by FEDER.

Disclosure of Interests: Cristina Regueiro : None declared, Luis Rodriguez Rodriguez: None declared, Raquel López-Mejías: None declared, Laura Nuño: None declared, Ana Triguero-Martínez: None declared, Eva Perez-Pampín: None declared, Alfonso Corrales: None declared, Alejandro Villalva: None declared, Yolanda Lopez-Golan: None declared, Lydia Abasolo: None declared, Sara Remuzgo Martínez: None declared, Ana Ortiz: None declared, Eva Herranz: None declared, ANA MARTÍNEZ-FEITO: None declared, Carmen Conde : None declared, ANTONIO MERA VARELA: None declared, Alejandro Balsa Grant/research support from: Abbvie, Pfizer, Novartis, BMS, Nordic, Sanofi, Consultant for: Abbvie, Pfizer, Novartis, BMS, Nordic, Sanofi, Sandoz, Lilly, Paid instructor for: Pfizer, Speakers bureau: Pfizer, Novartis, UCB, Nordic, Sanofi, Sandoz, Lilly, Isidoro González-Álvarez: None declared, Miguel A González-Gay Grant/research support from: Prof. MA Gonzalez-Gay received grants/research supports from Abbvie, MSD, Jansen and Roche., Speakers bureau: Consultation fees/participation in company sponsored speaker's bureau from Pfizer, Lilly, Sobi, Celgene, Novartis, Roche and Sanofi., Benjamin Fernandez: None declared, Antonio Gonzalez: None declared

DOI: 10.1136/annrheumdis-2019-eular.4607

SAT0064

EXPANSION OF CD16+ MONOCYTE SUBSETS AND ACE EXPRESSION ARE ASSOCIATED WITH ARTERIAL THICKENING AND VASCULAR FUNCTION IN VERY EARLY RHEUMATOID ARTHRITIS

Javier Rodríguez-Carrio^{1,2}, Mercedes Alperi-López³, Patricia López¹, Ángel Pérez-Álvarez⁴, Lorena Benavente⁴, Francisco Javier Ballina-García³, Ana Suárez¹.

¹University of Oviedo, ISPA, Area of Immunology, Oviedo, Spain; ²Hospital Universitario Central de Asturias, Bone and Mineral Research Unit, Instituto Reina Sofía de Investigación Nefrológica, REDinREN del ISCIII, Oviedo, Spain; ³Hospital Universitario Central de Asturias, Department of Rheumatology, Oviedo, Spain; ⁴Hospital Universitario Central de Asturias, Department of Neurology, Oviedo, Spain

Background: monocytes are largely recognized as drivers of the atherosclerosis development, the leading determinant of cardiovascular (CV) morbidity in rheumatoid arthritis (RA). Far from being a unique population, functionally and phenotypically subpopulations can be distinguished, although their relevance to CV disease is uncertain. Recently, the expression of angiotensin-converting enzyme (ACE) expression has been