seems to suppress Th17 and osteoclasts differentiation in RA patients in synergy with tocilizumab.

REFERENCE

- [1] Bellan M. Role of Vitamin D in Rheumatoid Arthritis. *Adv Exp Med Biol.* 2017;996:155-168

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3857

SAT0154

STABILIZATION OF RHEUMATOID ARTHRITIS ASSOCIATED INTERSTITIAL LUNG DISEASE BY RITUXIMAB: A SINGLE CENTER EXPERIENCE WITH LONG-TERM FOLLOW-UP

Nikolaos Kougkas, Nestor Avgustidis, Anastasios Eskitzis, Styliani Polia, Ioannis Papalopoulos, Irini Flouri, Argyro Repa, Ainour Molla Ismail Sali, George Bertias, Prodromos Sidiropoulos. *University Hospital of Heraklion, Crete, Greece, Rheumatology, Clinical Immunology and Allergy, Heraklion, Greece*

Background: Interstitial lung disease (ILD) is the most significant pulmonary manifestation of Rheumatoid Arthritis (RA)[1], contributing to decreased quality of life, progressive chronic disability and high utilization of healthcare resources. Treatment is challenging with mixed results from different biologics while Rituximab (RTX) has the most encouraging data based on retrospective observational studies. [2]

Objectives: To evaluate the efficacy and safety of RTX in patients with RA-ILD in the context of daily practice.

Methods: Observational retrospective study (2009-2019) of patients with a diagnosis of RA-ILD from a cohort of RTX-treated RA patients in the University Hospital of Heraklion. ILD was diagnosed by HRCT and monitored by means of pulmonary function tests (PFTs), HRCT and 6-minute walking test (6MWT, clinically important difference >50 m). RTX was used at standard dose (1 gr x2/6 months). ILD progression was defined by any of: decrease of pre-RTX forced vital capacity (FVC) >10%, or diffusion capacity of carbon monoxide (DLCO) >15% predicted score, or death from progressive ILD. Improvement was defined by any of: increase of pre-RTX FVC >10% or DLCO >15% predicted. All other patients were classified as having stable disease. RA activity was assessed by DAS28 (ESR).

TABLE

Baseline	End of follow-up
6MWT, n (%)	
– Stable	10/12(83%)
– Improvement	0/12(0%)
– Worsening	2/12(17%)
FVC, n (%)	
– Stable	10/13(76%)
– Improvement	1/13(7.7%)
– Worsening	2/13(15.3%)
DLCO, n (%)	
– Stable	10/13(76%)
– Improvement	1/13(7.7%)
– Worsening	2/13(15.3%)
HRCT, n (%)	
– Stable	13/16(81%)
– Improvement	1/16(6.25%)
– Worsening	2/16(12.75%)
DAS 28 (ESR) (median (IQR))	
5.05(2-6.9)	2.99(2.9-5.5)

Results: Of 247 RA patients treated with RTX, 16 (11 women, 5 men) had RA-ILD (prevalence 6.5%). 12/16 (75%) had radiographic pattern of UIP and 4/16 (25%) had NSIP. 9/16 (56%) were positive for ACPA and

or RF. The median (IQR) age at RTX initiation was 66 (54-77) years. ILD duration prior to RTX was 5.2 (1-15) years. Number of prior biologics was 1 ± 4 and 11/16 (69%) have been treated with concomitant DMARDs. 1/16 (6.25%) had previously received Cyclophosphamide. Concomitant COPD was present in 3/16 (19%) of the patients. Follow-up while on RTX was 37 (6-136) months. The overall dose of RTX was 13 (2-40) g. Following RTX treatment, new-onset ILD was diagnosed in 3/247 patients (incidence=1.2%). During follow-up, data for ILD were available for 13/16 patients. Post-RTX, 10/13(76%) showed stabilization, 1/13 (7.7%) had improvement, while 2/13(15.3%) had progression of RA-ILD. 1/16(6.25%) switched to Cyclophosphamide. During follow-up period, 6 admissions were recorded due to infections (4 for community-acquired pneumonia and 2 for hospital-acquired pneumonia) and 1/16(6.25%) patient died after progression of ILD.

Conclusion: In this cohort of RA-ILD, the majority of patients treated with RTX showed stabilization of lung disease, while RA improved significantly, over a prolonged follow-up period. RTX appears to be an acceptable and safe therapeutic choice for patients with RA-ILD. Nevertheless, randomized studies are warranted to prove any clinical effectiveness of RTX in RA-ILD.

REFERENCES

- [1] P.Spagnolo, et al. *Arthritis & Rheumatology* 2018 Oct; 70(10):1544-1554.
[2] Md Yuzaiful Md Yusof, et al. *Rheumatology* 2017; 56:1348-1357

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6828

SAT0155

THE PRESENCE OF ANTI-NUCLEAR ANTIBODIES BEFORE ANTI-TNF THERAPY IS A RISK FACTOR FOR THE APPEARANCE OF ANTI-DRUG ANTIBODIES IN PATIENTS WITH RHEUMATOID ARTHRITIS

Shunichi Kumagai^{1,2}, Ayano Mori², Toshiharu Saito², Miho Takahashi², Katsuhiko Yoneda¹, Yasuhiro Noda¹, Soushi Takahashi¹, Miwa Nishida¹, Saori Hatachi¹, Goh Tsuji¹. ¹Shinko Hospital, Center for Rheumatic Diseases, Kobe, Japan; ²Shinko Hospital, Shinko Institute for Medical Research, Kobe, Japan

Background: TNF inhibitors (TNFi) revolutionized the treatment of rheumatoid arthritis (RA), but have often developed anti-drug antibodies (ADrA) and also induced other autoimmune pathology^{1, 2}.

Objectives: To determine whether presence of anti-nuclear antibodies (ANA) predict the development of ADrA and clinical efficacy in RA patients treated with TNFi, infliximab (IFX) or adalimumab (ADA).

Methods: A total of 92 RA patients, 38 cases treated with IFX and 54 cases with ADA, were prospectively enrolled. Observation period of IFX-treated group was 3 years, and that of ADA-treated group was 1 years. ANA were measured before and serially during the therapy. ANA were measured by indirect fluorescent assay using a Computer-aided microscope system with HEp20-10 cells. The concentrations of anti-IFX antibodies (HACA) and anti-ADA antibodies (AAA) were measured by radioimmunoassay at Sanquin Diagnostic Services, Netherlands. Clinical response to therapy was judged by EULAR response criteria based on Disease Activity Score (DAS) 28-CRP.

Results: Thirteen (34.2%) of 38 patients in IFX group and 18 (33.3%) of 54 patients in ADA group became positive for HACA and AAA, respectively, during the treatment. All of the ADrA-positive patients were ANA-positive before the therapy. In contrast, none became ADrA-positive after treatment in 15 RA patients negative for ANA before therapy (6 cases in IFX group and 9 cases in ADA group), and the positive rate for ADrA was significantly different between ANA-positive and negative patients ($p=0.0018$). In addition, the positive rate for HACA was significantly higher in patients positive for ANA titers of ≥ 640 ($p=0.034$), but that for AAA was not related to ANA titers. Next, ANA became positive in 4 of 6 ANA-negative patients and ANA titers increased in 21 of 32 ANA-positive patients after IFX treatment. HACA positive rate was significantly higher in patients with ANA titers of ≥ 640 after IFX treatment ($p=0.0045$). ANA titers of patients in ADA group were mostly unchanging after treatment, and were not related to the appearance of AAA.

Continuance rate of TNFi (IFX and ADA) treatment was significantly lower in ADRa-positive patients than in those negative ($p=0.0066$ and $p=0.0127$, respectively). In IFX group, patients with ANA titers of ≥ 160 before treatment and those with ANA titers of ≥ 320 after treatment positive showed worse EULAR treatment response ($p=0.037$ and $p=0.033$, respectively). In ADA group, 7 of 9 ANA-negative patients before treatment showed moderate or good EULAR response, but positive ANA both before and after treatment was not connected with the clinical response.

Conclusion: The presence of ANA before IFX or ADA is a risk factor for the appearance of ADRa, while ADRa did not appear in any patient negative for ANA before treatment. ANA of high titers before and after IFX treatment predicted existence of ADRa and possibly leading to the treatment failure.

REFERENCES

- [1] Kalden JR, Shulze-Koops: Nat Rev Rheumatol, 2017, 13: 707-18.
- [2] Takase K, et al: Ann Rheum Dis, 2014, 73: 1695-99.

Disclosure of Interests: Shunichi Kumagai Grant/research support from: Mitsubishi Tanabe Pharm, Eisai Co.Ltd., Ayano Mori: None declared, Toshiharu Saito: None declared, Miho Takahashi: None declared, Katsuhiko Yoneda: None declared, Yasuhiro Noda: None declared, Soshi Takahashi: None declared, Miwa Nishida: None declared, Saori Hatachi: None declared, Goh Tsuji: None declared
DOI: 10.1136/annrheumdis-2019-eular.4649

SAT0156

METHOTREXATE DISCONTINUATION AND DOSE DECREASES AFTER THERAPY WITH TOCILIZUMAB: RESULTS FROM THE CORRONA RHEUMATOID ARTHRITIS REGISTRY

Dimitrios A. Pappas^{1,2}, Taylor Blachley², Steve Zlotnick³, Jennie H. Best³, Kelechi Emeanuru², Joel Kremer^{2,4}. ¹Columbia University, New York, United States of America; ²Corrona, LLC, Southborough, United States of America; ³Genentech, Inc, South San Francisco, United States of America; ⁴Albany Medical College and The Center for Rheumatology, Albany, United States of America

Background: Methotrexate (MTX) is frequently prescribed with biologic disease-modifying antirheumatic drugs. Evidence has shown that tocilizumab (TCZ) monotherapy is effective in the treatment of patients with rheumatoid arthritis (RA).^{1,2} Similar outcomes have been shown in patients responding to TCZ combination therapy who discontinued MTX or remained on combination therapy.³

Objectives: To examine MTX discontinuation and dose decreases in patients with RA initiating TCZ and to describe disease activity outcomes in a real-world setting.

Methods: TCZ-naïve patients enrolled in the Corrona RA Registry who initiated TCZ and had a 6-month follow-up visit without discontinuation of TCZ were included. Patients were grouped by MTX dose at the time of TCZ initiation (≤ 10 mg, > 10 to ≤ 15 mg, > 15 to ≤ 20 mg, > 20 mg). The primary outcome was the proportion of patients with changes in MTX use at 6 months. Changes in disease activity (Clinical Disease Activity Index [CDAI]) and patient-reported outcomes (PROs) over the follow-up period are described.

Results: Of 444 eligible patients, 82.7% were female, and 83.7% were white, with a mean (SD) disease duration of 11.6 (9.3) years and a baseline CDAI score of 24.0 (15.4). The mean (SD) MTX dose at baseline was 17.7 (5.8) mg. Overall, a total of 139 patients (31.3%) discontinued or decreased MTX at 6 months (overall mean [SD] dose change, -3.0 [7.5] mg); across baseline MTX dose groups, the proportion of patients who discontinued or decreased MTX at 6 months ranged from 28.2% to 38.2%. Improvements in CDAI scores and PROs were observed at 6 months in all baseline MTX dose groups and in patients who discontinued, decreased, maintained, or increased MTX doses at 6 months (Table 1). Similar patterns and results were observed at 12 months (not shown).

Conclusion: A considerable proportion of patients initiating TCZ were able to discontinue or decrease the dose of MTX after TCZ initiation. Patients who were able to discontinue or decrease MTX experienced similar improvements in disease activity and functionality. Discontinuing or decreasing MTX may be an effective treatment strategy for patients initiating TCZ combination therapy.

REFERENCES

- [1] Bykerk VP, et al. Clin Rheumatol. 2015;34(3):563-571.
- [2] Jones G, et al. Ann Rheum Dis. 2010;69(1):88-96.
- [3] Kremer J, et al. Arthritis Rheumatol. 2018;70(8):1200-1208.

Table 1. Change in Disease Activity and PROs* at 6 Months in Patients With RA

	All Patients (N = 444)	Discontinued MTX (n = 72)	Decreased MTX Dose (n = 67)	No Change in MTX Dose (n = 269)	Increased MTX Dose (n = 36)
CDAI	-7.8 (14.3)	-5.5 (13.1)	-7.1 (13.6)	-8.4 (14.7)	-9.8 (15.2)
DAS28	-1.2 (1.7)	-1.0 (1.7)	-1.1 (1.5)	-1.3 (1.6)	-0.9 (2.4)
Tender joint count	-3.2 (7.9)	-1.3 (7.1)	-1.8 (7.5)	-3.8 (8.1)	-4.2 (7.6)
Swollen joint count	-2.6 (5.6)	-2.2 (5.2)	-2.2 (5.1)	-2.7 (5.8)	-3.1 (6.1)
Physician global assessment	-13.1 (23.1)	-13.3 (23.4)	-17.0 (20.7)	-11.8 (23.4)	-16.2 (24.0)
Patient global assessment	-7.6 (25.9)	-6.4 (25.9)	-13.5 (28.5)	-6.4 (25.0)	-7.3 (26.4)
mHAQ	-0.1 (0.4)	-0.1 (0.4)	-0.1 (0.4)	0 (0.4)	-0.2 (0.5)
Patient pain	-7.3 (24.4)	-6.2 (23.7)	-8.2 (24.8)	-7.0 (24.3)	-10.8 (26.2)
Patient fatigue	-6.3 (26.6)	-7.3 (26.5)	-8.6 (30.7)	-5.0 (26.3)	-10.3 (20.2)
Morning stiffness present, n (%)	-29 (7.1)	-6 (9.4)	-5 (8.2)	-17 (6.8)	-1 (3.0)
Morning stiffness time, h	-0.1 (3.7)	-0.7 (2.8)	0 (2.1)	-0.1 (3.8)	0.3 (6.1)

CDAI, Clinical Disease Activity Index; DAS28, Disease Activity Score in 28 joints; mHAQ, modified Health Assessment Questionnaire; MTX, methotrexate; PRO, patient-reported outcome; RA, rheumatoid arthritis.
* Changes in disease activity and PRO measures are presented as mean (SD) unless otherwise indicated.

Acknowledgement: This study was sponsored by Corrona, LLC, and the analysis was funded by Genentech, Inc. Access to study data was limited to Corrona, and Corrona statisticians completed all of the analyses; all authors contributed to the interpretation of the results. Corrona has been supported through contracted subscriptions in the last 2 years by AbbVie, Amgen, Boehringer Ingelheim, Bristol-Myers Squibb, Celgene, Crescendo, Eli Lilly and Company, Genentech, Gilead, GSK, Janssen, Merck, Momenta Pharmaceuticals, Novartis, Pfizer Inc, Regeneron, Roche, Sun, UCB and Valeant.

Disclosure of Interests: Dimitrios A Pappas Grant/research support from: AbbVie, Consultant for: AbbVie, Employee of: Corrona, Taylor Blachley Employee of: Corrona, LLC, Steve Zlotnick Shareholder of: Genentech, Employee of: Genentech, Jennie H. Best Shareholder of: Genentech, Employee of: Genentech, Kelechi Emeanuru Employee of: Corrona, LLC, Joel Kremer Shareholder of: Corrona, Consultant for: AbbVie, Amgen, Bristol-Myers Squibb, Genentech, GSK, Lilly, Pfizer, Regeneron and Sanofi, Employee of: Corrona

DOI: 10.1136/annrheumdis-2019-eular.4200

SAT0157

REAL-WORLD EVALUATION OF PERSISTENCE WITH EARLY-LINE ABATACEPT VERSUS TUMOR NECROSIS FACTOR-INHIBITORS FOR RHEUMATOID ARTHRITIS COMPLICATED BY POOR PROGNOSTIC FACTORS

Damemarie Paul¹, Irina Yermilov², Sarah Gibbs², Michael Broder². ¹Bristol-Myers Squibb, Lawrenceville, NJ, United States of America; ²Partnership for Health Analytic Research, LLC, Beverly Hills, CA, United States of America

Background: Abatacept reduces signs and symptoms, inhibits the progression of structural damage, and improves physical function in adult patients with moderately to severely active rheumatoid arthritis (RA). Real-world data on its use as an early-line biologic agent are limited.

Objectives: To assess 12-month treatment persistence in early-line abatacept versus tumor necrosis factor inhibitor (TNFi) treated patients with RA complicated by poor prognostic factors.

Methods: We performed a multicenter retrospective medical record review of adult RA patients with poor prognostic factors treated at 5 United States clinics located in the West, Midwest, and Southeast. Patients were treated with abatacept or TNFi as the first biologic treatment at the clinic (defined as early line). Poor prognostic factors included positive anti-cyclic citrullinated peptide antibodies, positive rheumatoid factor antibodies, increased C-reactive protein levels, elevated erythrocyte sedimentation rate levels, or presence of joint erosions. TNFis included adalimumab, etanercept, infliximab (and their biosimilars), certolizumab pegol, or golimumab. Data were collected from biologic treatment initiation (between 8/9/11 and 11/14/16) for ≥ 1 year. Patients with Crohn's disease, ankylosing spondylitis, ulcerative colitis, psoriatic arthritis, or anal fistula were excluded. Demographic, disease, and treatment information (start, stop, reason for discontinuation) was abstracted. Treatment persistence (continuation of index treatment with gap ≤ 60 days) at 12 months and time to discontinuation were reported. Multivariate logistic and Cox regressions were used to compare 12-month persistence and overall discontinuation rate between abatacept and TNFi, controlling for demographic and clinical characteristics (age, sex, Charlson comorbidity index, RA duration), baseline utilization, and clinic.

Results: Data on 209 patients (88 abatacept, 121 TNFi) were collected. Abatacept patients were older than TNFi patients (64.36 vs. 57.23 years, $p<0.001$). There were no significant differences in either gender (79.55% vs. 72.73% female, $p=0.257$) or duration of treatment at the clinic (5.79 vs. 5.10 years, $p=0.226$) in the abatacept and TNFi cohorts. Median time to discontinuation was 1,672 days for abatacept vs. 612 days for TNFi ($p=0.002$). At 12 months, 86.36% of abatacept patients were persistent