SAT0253

PROGNOSTIC VALUE OF CARDIAC MAGNETIC RESONANCE IN SYSTEMIC SCLEROSIS

Cosimo Bruni¹, Luna Gargani², Alessia Pepe³, Antonella Meloni³, Daniele De Marchi³, Giancarlo Todiere³, Fabio Fulgenzi⁴, Yacob Ghebru⁴, Serena Guiducci¹, Raffaele De Caterina⁵, Marco Matucci-Cerinic¹. ¹Careggi University Hospital, Firenze, Italy; ²CNR IFC Istituto di Fisiologia Clinica, Pisa, Italy; ³Fondazione Toscana "Gabriele Monasterio", Pisa, Italy; ⁴Università degli Studi "Gabriele d'Annunzio" – sede di Chieti, Chieti, Italy; ⁵Ospedale Santa Chiara, Pisa, Italy

Background: Cardiac involvement is frequent in patients with systemic sclerosis (SSc). It is often subclinical, but significantly affects the prognosis of the disease. Cardiovascular magnetic resonance (CMR) is the non-invasive gold standard to quantify biventricular functional parameters and to perform a myocardial tissue characterization.

Objectives: To evaluate the prognostic value of CMR for cardiac events in SSc.

Methods: Two hundred and seventy-three SSc patients with a thorough clinical assessment underwent a CMR exam using a 1.5 T GE scanner. We quantified biventricular function parameter by SSFP cine images, oedema by STIR T2 images, and macroscopic fibrosis by late gadolinium enhancement (LGE) technique. Patients were followed-up and cardiac events were recorded as new onset of pulmonary arterial hypertension (PAH), new onset of heart failure (HF), or at least sustained ventricular tachycardia.

Results: Mean follow-up was 23.9 ± 17.0 months. During the follow-up a total of 14 events occurred (3 new onset PAH, 5 new onset HF, 6 ventricular tachycardia). CMR predictors of cardiac events by univariate analysis were left and right ventricular ejection fractions, indexed left and right atrial areas, and LGE (see Table). Myocardial fibrosis by LGE was the only independent predictor at multivariate analysis (hazard ratio 3.175; 95% C.I. 1.021-9.870, see Figure).

	Univariate analysis		Multivariate analysis	
	HR (95% CI)	p	HR (95% CI)	p
Left Ventricular Ejection Fraction	0.904 (0.843 – 0.969)	0.004	0.935 (0.868 – 1.007)	0.075
Indexed Left Atrial Area	1.244 (1.002 – 1.546)	0.048		
Right Ventricular Ejection Fraction	0.926 (0.884 – 0.969)	0.001	0.958 (0.904 – 1.014)	0.139
Indexed Right Atrial Area	1.384 (1.059 – 1.809)	0.017		
Right Ventricular End-Diastolic Mass	1.039 (0.991 – 1.091)	0.115		
Right Ventricular End-Systolic Mass	1.036 (0.988 – 1.086)	0.148		
T2 ratio	0.247 (0.019 – 3.251)	0.288		
Late Gadolinium Enhancement	3.871 (1.289 – 11.623)	0.016	3.175 (1.021 – 9.870)	0.046

Conclusion: Cardiac magnetic resonance anatomical and functional parameters of both the left and right heart have significant prognostic value in patients with SSc.

Disclosure of Interests: Cosimo Bruni: None declared, Luna Gargani: None declared, Alessia Pepe: None declared, Antonella Meloni: None declared, Daniele De Marchi: None declared, Giancarlo Todiere: None declared, Fabio Fulgenzi: None declared, Yacob Ghebru: None declared, Serena Guiducci: None declared, Raffaele De Caterina: None declared, Marco Matucci-Cerinic Grant/research support from: Actelion, MSD, Pfizer,

BMS, Chemomab, Sanipedia, Speakers bureau: Actelion, BMS; MSD, Janssen

DOI: 10.1136/annrheumdis-2019-eular.5583

SAT0254

VASODILATOR THERAPY IN THE LONG TERM PREVENTION OF MYOCARDIAL MANIFESTATIONS IN SYSTEMIC SCLEROSIS (SSc): RESULTS FROM DESSCIPHER INCEPTION COHORT STUDY

Antonella Riccardi¹, Dorte Husher², Serena Fasano¹, Valentina Messiniti¹, Yannick Allanore³, Jérôme Avouac³, László Czirájk⁴, Veronika Lorand⁴, Francesco Del Galdo⁵, Giuseppina Abignano⁵, Christopher Denton⁶, Svetlana Nihtyanova⁷, Oliver Distler⁸, Britta Maurer⁹, Marco Matucci-Cerinic¹⁰, Serena Guiducci¹⁰, Gabriela Riemekasten¹¹, Elise Siegert¹¹, Ulrich Walker¹², Veronika Jaeger¹³, Marc Freix¹⁴, Ingo Helmut Turner¹⁴, Ulf Müller-Ladner¹⁴, Gabriele Valentini¹.

¹University of Campania, Precision Medicine, Naples, Italy; ²University Hospital Charité, Biometry and Clinical Epidemiology, Charité, Germany; ³Cochin Hospital, Rheumatology, Paris, France; ⁴University of Pécs, Rheumatology and Immunology, Pécs, Hungary; ⁵University of Leeds, Biomedical Research Centre, Leeds, United Kingdom; ⁶University College London, Centre for Rheumatology and Connective Tissue Disease, London, United Kingdom; ⁷University College London, Centre for Rheumatology and Connective Tissue Disease, London, United Kingdom; ⁸University Hospital Zurich, Rheumatology, Zurich, Switzerland; ⁹University Hospital Zurich, Rheumatology, Zurich, Switzerland; ¹⁰University of Florence, Experimental and Clinical Medicine, Florence, Italy; ¹¹University of Lübeck, Rheumatology and Clinical Immunology, Lübeck, Germany; ¹²University of Basel, Rheumatology, Basel, Switzerland; ¹³University of Basel, Epidemiology, Basel, Switzerland; ¹⁴University Giessen, Rheumatology and Clinical Immunology, Bad Nauheim, Germany

Background: Calcium channel blockers (CCB) and angiotensin converting enzyme inhibitors (ACEinh) are known to improve in the short term, cardiac perfusion and function in patients with SSc (Parks JL et al. Rheum Dis Clin North Am 2014). No study has been carried out so far to investigate the long-term effectiveness of these drugs.

Objectives: To address this topic and other aspects of the management of SSc, the DeSScipher (To decipher the optimal treatment of SSc) project was submitted to and financed by the European Community (FP7-HEALTH n°305495).

Methods: From Dec.1, 2012 to Nov. 30, 2015, 512 SSc patients who, as inclusion criterion, had no significant gut or lung or kidney involvement, were treated for 0.5-4 years (median 2.31) with either (n= 359: group 1) vasodilator treatment (i.e. CCB, or ACE-inh or Angiotensin II receptor blockers [AglIrb] or a combination of them i.e. CCB plus either ACEinh or AglIrb) or (n= 153: group 2) no vasodilator therapy. The 296 patients of the 2 groups, who had been assessed at baseline and along the follow-up for conduction blocks (CB), ventricular arrhythmias (VA), pacemaker implantation (PMI), left ventricular ejection fraction (LVEF) < 55% and congestive heart failure (CHF), and sudden cardiac death (SCD) during follow-up, were investigated for the cumulative incidence rate (IR) of all these events. The IR of single manifestations were calculated in the patients investigated for each of them. Cox regression analysis was carried out in the 296 patients with no missing value to identify independent predictors of the occurrence of any myocardial manifestation.

Results: During 1164 patient-years follow-up, 6/508 were implanted a PM, 10/506 developed a LVEF < 55%, 2/492 a CHF, 11/325 a VA, 36/305 a CB, none underwent a SCD, any of these manifestations intervening in 59 out of the 296 investigated for all the manifestations. The IR of PMI and VA was greater in group 2 patients as compared with group 1 (1.3 vs 0.2x100 pts/year; p=0.02) and (2.7 vs 1.1x100 pts/year; p=0.077). In stepwise Cox regression analysis, male gender (HR 2.44, 95%CI 1.3-4.6; p=0.005), age at enrollment (HR 2.97, 95%CI 1.7-5.2; p=0.0002) and vasodilator therapy (HR 0.41, 95%CI 0.2-0.7; p=0.002) were independent positive and respectively, negative predictors of new onset cardiac manifestations.

Conclusion: Long term vasodilator therapy is likely to have a preventative role on the occurrence of myocardial manifestations of SSc.

Disclosure of Interests: Antonella Riccardi: None declared, Dorte Husher: None declared, SERENA FASANO: None declared, Valentina Messiniti: None declared, Yannick Allanore Grant/research support from: Inventiva, F Hoffman La-Roche, Sanofi, BMS, Pfizer, Consultant for: Actelion, Bayer, BMS, Boehringer, Roche, Sanofi, Jérôme Avouac Grant/research support from: research grant from Pfizer, László Czirájk: None declared, Veronika Lorand: None declared, Francesco Del Galdo: None declared, Giuseppina Abignano: None declared, Christopher Denton Grant/research support from: GlaxoSmithKline, Inventiva, CSF Behring, Consultant for: Roche-Genentech, Actelion, GlaxoSmithKline, Sanofi Aventis, Inventiva, CSL Behring, Boehringer Ingelheim, Bayer, Svetlana Nihtyanova: None

declared, Oliver Distler Grant/research support from: Prof. Distler received research funding from Actelion, Bayer, Boehringer Ingelheim and Mitsubishi Tanabe to investigate potential treatments of scleroderma and its complications, Consultant for: Prof. Distler has/had consultancy relationship within the last 3 years with Actelion, AnaMar, Bayer, Boehringer Ingelheim, ChemomAb, espeRare foundation, Genentech/Roche, GSK, Inventiva, Italfarmaco, iQvia, Lilly, medac, MedImmune, Mitsubishi Tanabe Pharma, Pharmacyclics, Novartis, Pfizer, Sanofi, Serodapharm and UCB in the area of potential treatments of scleroderma and its complications. In addition, he had/had consultancy relationship within the last 3 years with A. Menarini, Amgen, Abbvie, GSK, Mepha, MSD, Pfizer and UCB in the field of arthritides and related disorders, Britta Maurer Grant/research support from: Grant/research support from AbbVie, Protogen, Novartis; congress support from MSD, Pfizer, Roche, and Actelion, Marco Matucci-Cerinic Grant/research support from: Actelion, MSD, Pfizer, BMS, Chemomab, Sanipedia, Speakers bureau: Actelion, BMS; MSD, Janssen, Serena Guiducci: None declared, Gabriela Riemekasten Consultant for: Chugai, F. Hoffmann-La Roche, Speakers bureau: Chugai, F. Hoffmann-La Roche, Elise Siegert Shareholder of: Astra Zeneca, Grant/research support from: Actelion, Consultant for: AEC Partners, Speakers bureau: Actelion, Norsk, Ulrich Walker Grant/research support from: Unrestricted grant from AbbVie for the creation of the CoMPASS II app, Veronika Jaeger Grant/research support from: Has received an unrestricted grant from AbbVie to support the creation of the CoMPASS II app, Marc Frerix: None declared, Ingo Helmut Tarnier: None declared, Ulf Müller-Ladner Grant/research support from: Projekt supported by an unrestricted educational grant from Celgene GmbH., Gabriele Valentini Grant/research support from: MSD, Pfizer, Consultant for: MSD, Pfizer, biogen, Speakers bureau: MSD, amgen, biogen, lilly, sanofi, pfizer
DOI: 10.1136/annrheumdis-2019-eular.4784

SAT0255 EVALUATION OF THE CARDIOVASCULAR INVOLVEMENT OF SYSTEMIC SCLEROSIS USING NON-INVASIVE CARDIAC IMAGING TECHNIQUES

Carina Espinet^{1,2}, Maria Nazarena Pizzi¹, Alfredo Guillén del Castillo¹, Albert Roque¹, Victor Pineda¹, Isidro Sanz¹, Eduardo L. Callejas-Moraga², Maria Roca¹, Santiago Aguade¹, Carmen Pilar Simeón-Aznar¹. ¹Hospital Universitari Vall Hebron, Barcelona, Spain; ²Parc Taulí Hospital, Badalona, Spain

Background: Systemic sclerosis (SSc) is a systemic disease that may affect many organs; among them, cardiac involvement. The prevalence of cardiac involvement in SSc varies depending on the sensitivity of the methods used for its detection. Indirect evidence suggests that subclinical cardiac involvement may eventually occur in the vast majority of patients with SSc. Early detection and monitoring of myocardial involvement are integral to SSc management, as cardiovascular involvement is known to be a poor prognostic indicator when present

Objectives: To describe myocardial perfusion abnormalities and potentially associated coronary arteries lesions using non-invasive imaging techniques in a group of patients with SSc and suggestive symptoms of myocardial involvement (symptomatic) in comparison with a control group of patients with SSc without cardiac symptoms

Methods: A retrospective observational study was performed including a total of 61 patients diagnosed with SSc, 52 symptomatic, with dyspnea and/or chest pain (57.98 ± 12.3 years, 45 women) and 9 asymptomatic controls (50.2 ± 15.21 years, 8 women).

All patients underwent a post-stress (treadmill or pharmacological) myocardial perfusion gated-SPECT, a cold-induced stress SPECT, that were compared to a rest SPECT (to assess ischemia and/or necrosis), as well as a cardiac CT-angiography (to assess significant coronary arteries lesions, considering stenosis of more than 50%)

Results: Twenty-one out of the 52 symptomatic patients (50%) showed myocardial perfusion defects in the stress-rest SPECT: 13 (25%) showed ischemia, 13 (25%) fibrosis/necrosis, and 5 (9.6%) ischemia and necrosis. In the cold-induced SPECT, 17 patients (32.7%) had myocardial perfusion abnormalities: 10 (19.2%) showed ischemia, 13 (25%) fibrosis/necrosis and 6 (11.5%) ischemia and necrosis.

In the other hand, of the 9 asymptomatic patients only 1 (11%) had ischemia and necrosis in the stress-rest SPECT, being only positive for necrosis in the cold-induced SPECT images.

In the cardiac CT-angiography, 7/52 patients (13.4%) showed significant coronary lesions, 4 (57.2%) of them with perfusion defects in the SPECT images, and 3 (42.8%) without significant perfusion alterations

Of the 9 asymptomatic patients, 1 (11%) had significant coronary lesions, being the same patient who presented perfusion defects in myocardial SPECT images

Conclusion: The gated-SPECT is a sensitive tool for detecting myocardial perfusion alterations, normally with no associated significant coronary lesions, suggesting microvascular abnormalities. In this cohort, myocardial perfusion abnormalities were detected in 50% of symptomatic patients, whereas in only 11% of non-symptomatic patients

REFERENCES

- [1] Fernández-Codina A, Simeón-Aznar CP2, Pinal-Fernandez I, Rodríguez-Palomares J, Pizzi MN, Hidalgo CE, Guillén-Del Castillo A, Prado-Galbarro FJ, Sarria-Santamera A, Fonollosa-Plà V, Vilardell-Tarrés M. **Cardiac involvement in systemic sclerosis: differences between clinical subsets and influence on survival.** *Rheumatol Int.* 2017 Jan;37(1):75-84. doi: 10.1007/s00296-015-3382-2. Epub 2015 Oct 25.
- [2] Rangarajan V, Matiasz R, Freed BH. Cardiac complications of systemic sclerosis and management: recent progress. *Curr Opin Rheumatol.* 2017 Nov;29(6):574-584. doi: 10.1097/BOR.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5492

SAT0256 EFFICACY AND SAFETY OF ANTIFIBROTIC THERAPY IN DIFFUSE INTERSTITIAL PULMONARY DISEASE ASSOCIATED WITH SYSTEMIC AUTOIMMUNE DISEASES

Marta Aguilar-Zamora¹, J. Narváez², Juanjo J. Alegre-Sancho¹, Vanesa Vicens³, Maria Molina³, Joan Miquel Nolla². ¹Hospital Universitari Dr. Peset, Rheumatology, Valencia, Spain; ²Hospital Universitari de Bellvitge, Rheumatology, Bellvitge, Spain; ³Hospital Universitari de Bellvitge, Pneumology, Bellvitge, Spain

Objectives: Assess the efficacy and safety of antifibrotic agents (pirfenidone and nintedanib) in refractory diffuse interstitial lung disease (DILD) associated with systemic autoimmune diseases (SAD).

Methods: Open observational study in patients with active symptomatic DILD-SAD (evidence of clinical and functional impairment) despite treatment with glucocorticoids (GC) and immunosuppressant therapy.

Results: We included 13 patients (8 women) with a mean age (± SD) of 55 ± 12 years (range, 27-71). The baseline SAD were: eighth systemic sclerosis, one rheumatoid arthritis, one ankylosing spondylitis, one microscopic polyangiitis with pulmonary fibrosis and one interstitial lung disease with autoimmune characteristics (IPAF). Mean time from diagnosis to initiation of antifibrotic therapy was 3 ± 8.2 years (IQR 25-75%: 1.6-9.5). The histopathological patterns found were: seven cases (54%) of usual interstitial pneumonia (UIP), three (23%) nonspecific interstitial pneumonia (NSIP), two cases (15%) combined pattern of pulmonary fibrosis plus emphysema (CPFE), and one (8%) non-classifiable pulmonary fibrosis (without histospecific radiological pattern).

In addition to GC, the previous therapy tested for DILD was mycophenolate mofetil (MMF) in 77% of the patients, cyclophosphamide (8 intravenous boluses) in 46%, and rituximab (RTX) in 54% (the number of RTX cycles administered were 4 ± 3.3; range, 1-9). Despite this, during the year prior to start the antifibrotic drugs, all patients presented a worsening in the values of % pFVC (-8.77%) and % pDLCO (-12.41%).

Five (38%) patients received pirfenidone at doses of 3 tablets/8 hours (2403 mg/day) and eight (62%) nintedanib (5 at a dose of 150 mg/ 12 hours and 3 at a dose of 100 mg/ 12 hours).

Antifibrotic therapy was administered in combination with MMF (1.5-2 g/day) in 10 patients (77%) and with rituximab in 3 (23%), in addition to GC at doses between 5 and 20 mg/ prednisone per day. The mean

	Pre-treatment (mean ± SD)	Post-treatment (mean ± SD)	Delta (mean)	p
%pFVC	67.9 ± 15.6	74.2 ± 13.9	+ 5.13	NS
%pTLC	66 ± 10.8	77 ± 15.55	+ 4.86	NS
%pDLCO	37.8 ± 7.22	44 ± 7.32	+ 4.42	NS
Walking (meters)	409 ± 52	415 ± 103	+ 7.57	NS

The response in the Pulmonary function testing (PFT) in these 9 patients categorized according to the definitions of the American Thoracic Society was as follows: A) % pFVC: stabilization in 5 cases and improvement in 4; B) % pDLCO: stabilization in 7 cases and improvement in 3.